

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Cancer-Associated Venous Thromboembolic Disease

Version 1.2016

NCCN.org

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2016 Panel Members

Cancer-Associated Venous Thromboembolic Disease

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

***Michael B. Streiff, MD/Chair ‡**
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Shuwei Gao, MD †
The University of Texas
MD Anderson Cancer Center

Thomas L. Ortel, MD, PhD ‡
Duke Cancer Institute

Bjorn Holmstrom, MD †/Vice-Chair
Moffitt Cancer Center

Samuel Z. Goldhaber, MD λ
Dana-Farber/Brigham and Women's
Cancer Center

Rita Paschal, MD ‡
University of Alabama at Birmingham
Comprehensive Cancer Center

Aneel Ashrani, MD ‡
Mayo Clinic Cancer Center

Paul Hendrie, MD, PhD ‡
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Sanford Shattil, MD ‡
UC San Diego Moores Cancer Center

Paula L. Bockenstedt, MD ‡
University of Michigan
Comprehensive Cancer Center

Nicole Kuderer, MD, MS ‡ †
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Tanya Siddiqi, MD ‡
City of Hope Comprehensive Cancer Center

Carolyn Chesney, MD ‡
St. Jude Children's Research Hospital/
The University of Tennessee Health
Science Center

Alfred Lee, MD, PhD ‡
Yale Cancer Center/Smilow Cancer Hospital

Kristi J. Smock, MD ‡ ≠
Huntsman Cancer Institute
at the University of Utah

***John Fanikos, RPH, MBA Σ**
Dana-Farber/Brigham and Women's
Cancer Center

Jason T. Lee, MD ¶ λ
Stanford Cancer Institute

Gerald Soff, MD ‡
Memorial Sloan Kettering Cancer Center

Randolph B. Fenninger, JD ≠
National Blood Clot Alliance

Mirjana Lovrinevic, MD
Roswell Park Cancer Institute

Tzu-Fei Wang, MD ‡
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Annemarie E. Fogerty, MD ‡ †
Massachusetts General Hospital
Cancer Center

Michael M. Millenson, MD ‡ † †
Fox Chase Cancer Center

Gary C. Yee, PharmD Σ ‡
Fred & Pamela Buffett Cancer Center

Lalitha Nayak, MD, DO, PhD, RN ‡
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center and
Cleveland Clinic Taussig Cancer Institute

Anaadriana Zakarija, MD ‡
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

NCCN
Anita Engh, PhD
Karin G. Hoffmann, RN, CCM

Continue

¶ Surgery/Surgical oncology
‡ Hematology/Hematology oncology
Σ Pharmacology/Pharmacy
† Medical oncology
λ Cardiology
† Internal medicine
Nursing
≠ Patient advocacy
≠ Pathology
*Discussion section writing member

[NCCN Guidelines Panel Disclosures](#)



[NCCN Cancer-Associated Venous Thromboembolic Disease Panel Members](#) [Summary of the Guidelines Updates](#)

[Inpatient-Venous Thromboembolism Prophylaxis \(VTE-1\)](#)

[Acute Superficial Vein Thrombosis \(SVT-1\)](#)

[Acute Deep Vein Thrombosis \(DVT-1\)](#)

[Acute Pulmonary Embolism \(PE-1\)](#)

[Heparin-Induced Thrombocytopenia \(HIT-1\)](#)

[Splanchnic Vein Thrombosis \(SPVT-1\)](#)

[VTE Risk Factors in Cancer Patients \(VTE-A\)](#)

[Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment and](#)

[Contraindications to Mechanical Prophylaxis \(VTE-B\)](#)

[Inpatient/Outpatient Prophylactic Anticoagulation Treatment \(VTE-C\)](#)

[Therapeutic Anticoagulation for Venous Thromboembolism \(VTE-D\)](#)

[Reversal of Anticoagulation \(VTE-E\)](#)

[Elements for Consideration in Decision Not to Treat \(VTE-F\)](#)

[Clinical Scenarios Warranting Consideration of Filter Placement \(VTE-G\)](#)

[Therapeutic Anticoagulation Failure \(VTE-H\)](#)

[Thrombolytic Agents \(VTE-I\)](#)

[Contraindications to Thrombolysis and Indications for Thrombolysis \(VTE-J\)](#)

[Perioperative Management of Anticoagulation and Antithrombotic Therapy \(PMA-1\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical_trials/physician.html](#).

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise specified.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2016.



NCCN Guidelines Version 1.2016 Updates

Cancer-Associated Venous Thromboembolic Disease

Updates in Version 1.2016 of the NCCN Guidelines for Cancer-Associated Venous Thromboembolic Disease from Version 1.2015 include:

General

- “Pomalidomide” was added to “thalidomide/lenalidomide” as appropriate throughout the guidelines.

Venous Thromboembolism

VTE-1

- At-Risk Population
 - ▶ 3rd bullet: Revised “Providers are encouraged to discuss VTE risk factors, *risks and benefits* of VTE prevention, and the importance of patient adherence to care programs.” (Also for for VTE-2)
- Footnote “c” was revised: ~~“Discuss prevention and risks/benefits of VTE prophylactic anticoagulation by pharmacologic intervention. Discuss VTE prevention and the risks/benefits of pharmacologic and mechanical VTE prophylaxis. A systematic approach to patient risk assessment is recommended. Institutions are strongly encouraged to implement best practice programs to monitor provider and patient adherence to VTE prophylaxis.”~~

VTE-2

- At-Risk Population
 - ▶ Recommendations for “Surgical oncology patient” were revised to: “Out-of-hospital primary VTE prophylaxis is recommended for up to four weeks post-operation (particularly for high-risk abdominal or pelvic cancer surgery patients.)”
 - ▶ For “Medical oncology patient” treated in “Other outpatient settings”, footnote was removed: “Consider patient conversation about risks and benefits of VTE prophylaxis in patients with a Khorana score ≥ 3 . (See Khorana Predictive Model [VTE-A 3 of 3]).”

Acute Deep Vein Thrombosis (DVT) and Acute Superficial Vein Thrombosis (SVT)

- Deep or Superficial Vein Thrombosis (DVT/SVT) was divided into two new pages.

SVT-1

- “Acute Superficial Vein Thrombosis (SVT)” was extensively revised.

DVT-1

- “Acute Deep Vein Thrombosis (DVT)” was extensively revised.

DVT-3

- Treatment
 - ▶ DTV, no contraindication, 1st bullet was revised: “Anticoagulation for at least 3 months or as long as catheter CVAD is in place if ~~catheter removed, total duration of therapy is at least 3 months~~”
 - ▶ No DVT, bullet was revised from “Consider further imaging/testing with another modality if clinical suspicion is high and initial imaging failed to show” to “Consider further diagnostic imaging/testing if initial testing is unrevealing and clinical suspicion remains high”
 - ▶ Footnote “k” is new: “Consider longer duration anticoagulation in patients with poor flow, persistent symptoms, or unresolved thrombus.”

Acute Pulmonary Embolism (PE)

PE-1

- Diagnosis and Evaluation
 - ▶ Under “Diagnosis”, a bullet was removed: “Incidental PE”
 - ▶ “Unsuspected PE” pathway is new, with the evaluation: “If not already performed: Comprehensive medical history, and physical examination; CBC with platelet count; PT, aPTT; Liver and kidney function tests; EKG,” then referred to: “See PE Treatment (PE-2)”
 - ▶ Clinical suspicion of PE, Imaging, 3rd bullet was revised: “... (if patient has renal insufficiency or ~~uncorrectable allergy to contrast allergy refractory to anaphylaxis prophylaxis~~)”
 - ▶ Footnote “c” was revised: “Repeat imaging and diagnostic studies are ~~is~~ not routinely needed in patients with incidental PE. Consider outpatient management for these patients.”

PE-2

- Treatment
 - ▶ Footnote was removed from risk stratification procedures: “The Pulmonary Embolism Severity Index (PESI) clinical prediction rule can also be considered, but should not be substituted for the risk stratification procedures indicated above. (Donze J, Le Gal G, Fine MJ, et al. Prospective validation of the Pulmonary Embolism Severity Scale. *Throm Haemost* 2008;100:943-948). Consideration can also be given to the RIETE Cancer Score (den Exter P, Gómez V, Jiménez D, et al. A clinical prognostic model for the identification of low-risk patients with acute symptomatic pulmonary embolism and active cancer. *Chest* 2013;143:138-145.)”

[Continued](#)

UPDATES
1 OF 3

NCCN Guidelines Version 1.2016 Updates

Cancer-Associated Venous Thromboembolic Disease

Updates in Version 1.2016 of the NCCN Guidelines for Cancer-Associated Venous Thromboembolic Disease from Version 1.2015 include:

Heparin-Induced Thrombocytopenia (HIT)

HIT-1

• Diagnosis and Treatment of HIT

- ▶ “Low” pathway: Fourth bullet removed: “Consider HIT antibody test (enzyme-linked immunosorbent assay [ELISA]) for select patients (See HIT antibody test results HIT-2)”
- ▶ “Moderate/High” pathway:
 - ◊ 1st bullet was revised from “Eliminate unfractionated and LMWH exposure from all sources, including treatment, prophylaxis, flush doses, and coated catheters” to “Eliminate UFH/LMWH exposure from all sources (treatment, prophylaxis, line flushes, coated catheters)”
 - ◊ 2nd bullet was revised from “For patients receiving warfarin, discontinue it and reverse with vitamin K” to “Discontinue and reverse warfarin (and other vitamin K antagonists) with vitamin K”
- ▶ Footnote “c” was revised: “A “low” pre-test probability score combined with a negative antibody test is useful in ruling out a diagnosis of HIT; a positive test increases the suspicion for HIT. ~~Sending for the HIT antibody test should be individualized and based on clinical judgment. Patients with 4T scores <4 are very unlikely to have HIT, so routine HIT antibody testing in these patients is probably not advisable. In non-cancer patients with 4T scores of 1–3, the risk of HIT is small but not zero, but this has not been validated in cancer patients. Based on clinical judgment HIT antibody testing may be warranted in select patients of concern.~~”
- ▶ Footnote “e” is new: “Data supporting use of fondaparinux is limited; however, among experienced clinicians it is commonly used for outpatient management in low-risk patients.”

Therapeutic Options for HIT

HIT-B 1 of 2

- Direct Thrombin Inhibitors (Preferred); Argatroban:
 - ▶ 1st sub-bullet was revised: “Normal liver function, non-ICU patient: 4–2 mcg/kg/min adjusted to aPTT ratio (first check in 4 hours)”
 - ▶ 2nd sub-bullet was revised: “Abnormal liver function (total bilirubin 1.8–3.6 mg/dL; aspartate transaminase/alanine transaminase [AST/ALT] 150–600 IU/L) or ICU, heart, or multi-organ failure patient: ~~0–25~~ 0.5 mcg/kg/min”
- Footnote “3” is new: “Prescribing information: Argatroban injection, for intravenous infusion only. 2016.”

HIT-B 2 of 2

- Indirect Factor Xa Inhibitor, fondaparinux, 1st sub-bullet was revised: “For patients with C_{cr} 30–50 mL/min (clearance reduced by ≥40%): ~~Use caution~~ Consider using a DTI”
- Footnote “6” was added: “Prescribing information: Fondaparinux sodium, solution for subcutaneous injection. 2009”

Splanchnic Vein Thrombosis (SPVT)

SPVT-1

- Footnote “a”, Risk factors relevant to cancer population for SPVT; 8th sub-bullet was revised: “Myeloproliferative ~~disorder (polycythemia vera, essential thrombocythemia)~~ neoplasms associated with the JAK2 V617F mutation (most common) or CALR mutation (rare)”

SPVT-2

- Footnote “d” was revised: “Duration of anticoagulation should be at least 6 months for triggered events (eg, postsurgical) and indefinite if active cancer, ~~thrombophilic state, or idiopathic thrombosis~~ persistent thrombophilic state or unprovoked thrombotic event.”

VTE Risk Factors In Cancer Patients

VTE-A 1 of 3

- High-risk outpatients on chemotherapy, based on combinations of the following risk factors; 5th bullet was revised: “Use of ~~erythropoietic-erythropoiesis-stimulating agents~~”

VTE-A 2 of 3

- Risk Factors, Myeloma Therapy:
 - ▶ 1st bullet was revised: “~~Thalidomide or lenalidomide~~ IMiD in combination with:”
 - ▶ Footnote “2” is new: “Immunomodulator drugs (IMiDs), including thalidomide, lenalidomide, and pomalidomide.”
- Footnote “1” was revised: “~~Reproduced~~ Adapted with permission from Nature Publishing Group. Palumbo A, Rajkumar SV, Dimopoulos MA et al, Prevention of thalidomide- and lenalidomide-associated thrombosis in myeloma. Leukemia 2008;22:414-423. Copyright 2008.
<http://www.nature.com/leu/journal/v22/n2/full/2405062a.htm>”

VTE-A 3 of 3

- Title was revised: “VTE Risk Factors in Cancer Outpatients.”

[Continued](#)



NCCN Guidelines Version 1.2016 Updates

Cancer-Associated Venous Thromboembolic Disease

Updates in Version 1.2016 of the NCCN Guidelines for Cancer-Associated Venous Thromboembolic Disease from Version 1.2015 include:

VTE-B

- **Contraindications To Prophylactic or Therapeutic Anticoagulation Treatment**
 - ▶ 8th bullet under “Relative” was added: *“Interventional spine and pain procedures”*
 - ▶ Footnote “1” was revised: “Refer to institutional-specific anesthesia practice guidelines, if available. Twice daily prophylactic dose UFH (5000 units every 12 hours) and once daily LMWH (eg, enoxaparin 40 mg once daily) may be used with neuraxial anesthesia. Twice daily prophylactic dose LMWH (eg, enoxaparin 30 mg every 12 h), prophylactic dose fondaparinux (2.5 mg daily), and therapeutic dose anticoagulation should be used with extreme caution with neuraxial anesthesia. The safety of thrice daily prophylactic dose UFH in conjunction with neuraxial anesthesia has not been established. (Horlocker TT, Wedel DJ, Rowlingson JC, et al. *Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Third Edition)*. *Reg Anesth Pain Med* 2010;35:64-101.)”

Therapeutic Anticoagulation for Venous Thromboembolism

VTE-D 1 of 2

- Footnote “2” was revised: “Follow institutional standard operating procedures (SOPs) for dosing schedules. If no SOPs then use the American College of Chest Physicians (ACCP) recommendations. (Kearon C, Kahn SR, Agnelli G, Goldhaber S, et al. *Antithrombotic therapy for venous thromboembolic disease: American College of Physicians Evidence-Based Clinical Practice Guidelines (8th edition)*. *Chest* 2008;133[suppl]:454S-545S. [www.chestjournal.org]) Garcia DA, Baglin TP, Weitz JI, Samama MM. *Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines*. *Chest* 2012; 141:e24S-43S.”

Reversal of Anticoagulation in the Event of Life-Threatening Bleeding or Emergent Surgery

VTE-E 1 of 9

- Statement was revised: “In the event of bleeding or the need for urgent/emergent invasive procedures, anticoagulant effect must be reversed promptly. *It is incumbent on the provider to keep in stock the recommended reversal agents for all anticoagulants included in this table. This includes: protamine; vitamin K oral (phytonadione), and IV solution; fresh frozen plasma (FFP); 4-factor prothrombin complex concentrate (4-factor PCC), 3-factor PCC, rhFVIIa, activated prothrombin complex concentrates (aPCC) (anti-inhibitor coagulant complex, vapor heated), desmopressin (DDAVP), idarucizumab, and oral charcoal.*”
- Low-molecular-weight heparin (LMWH) (Half-life 3–7 hours); 4th bullet was revised: “Administer protamine by slow IV infusion (≤ 5 mg/min) (no faster than 5 mg per min)”

VTE-E 6 of 9

- **Reversal of Anticoagulation for “Dabigatran”**
 - ▶ 2nd bullet is new: *“Administer idarucizumab, 5 g IV”*
 - ▶ 3rd bullet was revised: ~~“No specific antidote exists, but beneficial effects have been ascribed to the following~~ *For special situations with slow or incomplete clearance (eg, renal dysfunction or failure), consider adding to idarucizumab:”*
 - ▶ Statement was removed: “May be helpful based on in vitro or animal models aPCC (anti-inhibitor coagulant complex, vapor heated 25–50 units/kg IV) rhFVIIa 90 mcg/kg IV”
- **Precautions/Additional Considerations for “Dabigatran”**
 - ▶ 3rd bullet revised: “In patients with renal failure/severe renal insufficiency, dialysis may be the most helpful in addition to idarucizumab.”

VTE-E 8 of 9

- This is a new page with recommendations for reversal of edoxaban anticoagulation in the event of life-threatening bleeding or emergent surgery.

VTE-J

- A new box was added listing indications for thrombolysis:
 - ▶ *“Limb-threatening/life-threatening acute proximal DVT*
 - ▶ *Symptomatic ileal femoral thrombosis*
 - ▶ *Massive/life-threatening PE*
 - ▶ *Intestinal SPVT with high risk of ischemia”*

Perioperative Management of Anticoagulation and Antithrombotic Therapy

PMA-A 1 of 2

- **Types of Surgery or Procedures**
 - ▶ 8th bullet for high bleeding risk was added: *“Head and neck surgery”*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

INPATIENT VENOUS THROMBOEMBOLISM PROPHYLAXIS

AT-RISK POPULATION

- Adult medical or surgical patient
- Diagnosis of cancer or clinical suspicion of cancer^a
- Providers are encouraged to discuss VTE risk factors, risks and benefits of VTE prevention, and the importance of patient adherence to care programs

WORKUP

- Initial Workup:**
- History and physical
 - Complete blood count (CBC) with platelet count
 - Prothrombin time (PT)
 - Activated partial thromboplastin time (aPTT)
 - Liver and kidney function tests

Contraindication to anticoagulation^b



INITIAL PROPHYLAXIS

- Prophylactic anticoagulation therapy^c (category 1)**
(Consider preoperative dosing with unfractionated heparin [UFH] or low-molecular-weight heparin [LMWH] for high-risk surgery [eg, abdominal/pelvic] patients) ± Intermittent pneumatic compression (IPC) device^{d,e} ± Graduated compression stockings (GCS)^{d,e}
- Mechanical prophylaxis^e**
• IPC ± GCS^{d,e}

[VTE Prophylaxis following discharge \(VTE-2\)](#)

^aSee [VTE Risk Factors in Cancer Patients \(VTE-A\)](#).

^bSee [Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment \(VTE-B\)](#).

^cDiscuss VTE prevention and the risks/benefits of pharmacologic and mechanical VTE prophylaxis. A systematic approach to patient risk assessment is recommended. Institutions are strongly encouraged to implement best practice programs to monitor provider and patient adherence to VTE prophylaxis. See [Inpatient/Outpatient Prophylactic Anticoagulation Treatment \(VTE-C\)](#).

^dIPC device is preferred over GCS. Patients should be appropriately measured for stockings and monitored for adverse effects including skin ulcerations, especially in immobilized patients with peripheral neuropathy. See [Contraindications to Mechanical Prophylaxis \(VTE-B\)](#). (CLOTS Trial Collaboration. Lancet 2009. 373:1958-1965.)

^eMost data come from surgical patients; this is an extrapolation to the medical population. See [Contraindications to Mechanical Prophylaxis \(VTE-B\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

VTE PROPHYLAXIS FOLLOWING DISCHARGE AND FOR AMBULATORY CANCER PATIENTS AT RISK^a

AT-RISK POPULATION

- Adult medical or surgical patient
- Diagnosis of cancer
- Patient received VTE prophylaxis during hospitalization
- Cancer inpatient intended for discharge
- Outpatients at risk
- Providers are encouraged to discuss VTE risk factors, risks and benefits of VTE prevention, and the importance of patient adherence to care programs

Surgical oncology patient

Out-of-hospital primary VTE prophylaxis is recommended for up to four weeks post-operation for high-risk abdominal or pelvic cancer surgery patients^f
[See Inpatient/Outpatient Prophylactic Anticoagulation Treatment \(VTE-C\)](#)

Medical oncology patient

Multiple myeloma patients receiving thalidomide, lenalidomide or pomalidomide:
• High risk: Recommend anticoagulant VTE prophylaxis^g
• Low risk: Recommend aspirin^g

Other outpatient settings
• No routine VTE prophylaxis recommended outside of a clinical trial setting

^a[See VTE Risk Factors in Cancer Patients \(VTE-A\).](#)

^fHigh-risk abdominal/pelvic cancer surgery patients include patients undergoing surgery for gastrointestinal malignancies, patients with a previous history of VTE, anesthesia time greater than 2 hours, bed rest >4 days, advanced-stage disease, and patient age greater than 60 years.

^gMultiple myeloma patients receiving thalidomide/lenalidomide/pomalidomide: in combination with high-dose dexamethasone (≥480 mg per month) or doxorubicin or multi-agent chemotherapy or for myeloma patients with two or more individual or myeloma risk factors ([See VTE Risk Factors in Cancer Patients \[VTE-A 2 of 3\]](#)), recommended prophylaxis is LMWH (enoxaparin 40 mg subcutaneous every 24 hours or its equivalent) or warfarin (adjusted to INR 2-3). For low-risk myeloma patients with one or fewer individual or myeloma risk factors, aspirin 81–325 mg daily may be used. Aspirin should not be used in non-myeloma patients for VTE prevention.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Acute Superficial Vein Thrombosis (SVT)

DIAGNOSIS

Clinical suspicion of SVT:

- Swelling of unilateral extremity
- Heaviness in extremity
- Pain in extremity
- Unexplained persistent calf cramping
- Swelling in face, neck, or supraclavicular space

WORKUP/IMAGING

- Comprehensive medical history and physical examination
- CBC with platelet count
- PT, aPTT
- Liver and kidney function tests
- Consider venous US based on clinical judgment

IMAGING FINDINGS

SVT

Peripheral catheter related

Non-catheter related

SVT TREATMENT

Remove catheter

Not close proximity to deep venous system

Close proximity to deep venous system

- Symptomatic treatment, including warm compresses, anti-inflammatory medications, and elevation
- Re-evaluate if there are progressive symptoms
- Recommend anticoagulation^a for at least 6 weeks (category 2B)
 - ▶ Consider treating up to 12 weeks for close proximity to the common femoral vein

^aSee [Therapeutic Anticoagulation for Venous Thromboembolism \(VTE-D\)](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Acute Deep Vein Thrombosis (DVT)



^aIn cases with high suspicion of DVT and no contraindications, consider initiating early anticoagulation while awaiting imaging results.


Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Acute Deep Vein Thrombosis (DVT)

DVT LOCATION



^bSee [Therapeutic Anticoagulation for Venous Thromboembolism \(VTE-D\)](#).

^cSee [Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment \(VTE-B\)](#).

^dSee [Therapeutic Anticoagulation Failure \(VTE-H\)](#), if extension of VTE or new VTE while on recommended anticoagulation therapy.

^eChoice of regimen should be made based on institutional expertise/preferences in conjunction with interventional radiology or vascular surgery colleagues. (See [Thrombolytic Agents \[VTE-I\]](#)).

^fThe SOX trial was unable to demonstrate that GCS reduced the rate of post thrombotic syndrome. (Kahn SR, Shapiro S, Wells PS, et al. Lancet 2014;383:880-888.)

^gConsider permanent filters only for rare patients with permanent contraindications to anticoagulation.

^hSee [Contraindications to Thrombolysis \(VTE-J\)](#).

ⁱRecommend IVC filter removal, if tolerating anticoagulation.

^jSee [Elements for Consideration in Decision Not To Treat \(VTE-F\)](#).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Acute Deep Vein Thrombosis (DVT)

CATHETER-RELATED DVT: DIAGNOSIS AND TREATMENT



^bSee [Therapeutic Anticoagulation for Venous Thromboembolism \(VTE-D\)](#).

^cSee [Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment \(VTE-B\)](#).

^dSee [Therapeutic Anticoagulation Failure \(VTE-H\)](#), if extension of VTE or new VTE while on recommended anticoagulation therapy.

^eChoice of regimen should be made based on institutional expertise/preferences in conjunction with interventional radiology or vascular surgery colleagues. (See [Thrombolytic Agents \[VTE-I\]](#)).

^hSee [Contraindications to Thrombolysis \(VTE-J\)](#).

^jSee [Elements for Consideration in Decision Not To Treat \(VTE-F\)](#).

^kConsider longer duration anticoagulation in patients with poor flow, persistent symptoms, or unresolved thrombus.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016


Acute Pulmonary Embolism (PE)

PE: DIAGNOSIS

DIAGNOSIS

EVALUATION^a

IMAGING



^aD-dimer has limited utility in cancer patients.


^bIn cases with high suspicion of PE and no contraindications, consider initiating early anticoagulation while waiting for imaging results.

^cRepeat imaging and diagnostic studies are not routinely needed in patients with incidental PE. Consider outpatient management for these patients.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016 Acute Pulmonary Embolism (PE)



^dSee [Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment \(VTE-B\)](#).

^eSee [Therapeutic Anticoagulation for Venous Thromboembolism \(VTE-D\)](#).

^fSee [Therapeutic Anticoagulation Failure \(VTE-H\)](#), if extension of VTE or new VTE while on recommended anticoagulation therapy.

^gIt is unclear if IVC filter placement is beneficial in the absence of lower extremity, IVC, or pelvic DVT. See [Clinical Scenarios Warranting Consideration of Filter Placement \(VTE-G\)](#).

^hConsider permanent filters only for rare patients with permanent contraindications to anticoagulation.

ⁱConsider embolectomy for treatment of massive PE (category 2B).

^jRecommend IVC filter removal, if tolerating anticoagulation therapy.

^kSee [Elements for Consideration in Decision Not To Treat \(VTE-F\)](#).

^lSee [Thrombolytic Agents \(VTE-I\)](#).

^mSee [Contraindications to Thrombolysis \(VTE-J\)](#).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Heparin-Induced Thrombocytopenia (HIT)

DIAGNOSIS AND TREATMENT OF HIT



^aSee [HIT Pre-Test Probability Score Assessment \(HIT-A\)](#).

^bThe 4 T Score has not been validated in cancer patients, so it may have less utility, particularly in patients receiving chemotherapy.

^cA “low” pre-test probability score combined with a negative antibody test is useful in ruling out a diagnosis of HIT; a positive test increases the suspicion for HIT. Patients with 4T scores <4 are very unlikely to have HIT, so routine HIT antibody testing in these patients is probably not advisable. In non-cancer patients with 4T scores of 1–3, the risk of HIT is small but not zero, but this has not been validated in cancer patients. Based on clinical judgment, HIT antibody testing may be warranted in select patients.

^dSee [Therapeutic Options for Heparin-Induced HIT \(HIT-B\)](#).

^eData supporting use of fondaparinux is limited; however, among experienced clinicians it is commonly used for outpatient management in low-risk patients.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2016

Heparin-Induced Thrombocytopenia (HIT)

DIAGNOSIS AND TREATMENT OF HIT

HIT ANTIBODY TEST RESULTS (ELISA)



^dSee Therapeutic Options for HIT; section on warfarin (HIT-B 2 of 2).

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2016

Heparin-Induced Thrombocytopenia (HIT)

HIT PRE-TEST PROBABILITY SCORE ASSESSMENT¹

Suspicion of HIT based on the “4 T’s”	Score	HIT Pre-test Probability Score Criteria		
		2	1	0
Thrombocytopenia	<input type="checkbox"/>	Nadir 20,000–100,000/mcL or >50% platelet fall	Nadir 10,000–19,000/mcL or 30%–50% platelet fall	Nadir <10,000/mcL or <30% platelet fall
Timing of onset platelet fall (days of heparin therapy)	<input type="checkbox"/>	Days 5–10 or ≤ day 1 with recent heparin ²	> day 10 or timing unclear (but fits with HIT)	≤ day 1 (no recent heparin)
Thrombosis or other sequelae	<input type="checkbox"/>	Proven thrombosis, skin necrosis, or ASR ³	Progressive, recurrent, or silent thrombosis; erythematous skin lesions	None
Other cause of platelet fall	<input type="checkbox"/>	None evident	Possible	Definite
Total Pre-test Probability Score	<input type="checkbox"/>	Periodic reassessment as new information can change pre-test probability (eg, positive blood cultures)		



¹Modified with permission from Warkentin TEW, Aird C, and Rand, JH. Platelet-endothelial interactions: sepsis, HIT, and antiphospholipid syndrome. Hematology, 2003;497-519.

²Recent heparin indicates exposure within the past 30 days (2 points) or past 30–100 days (1 point).

³Acute systemic reaction (ASR) following IV heparin bolus.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Heparin-Induced Thrombocytopenia (HIT)

THERAPEUTIC OPTIONS FOR HEPARIN-INDUCED HIT¹

Direct Thrombin Inhibitors (Preferred)²

- **Argatroban (half-life 45 minutes with normal liver function; aPTT ratio 1.5–3)³**
 - ▶ Normal liver function, non-ICU patient: 2 mcg/kg/min adjusted to aPTT ratio (first check in 4 hours)
 - ▶ Abnormal liver function (total bilirubin 1.8–3.6 mg/dL; aspartate transaminase/alanine transaminase [AST/ALT] 150–600 IU/L) or ICU, heart, or multi-organ failure patient: 0.5 mcg/kg/min
 - ▶ Severe liver dysfunction (total bilirubin >3.6 mg/dL or AST/ALT >600 IU/L): Use bivalirudin or fondaparinux

- **Bivalirudin (half-life 25 minutes with normal renal function; aPTT ratio 1.5–2.5)⁴**
 - ▶ Consider strongly for patients with combined hepatic and renal dysfunction
 - ▶ Dosing:
 - ◇ Estimated creatinine clearance (C_{Cr}) >60 mL/min: 0.15 mg/kg/h – adjust to aPTT (first check 2 hours)
 - ◇ Estimated C_{Cr} 45–60 mL/min: 0.1 mg/kg/h
 - ◇ Estimated C_{Cr} 31–44 mL/min: 0.075 mg/kg/h
 - ◇ Estimated C_{Cr} <30 mL/min (no renal replacement therapy): 0.05 mg/kg/h
 - ◇ Renal replacement therapy or combined hepatic/renal failure: Consider argatroban for isolated renal failure or use 0.03 mg/kg/h

[Continued](#)

¹The NCCN Guidelines Panel encourages the development of protocols or order sets for HIT treatment that includes DTI dosing, adjustment in renal or hepatic dysfunction, nursing instructions, and monitoring parameters.

²A DTI is preferred over fondaparinux for the immediate treatment of patients with acute HIT with thrombosis.

³Prescribing information: Argatroban injection, for intravenous infusion only. 2016.

⁴Anaphylaxis has occurred with bivalirudin.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Heparin-Induced Thrombocytopenia (HIT)

THERAPEUTIC OPTIONS FOR HIT

Indirect Factor Xa Inhibitor⁵ (category 2B)

- **Fondaparinux (half-life 17–21 hours with normal renal function)**
 - ▶ For patients with C_{Cr} 30–50 mL/min (clearance reduced by 40%): Consider using a DTI⁶
 - ▶ For patients with C_{Cr} <30 mL/min: Avoid
 - ▶ **Dose**
 - ◇ Body weight <50 kg: 5 mg subcutaneous [SC] daily
 - ◇ Body weight 50–100 kg: 7.5 mg SC daily
 - ◇ Body weight >100 kg: 10 mg SC daily

Warfarin

- Initiate once platelets $\geq 150,000/\text{mcL}$ or return to baseline
- Initial dose 5 mg (consider lower dose for patients: Age >75 years, CYP2C9 inhibitors, poor oral intake, liver disease)
- DTI/fondaparinux-warfarin overlap should be at least 5 days: Continue warfarin until $\text{INR } 2 \geq$ for 24 hours
- Argatroban can increase the INR substantially during warfarin co-therapy; therefore, a higher target INR (approx 4.0) should be achieved before argatroban is discontinued. Bivalirudin slightly prolongs the INR during co-therapy.
- INR and aPTT should be repeated within 2–6 hours after DTI has been discontinued to ensure the INR is still therapeutic when the effects of the DTI are no longer present.
- Alternatively, chromogenic factor X activity, which is not affected by DTIs, can be used to monitor warfarin during co-therapy.
- Treat for at least 4 weeks (no thrombosis) or at least 3 months as dictated by thrombotic event.

Platelet Transfusions

- Avoid unless active bleeding or invasive procedure necessary and platelet count <50,000/mcL.

⁵Used as a second-line agent. Fondaparinux has been rarely associated with HIT.

⁶Prescribing Information: Fondaparinux sodium, solution for subcutaneous injection. 2009.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Splanchnic Vein Thrombosis (SPVT)

SPVT: DIAGNOSIS

CLINICAL SUSPICION OF SPVT^a

- Abdominal or mid-abdominal colicky pain
- Abdominal distention
- Rebound tenderness
- Guarding
- Fever
- Anorexia
- Nausea, vomiting
- Diarrhea
- GI bleeding
- Hepatomegaly
- Ascites
- Lower extremity edema
- Incidental SPVT^b

DIAGNOSTIC EVALUATION

History and physical

- Based on H&P consider further diagnostic testing

Lab testing

- CBC with differential
- PT, aPTT
- Basic metabolic profile
- Hepatic profile
- Serum lactate

Imaging:

- Abdominal duplex US
- Abdominal CTA
- Abdominal MRV

Negative or indeterminate

Continued suspicion

No

Investigate other causes

Yes

Repeat imaging

Positive

[See Treatment \(SPVT-2\)](#)

^aRisk factors relevant to cancer population for SPVT:

- Recent abdominal surgery (eg, splenectomy)
- Abdominal mass
- Pancreatitis
- Cirrhosis
- Thrombophilia
- Exogenous estrogens
- Paroxysmal nocturnal hemoglobinuria (PNH)
- Myeloproliferative neoplasms associated with the *JAK2* V617F mutation (most common) or *CALR* mutation (rare)

^bFor incidental SPVT, weigh the risks and benefits of anticoagulation therapy on an individual basis.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Splanchnic Vein Thrombosis (SPVT)

SPVT: TREATMENT



^cSee [Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment \(VTE-B\)](#).

^dWeigh risks/benefits of anticoagulation, particularly for chronic thromboses. Duration of anticoagulation should be at least 6 months for triggered events (eg, postsurgical) and indefinite if active cancer, persistent thrombophilic state or unprovoked thrombotic event.

^eSee [Therapeutic Anticoagulation for Venous Thromboembolism \(VTE-D\)](#).

^fDecision to offer thrombolysis should be based on local availability/expertise, location of thrombus, and risk of bleeding. Choice of regimen should be made based on institutional expertise/preferences in conjunction with interventional radiology or vascular surgery colleagues.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

VTE RISK FACTORS IN CANCER PATIENTS

General patient risk factors

- Active cancer
- Advanced stage of cancer
- Cancer types at higher risk:
 - ▶ Brain
 - ▶ Pancreas
 - ▶ Stomach
 - ▶ Bladder
 - ▶ Gynecologic
 - ▶ Lung
 - ▶ Lymphoma
 - ▶ Myeloproliferative neoplasms
 - ▶ Kidney
 - ▶ Metastatic cancers
- Regional bulky lymphadenopathy with extrinsic vascular compression
- Familial and/or acquired hypercoagulability (including pregnancy)
- Medical comorbidities: Infection, renal disease, pulmonary disease, congestive heart failure, arterial thromboembolism
- Poor performance status
- Older age

High-risk outpatients on chemotherapy, based on combinations of the following risk factors¹

- Active cancers associated with high incidence of VTE: Stomach, pancreas, lung, lymphoma, gynecologic, bladder, and testicular
- Prechemotherapy platelet count >300,000/mcL
- Prechemotherapy white blood cell (WBC) >11,000/mcL
- Hemoglobin <10 g/dL
- Use of erythropoiesis-stimulating agents
- Body mass index (BMI) 35 kg/m² or greater
- Prior VTE

Treatment-related risk factors

- Major surgery
- Central venous catheter/IV catheter
- Chemotherapy such as:
 - ▶ Thalidomide/lenalidomide/pomalidomide plus high-dose dexamethasone
- Exogenous hormonal therapies such as:
 - ▶ Hormone replacement therapy (HRT)
 - ▶ Contraceptives
 - ▶ Tamoxifen/raloxifene
 - ▶ Diethylstilbestrol

Modifiable risk factors

- Smoking, tobacco
- Obesity
- Activity level/exercise

¹Additional prospective randomized data are required to assess the benefit and safety of routine VTE prophylaxis in a cancer outpatient population with a favorable risk-benefit ratio. Listed risk factors are limited to cancer populations included in recent prospective, observational studies of solid tumor or lymphoma outpatients receiving chemotherapy (Khorana AA, Kuderer NM, Culakova E, et al. Development and validation of a predictive model for chemotherapy-associated thrombosis. *Blood* 2008;111(10):4902-4907; and Mandalà M, Barni S, Prins M, et al. Acquired and inherited risk factors for developing venous thromboembolism in cancer patients receiving adjuvant chemotherapy: a prospective trial. *Ann Oncol* 2010;21:871-876.)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

VTE RISK FACTORS IN CANCER PATIENTS

Risk Assessment Model for the Management of Venous Thromboembolism in Multiple Myeloma Patients Treated with Thalidomide, Lenalidomide or Pomalidomide¹

Risk factors	Recommended action
Individual risk factors <ul style="list-style-type: none"> • Obesity (BMI ≥30 kg/m²) • Prior VTE • CVAD or pacemaker • Associated disease: <ul style="list-style-type: none"> ▸ Cardiac disease ▸ Chronic renal disease ▸ Diabetes ▸ Acute infection ▸ Immobilization • Surgery: <ul style="list-style-type: none"> ▸ General surgery ▸ Any anesthesia ▸ Trauma • Use of erythropoietin • Blood clotting disorders 	No risk factor or only one individual/myeloma risk factor: <ul style="list-style-type: none"> • Aspirin 81–325 mg once daily ≥2 individual/myeloma risk factors: <ul style="list-style-type: none"> • LMWH (equivalent to enoxaparin 40 mg once daily); or • Full-dose warfarin (target INR 2–3)
Myeloma-related risk factors <ul style="list-style-type: none"> • Diagnosis of myeloma, <i>per se</i> • Hyperviscosity 	
Myeloma therapy <ul style="list-style-type: none"> • IMiD² in combination with: <ul style="list-style-type: none"> ▸ High-dose dexamethasone (≥480 mg per month) ▸ Doxorubicin ▸ Multiagent chemotherapy 	Therapies as described in the left column: <ul style="list-style-type: none"> • LMWH (equivalent to enoxaparin 40 mg once daily); or • Full-dose warfarin (target INR 2–3)

¹Adapted with permission from Nature Publishing Group. Palumbo A, Rajkumar SV, Dimopolous MA et al, Prevention of thalidomide-and lenalidomide-associated thrombosis in myeloma. *Leukemia* 2008;22:414-423. Copyright 2008. <http://www.nature.com/leu/journal/v22/n2/full/2405062a.html>

²Immunomodulator drugs (IMiDs), including thalidomide, lenalidomide, and pomalidomide.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

VTE RISK FACTORS IN CANCER OUTPATIENTS

Khorana Predictive Model For Chemotherapy-Associated VTE¹

<u>Patient Characteristic</u>	<u>Risk Score</u>
• Site of primary cancer	
▶ Very high risk (stomach, pancreas)	2
▶ High risk (lung, lymphoma, gynecologic, bladder, testicular)	1
• Prechemotherapy platelet count 350 x 10 ⁹ /L or higher	1
• Hemoglobin level less than 10 g/dL or use of red cell growth factors	1
• Prechemotherapy leukocyte count higher than 11 x 10 ⁹ /L	1
• BMI 35 kg/m ² or higher	1

<u>Total Score</u>	<u>Risk Category</u>	<u>Risk of Symptomatic VTE²</u>
0	Low	0.8%–3%
1, 2	Intermediate	1.8%–8.4%
3 or higher	High	7.1%–41%

¹Reproduced and adapted with permission from Khorana AA, Kuderer NM, Culakova E, et al. Development and validation of a predictive model for chemotherapy-associated thrombosis. Blood 2008;111:4902-4907; and Tafur AJ, Kalsi H, Wysokinski WE, et al. The association of active cancer with venous thromboembolism location: a population-based study. Mayo Clin Proc 2011;86:25-30.

²Kearney JC, Rossi S, Glinert K, Henry DH. Venous Thromboembolism (VTE) and Survival in a Cancer Chemotherapy Outpatient Clinic: A Retrospective Chart Review Validation of a VTE Predictive Model. Blood 2009;114: Abstract 2503; Price LH, Nguyen MB, Picozzi VJ, Kozarek RA. Portal vein thrombosis in pancreatic cancer: Natural history, risk factors, and implications for patient management. Proceedings of the 2010 ASCO Gastrointestinal Cancers Symposium: Abstract 143; and Ay C, Dunkler D, Marosi C, et al. Prediction of venous thromboembolism in cancer patients. Blood 2010;116:5377-5382.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

CONTRAINDICATIONS TO PROPHYLACTIC OR THERAPEUTIC ANTICOAGULATION TREATMENT

Absolute

- Recent central nervous system (CNS) bleed, intracranial or spinal lesion at high risk for bleeding
- Active bleeding (major): more than 2 units transfused in 24 hours

Relative

- Chronic, clinically significant measurable bleeding >48 hours
- Thrombocytopenia (platelets <50,000/mcL)
- Severe platelet dysfunction (uremia, medications, dysplastic hematopoiesis)
- Recent major operation at high risk for bleeding
- Underlying hemorrhagic coagulopathy
- High risk for falls (head trauma)
- Neuraxial anesthesia/lumbar puncture^{1,2}
- Interventional spine and pain procedures³

CONTRAINDICATIONS TO MECHANICAL PROPHYLAXIS

Absolute

- Acute DVT
- Severe arterial insufficiency (pertains to GCS only)

Relative

- Large hematoma
- Skin ulcerations or wounds⁴
- Thrombocytopenia (platelets <20,000/mcL) or petechiae
- Mild arterial insufficiency (pertains to GCS only)
- Peripheral neuropathy (pertains to GCS only)

¹Refer to institutional-specific anesthesia practice guidelines, if available. Twice daily prophylactic dose UFH (5000 units every 12 hours) and once daily LMWH (eg, enoxaparin 40 mg once daily) may be used with neuraxial anesthesia. Twice daily prophylactic dose LMWH (eg, enoxaparin 30 mg every 12 h), prophylactic dose fondaparinux (2.5 mg daily), and therapeutic dose anticoagulation should be used with extreme caution with neuraxial anesthesia. The safety of thrice daily prophylactic dose UFH in conjunction with neuraxial anesthesia has not been established. (Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Third Edition). Reg Anesth Pain Med 2010;35:64-101.)

²Timing of LMWH:

For LMWH, placement or removal of a neuraxial catheter should be delayed for at least 12 hours after administration of prophylactic doses such as those used for prevention of DVT. Longer delays (24 hours) are appropriate to consider for patients receiving therapeutic doses of LMWH. A post-procedure dose of LMWH should usually be given no sooner than 4 hours after catheter removal. (FDA Drug Safety Communication. Updated recommendations to decrease risk of spinal column bleeding and paralysis in patients on low molecular weight heparins. November 6, 2013: <http://www.fda.gov/downloads/Drugs/DrugSafety/UCM373735.pdf>.)

In all cases, a benefit-risk assessment should consider both the risk for thrombosis and the risk for bleeding in the context of the procedure and patient risk factors.

³Narouze S, Benzon HT, Provenzano DA, et al. Interventional spine and pain procedures in patients on antiplatelet and anticoagulant medications: guidelines from the American Society of Regional Anesthesia and Pain Medicine, the European Society of Regional Anaesthesia and Pain Therapy, the American Academy of Pain Medicine, the International Neuromodulation Society, the North American Neuromodulation Society, and the World Institute of Pain. Reg Anesth Pain Med 2015;40:182-212.

⁴Skin ulcerations and wounds are more problematic with the use of GCS.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

INPATIENT/OUTPATIENT PROPHYLACTIC ANTICOAGULATION TREATMENT^{1,2,3}

Agent	Standard Dosing	Obesity Dosing (BMI ≥40 kg/m ²) ⁴
LMWH⁵		
• Dalteparin	5,000 units SC daily (category 1 for inpatient)	Consider 7500 units SC daily (limited data)
• Enoxaparin	40 mg SC daily (category 1 for inpatient)	Consider 40 mg SC every 12 hours
Fondaparinux ⁶	2.5 mg SC daily (category 1 for inpatient)	Consider 5 mg SC daily (limited data)
UFH	5,000 units SC every 8–12 hours (category 1 for inpatient)	Consider 7500 units SC every 8 hours
Aspirin	81–325 mg daily (for low-risk multiple myeloma outpatients only) ⁷	
Warfarin	Adjusted to INR 2–3 ⁸	

[For Diagnosis and Treatment of HIT See \(HIT-1\)](#)

¹Agent selection based on: Renal failure ($C_{cr} < 30$ mL/min), FDA approval, cost, ease of administration, monitoring, and ability to reverse anticoagulation.

²Follow institutional standard operating procedures (SOPs) for dosing schedules. If no SOPs then use the American College of Chest Physicians (ACCP) recommendations. (Kahn SR, Lim W, Dunn AS, et al. Prevention of VTE in nonsurgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:e195S-226S; and Garcia DA, Baglin TP, Weitz JI, Samama MM. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:e24S-43S www.chestjournal.org).

³Following initiation of heparin: Hemoglobin, hematocrit, and platelet count every 2–3 days up to at least day 14 and every two weeks thereafter or as clinically indicated.

⁴Given the impact of renal insufficiency on clearance of enoxaparin and fondaparinux, UFH or dalteparin are recommended for obese patients with severe renal impairment ($C_{cr} < 30$ mL/min).

⁵LMWHs should be used with caution in patients with renal dysfunction. Dose adjustments and Anti-Xa monitoring may be required. Follow package insert for renal dysfunction dosing.

⁶Fondaparinux is contraindicated in patients with creatinine clearance < 30 mL/min. Use with caution in patients with moderate renal insufficiency (creatinine clearance 30–50 mL/min), weight < 50 kg, or age > 75 years.

⁷Use only for lower risk multiple myeloma outpatients with one or fewer individual or myeloma risk factors ([See VTE Risk Factors in Cancer Patients \[VTE-A\]](#)).

⁸Warfarin (INR 2–3) or LMWH (eg, enoxaparin 40 mg SC every 24 hours) are prophylaxis options for select high-risk myeloma outpatients receiving highly thrombotic anti-angiogenic therapy (ie, multiple myeloma patients receiving thalidomide/lenalidomide/pomalidomide in combination with high-dose dexamethasone [≥ 480 mg per month] or doxorubicin or multi-agent chemotherapy) or for myeloma patients with two or more individual or myeloma risk factors ([See VTE Risk Factors in Cancer Patients \[VTE-A\]](#)).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

THERAPEUTIC ANTICOAGULATION FOR VENOUS THROMBOEMBOLISM^{1,2,3}

Acute Management: At diagnosis or during diagnostic evaluation:

- **LMWH (preferred)**⁴
 - ▶ Dalteparin (200 units/kg SC daily)⁵
 - ▶ Enoxaparin (1 mg/kg SC every 12 hours)
- Fondaparinux⁶ (5 mg [<50 kg]; 7.5 mg [50–100 kg]; 10 mg [>100 kg] SC daily)
- Unfractionated heparin IV (80 units/kg load, then 18 units/kg/h, target aPTT of 2–2.5 x control or per hospital standard operating procedures [SOPs])
- Unfractionated heparin SC 333 units/kg load, then 250 units/kg every 12 hours⁷
- Direct oral anticoagulants (DOACs) are not recommended (apixaban, dabigatran, edoxaban, and rivaroxaban require additional clinical experience and research to provide data regarding risks/benefits and guidance for their safe and effective use in cancer patients).

[Continued](#)

[For Diagnosis and Treatment of HIT, see \(HIT-1\)](#)

[For Reversal of Anticoagulation, see \(VTE-E\)](#)

¹Agent selection based on: Renal failure ($C_{cr} <30$ mL/min), inpatient/outpatient, FDA approval, cost, ease of administration, monitoring, and ability to reverse anticoagulation.

²Follow institutional standard operating procedures (SOPs) for dosing schedules. If no SOPs then use the American College of Chest Physicians (ACCP) recommendations.

Garcia DA, Baglin TP, Weitz JI, Samama MM. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012; 141:e24S-43S.

³Following initiation of heparin: Hemoglobin, hematocrit, and platelet count every 2–3 days for the first 14 days and every two weeks thereafter or as clinically indicated.

⁴LMWH should be used with caution in patients with renal dysfunction. Dose adjustments should be considered in patients with severe renal dysfunction (creatinine clearance <30 mL/min). Follow package insert for renal dysfunction and body weight dosing. Anti-Xa monitoring (peak and trough) of LMWH has been recommended for patients with severe renal dysfunction, although limited data are available to support the clinical relevance of anti-Xa levels.

⁵For chronic treatment, dalteparin 150 units/kg daily after 30 days. Although each of the LMWH agents has been studied in randomized controlled trials in cancer patients, the efficacy of dalteparin in this population is supported by the highest quality evidence and is the only LMWH approved by the FDA for this indication. Lee AYY, Levine MN, Baker RI, et al. Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. New Eng J Med 2003;349:146-153.

⁶Fondaparinux is contraindicated in patients with creatinine clearance <30 mL/min. It should be used with caution in patients with moderate renal insufficiency (creatinine clearance 30–50 mL/min), weight <50 kg, or age >75 y.

⁷Kearon C, et al. Comparison of fixed-dose weight-adjusted unfractionated heparin and low-molecular-weight heparin for acute treatment of venous thromboembolism. JAMA 2006;296:935-942.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

THERAPEUTIC ANTICOAGULATION FOR VENOUS THROMBOEMBOLISM^{1,2,3}

Chronic Management:¹

- LMWH⁴ (category 1) is preferred for the first six months as monotherapy without warfarin in patients with proximal DVT or PE and prevention of recurrent VTE in patients with advanced or metastatic cancer.
- Warfarin (2.5–5 mg every day initially, subsequent dosing based on INR value; target INR 2–3)
 - ▶ If warfarin is selected for chronic anticoagulation, initiate warfarin concurrently with the parenteral agent used for acute therapy and continue both therapies for at least 5 days and until the INR ≥ 2 for 24 hours.
 - ▶ During the transition to warfarin monotherapy, the INR should be measured at least twice weekly. Once the patient is on warfarin alone, the INR should be measured initially at least once weekly. Once the patient is on a stable dose of warfarin with an INR between 2 and 3, INR testing can be gradually decreased to a frequency no less than once monthly.
- DOAC are not recommended (apixaban, dabigatran, edoxaban, and rivaroxaban require additional clinical experience and research to provide data regarding risks/benefits and guidance for their safe and effective use in cancer patients).

Duration of Anticoagulation as Recommended by Guideline:

- Minimum time of 3 months
- For non-catheter-associated DVT or PE recommend indefinite anticoagulation while cancer is active, under treatment, or if risk factors for recurrence persist.
- For catheter-associated thrombosis, anticoagulate as long as catheter is in place. Recommended total duration of therapy is at least 3 months.
- Providers should continue to discuss with patients the risks/benefits of anticoagulation to determine the appropriate duration of therapy. (See [Elements for Consideration in Decision Not to Treat \[VTE-F\]](#))

[For Diagnosis and Treatment of HIT, see \(HIT-1\)](#)

[For Reversal of Anticoagulation, see \(VTE-E\)](#)

¹Agent selection based on: Renal failure ($C_{cr} < 30$ mL/min), inpatient/outpatient, FDA approval, cost, ease of administration, monitoring, and ability to reverse anticoagulation.

²Follow institutional standard operating procedures (SOPs) for dosing schedules. If no SOPs then use the American College of Chest Physicians (ACCP) recommendations. Garcia DA, Baglin TP, Weitz JI, Samama MM. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012; 141:e24S-43S.

³Following initiation of heparin: Hemoglobin, hematocrit, and platelet count every 2–3 days for the first 14 days and every two weeks thereafter or as clinically indicated.

⁴LMWHs should be used with caution in patients with renal dysfunction. Dose adjustments should be considered in patients with severe renal dysfunction (creatinine clearance < 30 mL/min). Follow package insert for renal dysfunction and body weight dosing. Anti-Xa monitoring (peak and trough) of LMWH has been recommended for patients with severe renal dysfunction, although limited data are available to support the clinical relevance of anti-Xa levels.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

In the event of bleeding or the need for urgent/emergent invasive procedures, anticoagulant effect must be reversed promptly. It is incumbent on the provider to keep in stock the recommended reversal agents for all anticoagulants included in this table. This includes: protamine; vitamin K oral (phytonadione), and IV solution; fresh frozen plasma (FFP); 4-factor prothrombin complex concentrate (4-factor PCC); 3-factor PCC; rhFVIIa activated prothrombin complex concentrates (aPCC) (anti-inhibitor coagulant complex, vapor heated); desmopressin (DDAVP); idarucizumab; and oral charcoal. The reversal guidelines for different anticoagulants are displayed in the following tables:

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
Unfractionated heparin (UFH) (Half-life 1 hour)	<ul style="list-style-type: none"> • Protamine 1 mg/100 units of UFH (taking into account UFH ~1 hour half-life) by slow IV infusion (no faster than 5 mg per min) • Follow aPTT closely • Maximum dose: 50 mg (Examples- Patient bleeds immediately after 5000 unit bolus is given 50 mg of protamine. Patient on 1250 units per hour bleeds and is given 24 mg of protamine to reverse the UFH remaining from the last four hours of the infusion) 	<ul style="list-style-type: none"> • Protamine can cause anaphylaxis if administered too rapidly. • Patients with fish allergies, previous exposure to protamine (eg, NPH insulin), or vasectomized or infertile men are at increased risk. • Excessive protamine (protamine: heparin ratios >1.3:1 mg/U) are associated with platelet dysfunction and decreased thrombin activity, resulting in bleeding. • Protamine reverses a variable amount of LMWH anti-Xa activity (enoxaparin 54%, dalteparin 74%)
Low-molecular-weight heparin (LMWH) (Half-life 3–7 hours)	<ul style="list-style-type: none"> • Protamine 1 mg/mg of enoxaparin or 1 mg/100 units of dalteparin within 8 hours of dose • Protamine 0.5 mg/mg of enoxaparin or 0.5 mg/100 units of dalteparin if dose administered >8 hours prior • If >12 hours since dose, consider clinical scenario (eg, LMWH dose, renal function, bleeding severity) when deciding whether protamine is indicated • Administer protamine by slow IV infusion (no faster than 5 mg per min) • Maximum dose: 50 mg 	

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION: MANAGEMENT OF SUPRATHERAPEUTIC INR

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
<p>Warfarin (Effective half-life 20–60 hours)</p> <ul style="list-style-type: none"> • INR 4.5–10, no bleeding 	<ul style="list-style-type: none"> • Hold warfarin dose • Look for drug or dietary interactions and eliminate them if possible • Follow INR closely¹ (at least daily as an inpatient, every 1–2 days as outpatient) • When INR approaches therapeutic range (INR <4) re-start warfarin at reduced dose (10%–20% dose reduction) if causal factor not present or cannot be eliminated • Recheck INR within 4–7 days • Adjust warfarin dose based on weekly INR until stable 	
<ul style="list-style-type: none"> • INR >10, no bleeding 	<ul style="list-style-type: none"> • Hold warfarin dose • Consider small dose of oral vitamin K₁ (phytonadione) 1–2.5 mg slowly, no faster than 1 mg/min, in patients at high risk of bleeding (may repeat dose in 24 h if INR remains elevated) • Look for drug or dietary interactions and eliminate them if possible • Follow INR closely¹ (at least daily as an inpatient, every 1–2 days as outpatient) • When INR approaches therapeutic range (INR <4) restart warfarin at reduced dose (at least 20% dose reduction) if causal factor not present or cannot be eliminated • Recheck INR within 4–7 days • Adjust warfarin dose based on weekly INR until stable 	<ul style="list-style-type: none"> • Avoid vitamin K₁ SC administration due to erratic absorption and delayed onset compared with oral administration • Vitamin K₁ IV administration can be used for more rapid absorption than tablets

¹The impact of warfarin dose changes can take at least 5–7 days to be fully manifested in the INR.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
<p>Warfarin (continued) (Effective half-life 20–60 hours)</p> <ul style="list-style-type: none"> • Management of urgent surgery (within 24–48 hours) 	<p><u>Within 24 hours:</u></p> <ul style="list-style-type: none"> • Hold warfarin • Vitamin K 1–2.5 mg IV slowly (over 1 hour) • Repeat INR pre-operation to determine need for supplemental fresh frozen plasma (FFP) <p><u>Within 48 hours:</u></p> <ul style="list-style-type: none"> • Hold warfarin • Vitamin K 2.5 mg orally • Repeat INR at 24 and 48 hours to assess need for supplemental vitamin K or FFP 	<ul style="list-style-type: none"> • Infection due to pathogen transmission (all plasma-derived agents; greater risk with FFP compared with solvent/detergent-treated products [3- or 4-factor PCC, aPCC]) • Immune reactions, including allergic/anaphylactic, alloimmunization (vitamin K₁ and all plasma-derived agents; greater risk with FFP compared with solvent/detergent-treated products [3- or 4-factor PCC, aPCC]) • Thromboembolic events (all agents) • Excessive intravascular volume (FFP) • Transfusion-related acute lung injury (FFP) • Pulmonary edema (FFP) • Agglutination reactions/hemolysis due to blood-type incompatibility (FFP) • Transfusion-associated graft-versus-host disease (if not irradiated FFP) • Febrile nonhemolytic transfusion reactions (FFP)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
<p>Warfarin (continued) (Effective half-life 20–60 hours)</p> <ul style="list-style-type: none"> • Life-threatening bleeding 	<ul style="list-style-type: none"> • Hold warfarin dose • Administer vitamin K₁ (phytonadione) 10 mg IV slowly (no faster than 1 mg/min) • Administer 4-factor prothrombin complex concentrate (4-factor PCC) <ul style="list-style-type: none"> ▶ 4-factor PCC dosing (based on units of Factor IX per kg of actual body weight) <ul style="list-style-type: none"> ◇ INR 2–<4: 25 units/kg (maximum 2,500 units) ◇ INR 4–6: 35 units/kg (maximum 3,500 units) ◇ INR >6: 50 units/kg (maximum 5,000 units) ▶ If 4-factor PCC unavailable or patient is allergic to heparin and/or a history of HIT in the last 12 months: <ul style="list-style-type: none"> ◇ INR <4: 3-factor PCC 25 units/kg + FFP 2–3 units ◇ INR >4: 3-factor PCC 50 units/kg + FFP 2–3 units ▶ FFP 15 mL/kg (consider if PCC not available) ▶ rhFVIIa 25 mcg/kg (consider if PCC unavailable or bleeding is unresponsive to PCC). • Monitor INR closely 	<ul style="list-style-type: none"> • Administer IV vitamin K₁ (phytonadione) slowly (less than 1 mg/min). 3 hours or longer may be required for phytonadione to halt or slow active bleeding. Rapid administration of IV vitamin K₁ associated with a higher risk of anaphylaxis (risk ~1 in 3000 doses). • Monitor vital signs closely • PCC and rhFVIIa have been associated with thromboembolic events • Administer 4-factor PCC IV push at a rate not exceeding 5 mL/minute • Administer 3-factor PCC IV push at a rate not exceeding 10 mL/minute • Administer rhFVIIa IV push over 2–5 minutes • For patients with a history of HIT use 3-factor PCC without heparin² (Factor IX complex)

²Prescribing information: Factor IX complex Profilnine. 2014.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

Anticoagulant	Reversal of Anticoagulation	Precautions/Additional Considerations
<p><u>Direct thrombin inhibitors</u>³</p> <ul style="list-style-type: none"> • Bivalirudin (Half-life 25 minutes with normal renal function) 	<ul style="list-style-type: none"> • Discontinue drug. • No specific antidote exists, but beneficial effects have been ascribed to the following: <ul style="list-style-type: none"> ▶ Hemofiltration and hemodialysis are effective in removal of bivalirudin ▶ Animal models and ex-vivo experiments suggest activated prothrombin complex concentrates (aPCCs) (anti-inhibitor coagulant complex, vapor heated 50–100 units/kg IV) or rhFVIIa (90 mcg/kg IV) may be effective ▶ Desmopressin (DDAVP) 0.3 mcg/kg reduced bleeding in animal and ex-vivo models 	<ul style="list-style-type: none"> • Limited data exist to support all reversal strategies for bivalirudin • Repeated doses (more than 3 or 4 doses) of DDAVP associated with tachyphylaxis and hyponatremia • aPCC (eg, anti-inhibitor coagulant complex, vapor heated) and rhFVIIa have been associated with thromboembolic events

³Limited information is available on the clinical efficacy of all these proposed reversal strategies. For life-threatening bleeding, the NCCN Guidelines Panel currently favors use of rhFVIIa as the first-line agent. Hemofiltration or hemodiafiltration can accelerate the clearance of bivalirudin.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
<p><u>Direct thrombin inhibitors⁴</u></p> <ul style="list-style-type: none"> • Argatroban (Half-life 45 minutes with normal hepatic function) 	<ul style="list-style-type: none"> • Discontinue drug. • No specific antidote exists, but beneficial effects have been ascribed to the following: <ul style="list-style-type: none"> ▶ Animal models and case reports suggest unactivated and activated prothrombin complex concentrates (aPCC, vapor heated 50–100 units/kg IV) may be effective ▶ Ex-vivo studies suggest rhFVIIa (90 mcg/kg IV) also may be effective ▶ Desmopressin (DDAVP) (0.3 mcg/kg) reduced bleeding in animal and ex-vivo models 	<ul style="list-style-type: none"> • Limited data exist to support all reversal strategies for argatroban • Repeated doses (more than 3 or 4) of DDAVP associated with tachyphylaxis and hyponatremia • aPCC (eg, anti-inhibitor coagulant complex, vapor heated) and rhFVIIa have been associated with thromboembolic events
<ul style="list-style-type: none"> • Dabigatran (Half-life 14–17 hours) 	<ul style="list-style-type: none"> • Discontinue drug. • Administer idarucizumab, 5 g IV • For special situations with slow or incomplete clearance (eg, renal dysfunction or failure), consider adding to idarucizumab: <ul style="list-style-type: none"> ▶ Oral charcoal if dose within 2 hours of ingestion: standard initial adult dose 50–100 g followed by doses every 1, 2, or 4 hours equivalent to 12.5 g/h ▶ Hemodialysis ▶ Hemodialysis with a charcoal filter 	<ul style="list-style-type: none"> • Limited data exist to support all reversal strategies for dabigatran • aPCC (eg, anti-inhibitor coagulant complex, vapor heated) and rhFVIIa have been associated with thromboembolic events • In patients with renal failure/severe renal insufficiency, dialysis may be helpful in addition to idarucizumab

⁴Limited information is available on the clinical efficacy of all these proposed reversal strategies. For life-threatening bleeding, the NCCN Guidelines Panel currently favors use of aPCC or rhFVIIa as the first-line agent.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
<u>Factor Xa Inhibitor</u> <ul style="list-style-type: none"> Fondaparinux (Half-life 17–21 hours) 	Discontinue drug. No specific antidote exists; however, limited data suggest rhFVIIa (90 mcg/kg IV) may be beneficial.	rhFVIIa has been associated with thromboembolic events
<u>Direct Factor Xa inhibitor</u> <ul style="list-style-type: none"> Rivaroxaban (Half-life 9–12 hours; upper level for elderly) 	Discontinue drug. No specific antidote exists. Beneficial effects have been ascribed to the following: <ul style="list-style-type: none"> Consider oral charcoal if dose within 2 hours of ingestion and repeat within 6 hours: standard initial adult dose 50–100 g followed by doses every 1, 2, or 4 hours equivalent to 12.5 g/h May be helpful based on in vitro and animal models Administer: <ul style="list-style-type: none"> ▶ Anti-inhibitor coagulant complex vapor heated 25–50 units/kg IV or ▶ 4-Factor PCC 25–50 units per kg (based on units of Factor IX per kg of actual body weight) or ▶ rhFVIIa 20–120 mcg/kg IV ▶ If 4-factor PCC is unavailable or patient is allergic to heparin and/or a history of HIT in the last 12 months then administer <ul style="list-style-type: none"> ◊ 3-factor PCC 50 units/kg kg (based on units of Factor IX per kg of actual body weight). 	aPCC (eg, anti-inhibitor coagulant complex, vapor heated) and rhFVIIa have been associated with thromboembolic events
<ul style="list-style-type: none"> Apixaban (Half-life 12 hours) 	Discontinue drug. No specific antidote exists. Beneficial effects have been ascribed to the following: <ul style="list-style-type: none"> Consider oral charcoal if dose within 2 hours of apixaban dose and repeat within 6 hours: standard initial adult dose 50–100 g followed by doses every 1, 2, or 4 hours equivalent to 12.5 g/h May be helpful based on in vitro and animal models Administer: <ul style="list-style-type: none"> ▶ Anti-inhibitor coagulant complex vapor heated 25–50 units/kg IV or ▶ 4-Factor PCC 25–50 units per kg (based on units of Factor IX per kg of actual body weight) or ▶ rhFVIIa 20–120 mcg/kg IV ▶ If 4-factor PCC is unavailable or patient is allergic to heparin and/or a history of HIT in the last 12 months then administer <ul style="list-style-type: none"> ◊ 3-factor PCC 50 units/kg kg (based on units of Factor IX per kg of actual body weight). 	

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY

<u>Anticoagulant</u>	<u>Reversal of Anticoagulation</u>	<u>Precautions/Additional Considerations</u>
<ul style="list-style-type: none"> • Edoxaban (Half-life 10–14 hours) 	<p>Discontinue drug. No specific antidote exists. Beneficial effects have been ascribed to the following:</p> <ul style="list-style-type: none"> • Consider oral charcoal if dose within 2 hours of edoxaban dose and repeat within 6 hours: standard initial adult dose 50–100 g followed by doses every 1, 2, or 4 hours equivalent to 12.5 g/h • May be helpful based on in vitro and animal models • Administer: <ul style="list-style-type: none"> ▶ Anti-inhibitor coagulant complex vapor heated 25–50 units/kg IV or ▶ 4-Factor PCC 25–50 units per kg (based on units of Factor IX per kg of actual body weight) or ▶ rhFVIIa 20–120 mcg/kg IV ▶ If 4-factor PCC is unavailable or patient is allergic to heparin and/or a history of HIT in the last 12 months then administer: <ul style="list-style-type: none"> ◊ 3-factor PCC 50 units/kg (based on units of Factor IX per kg of actual body weight). 	

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**REVERSAL OF ANTICOAGULATION IN THE EVENT OF LIFE-THREATENING BLEEDING OR EMERGENT SURGERY****The preceding tables are comprised of data from the following references:**

- Bhagirath VC, O'Malley L, Crowther MA. Management of bleeding complications in the anticoagulated patient. *Semin Hematol* 2011;48:285-294.
- Bucklin MH, Acquisto NM, Nelson C. The effects of recombinant activated factor VII dose on the incidence of thromboembolic events in patients with coagulopathic bleeding. *Thromb Res* 2014;133:768-771.
- Burbury KL, Milner A, Snooks B, et al. Short-term warfarin reversal for elective surgery--using low-dose intravenous vitamin K: safe, reliable and convenient. *Br J Haematol* 2011;154:626-634.
- Crowther MA and Warkentin TE. Bleeding risk and management of bleeding complications in patients undergoing anticoagulant therapy: focus on new anticoagulant agents. *Blood* 2008;111:4871-4879.
- Elmer J, Wittels KA. Emergency reversal of pentasaccharide anticoagulants: a systematic review of the literature. *Transfus Med* 2012;22:108-115.
- Garcia DA, Baglin TP, Weitz JI, Samama MM. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012;141:e24S-43S.
- Goldstein JN, Refaai MA, Milling TJ, Jr., et al. Four-factor prothrombin complex concentrate versus plasma for rapid vitamin K antagonist reversal in patients needing urgent surgical or invasive interventions: a phase 3b, open-label, non-inferiority, randomised trial. *Lancet* 2015;385:2077-2087.
- Ghanny S, Warkentin TE, Crowther MA. Reversing anticoagulant therapy. *Curr Drug Discovery Technologies* 2012;9:143-149.
- Holbrook A, Schulman S, Witt DM, et al. Evidence-based management of anticoagulant therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. American College of Chest Physicians. *Chest* 2012;141(2 Suppl):e152S-84S.
- Holland L, Warkentin TE et al. Suboptimal effect of a three-factor prothrombin complex concentrate (Profilnine-SD) in correcting supratherapeutic international normalized ratio due to warfarin overdose. *Transfusion* 2009;49:1171-1177.
- Kaatz S, Kouides PA, Garcia DA, et al. Guidance on the emergent reversal of oral thrombin and factor Xa inhibitors. *Am J Hematol* 2012;87 Suppl 1:S141-145.
- Koster A, Buz S, Krabatsch T, et al. Effect of modified ultrafiltration on bivalirudin elimination and postoperative blood loss after on-pump coronary artery bypass grafting: assessment of different filtration strategies. *J Card Surg.* 2008;23:655-658.
- Lubetsky A, Yonath H, Olchovsky D, et al. Comparison of oral vs intravenous phytonadione (vitamin K1) in patients with excessive anticoagulation: a prospective randomized controlled study. *Arch Intern Med* 2003;163:2469-2473.
- Marlu R, Enkelejda H, Paris A, et al. Effect of non-specific reversal agents on anticoagulant activity of dabigatran and rivaroxaban. A randomised crossover ex vivo study in healthy volunteers. *Thromb Haemost* 2012;108:217-224.
- Nutescu EA, Dager WE, Kalus JS, et al. Management of bleeding and reversal strategies for oral anticoagulants: Clinical practice considerations. *Am J Health-Syst Pharm* 2013;70:1914-1929.
- Pernod G, Albaladejo P, Godier A, et al. Management of major bleeding complications and emergency surgery in patients on long-term treatment with direct oral anticoagulants, thrombin or factor-Xa inhibitors: proposals of the working group on perioperative haemostasis (GIHP) - March 2013. *Arch Cardiovasc Dis* 2013;106:382-393.
- Sarode R, Milling TJ, Jr., Refaai MA, et al. Efficacy and safety of a 4-factor prothrombin complex concentrate in patients on vitamin K antagonists presenting with major bleeding: a randomized, plasma-controlled, phase IIIb study. *Circulation* 2013;128:1234-1243.
- Schulman S and Bjisterveld NR. Anticoagulants and their reversal. *Transfus Med Rev* 2007;21:37-48.
- Tanaka KA, Mazzeffi M, Durila M. Role of prothrombin complex concentrate in perioperative coagulation therapy. *J Intensive Care* 2014;2:60.
- Zatta A, McQuilten Z, Kandane-Rathnayake R, et al. The Australian and New Zealand Haemostasis Registry: ten years of data on off-licence use of recombinant activated factor VII. *Blood Transfus* 2015;13:86-99.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

ELEMENTS FOR CONSIDERATION IN DECISION NOT TO TREAT

- Patient refusal
- No therapeutic advantage
 - ▶ Limited survival
 - ▶ High risk
 - ▶ No planned oncologic intervention
- No palliative benefit (eg, alleviate dyspnea, prevent leg swelling)
- Unreasonable burden of anticoagulation treatment
 - ▶ Painful injections
 - ▶ Frequent monitoring with phlebotomy

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

CLINICAL SCENARIOS WARRANTING CONSIDERATION OF FILTER PLACEMENT^{1,2}

- **Absolute contraindication to therapeutic anticoagulation**³
- **Failure of anticoagulation (category 2B)**⁴
- **Patient non-compliance with prescribed anticoagulation (category 2B)**
- **Baseline cardiac or pulmonary dysfunction severe enough to make any new or recurrent PE life threatening (category 2B)**
- **Patient with documented multiple PE and chronic pulmonary hypertension (category 2B)**

¹It is unclear if IVC filter placement is beneficial in the absence of lower-extremity or intra-abdominal/pelvic DVT.

²Retrievable filter preferred. Consider permanent filters only for rare patients with permanent contraindications to anticoagulation or with chronic comorbidities.

³[See Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment \(VTE-B\).](#)

⁴[See Therapeutic Anticoagulation Failure \(VTE-H\).](#)


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease



¹Anticoagulation failure is defined as an extension of DVT or new DVT or PE, while on therapeutic levels of recommended anticoagulation therapy ([VTE-D](#)).
²If failure of anticoagulation involves a PE or central DVT progression, recommend placement of a filter to prevent fatal PE, and consider thrombolysis for high-risk patients (for submassive or massive PE or massive DVT that is life- or limb-threatening).
³Evaluate for HIT ([HIT-1](#)). If clinical suspicion of HIT is high, see ([HIT-1](#)).
⁴Therapeutic aPTT range based on hospital SOP range or 2.0–2.5 x control, if local ranges are unavailable.
⁵Retrievable filter preferred. Consider permanent filters only for rare patients with permanent contraindications to anticoagulation.
⁶Anti-Xa levels may be considered in patients who are underweight, obese, renally impaired, or for whom compliance is a concern. Anti-Xa levels should be checked at their peak at 4 hours after dosing for both twice-daily or once-daily dosing regimens. Reference ranges are not clinically validated and can vary by facility. Suggested “therapeutic range” is usually 0.6–1.0 units/mL for twice-daily dosing or 1–2 units/mL for once-daily dosing.
⁷Although data are limited, doses are generally increased by 25%.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

THROMBOLYTIC AGENTS

- **Deep Vein Thrombosis:**¹
 - ▶ **Alteplase (tPA) 0.5–1.0 mg/h IV**
 - ▶ **Retepase 0.25–0.75 units/h IV**
 - ▶ **Tenecteplase 0.25–0.5 mg/h IV**

- **Pulmonary Embolism:**
 - ▶ **Alteplase (tPA) 100 mg IV over 2 hours**
 - ▶ **Ultrasound-assisted catheter-directed thrombolysis**²

¹Thrombolytic therapy for DVT is generally delivered by catheter-based techniques in conjunction with mechanical thrombectomy.

²Ultrasound-assisted catheter-directed thrombolysis is a treatment option for PE in patients with ≥50% clot burden in one or both main pulmonary arteries or lobar pulmonary arteries, and evidence of right heart dysfunction based on right heart pressures (mean pulmonary artery pressure ≥25 mmHg) or echocardiographic evaluation. The panel recommends alteplase infusion at a rate of 1 mg/h per drug delivery catheter (2 mg/h for bilateral PE) with a total dose of 24 mg. The alteplase regimen is infused for 24 hours with one catheter and 12 hours for two catheters.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

CONTRAINDICATIONS TO THROMBOLYSIS^{1,2}

Absolute

- History of hemorrhagic stroke or stroke of unknown origin
- Intracranial tumor
- Ischemic stroke in previous 3 months
- History of major trauma, surgery, or head injury in previous 3 weeks
- Platelet count below 100,000/mm³
- Active bleeding
- Bleeding diathesis

Relative

- Age greater than 75 years
- Pregnancy or first post-partum week
- Non-compressible puncture sites
- Traumatic resuscitation
- Refractory hypertension (systolic pressure greater than 180 mmHg; diastolic blood pressure greater than 100 mmHg)
- Advanced liver disease
- Infective endocarditis
- Recent GI bleed (last 3 months)
- Life expectancy less than or equal to 1 year

INDICATIONS FOR THROMBOLYSIS

- Limb-threatening/life-threatening acute proximal DVT
- Symptomatic ilial femoral thrombosis
- Massive/life-threatening PE
- Intestinal SPVT with high risk of ischemia

¹Reproduced and adapted with permission from Kearon C, Akl E, Comerota AJ, et al. Antithrombotic therapy for VTE disease: Antithrombotic therapy prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141(2)(Supple):e419S-e494S.


²The risks and benefits of thrombolysis should be assessed on a case-by-case basis by the clinician caring for the patient.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



AT-RISK POPULATION



^aSee [Reversal of Anticoagulation in the Event Of Life-Threatening Bleeding or Emergent Surgery \(VTE-E\)](#).

^bSee [Table 1: Bleeding Risk Assessment \(PMA-A\)](#).

^cSee [Table 2: Perioperative Anticoagulation Management Guideline \(PMA-B\)](#).


^dSee [Table 3: Thromboembolic Risk Assessment for Arterial and Venous Thromboembolism \(PMA-C\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PERIOPERATIVE THROMBOEMBOLISM RISK ASSESSMENT^d



^cSee Table 2: Perioperative Anticoagulation Management Guideline (PMA-B).

^dSee Table 3: Thromboembolic Risk Assessment for Arterial and Venous Thromboembolism (PMA-C).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



Table 1: Bleeding Risk Assessment
Estimated Bleeding Risk of Various Surgical Procedures

Bleeding Risk Category	Type of Surgery or Procedure	Anticoagulation Recommendation ¹
Very high	<ul style="list-style-type: none"> • Neurosurgical procedure (intracranial or spinal) • Prostatectomy or bladder surgery • Heart valve replacement • Coronary artery bypass grafting 	<ul style="list-style-type: none"> • Can initiate prophylactic dose anticoagulation at 24 hours • Consider therapeutic dose anticoagulation at 72 hours
High	<ul style="list-style-type: none"> • Pacemaker or automatic implantable cardioverter defibrillator (AICD) placement • Major cancer surgery • Major vascular surgery (abdominal aortic aneurysm [AAA] repair, peripheral artery bypass) • Reconstructive plastic surgery • Renal or hepatic biopsy • Bowel polypectomy (if part of a colonoscopy) • Major orthopedic surgery • Head and neck surgery 	<ul style="list-style-type: none"> • Can initiate prophylactic dose anticoagulation within 12–24 hours • Consider therapeutic dose anticoagulation at 48–72 hours
Moderate	<ul style="list-style-type: none"> • Major intra-abdominal surgery • Major intra-thoracic surgery 	<ul style="list-style-type: none"> • Can initiate prophylactic dose anticoagulation within 12–24 hours • Consider therapeutic dose anticoagulation at 24–48 hours
Low	<ul style="list-style-type: none"> • Laparoscopic cholecystectomy or hernia repair • Coronary angiography • Arthroscopy • Biopsy (prostate, bladder, thyroid, lymph node) • Bronchoscopy ± biopsy • Central venous catheter removal • Multiple dental extraction or gum surgery 	<ul style="list-style-type: none"> • Can initiate prophylactic dose anticoagulation within 12 hours • Consider therapeutic dose anticoagulation at 24–48 hours
Very low	<ul style="list-style-type: none"> • Minor dental procedures (single tooth extractions or root canals) • Minor dermatologic procedures (excisions of basal and squamous cell carcinomas, actinic keratoses, and malignant or premalignant nevi) • Cataract removal • Electroconvulsive therapy (ECT) • Arthrocentesis • Joint or soft tissue injections • GI endoscopy without biopsy 	<ul style="list-style-type: none"> • Interruption of anticoagulation typically not necessary

¹Reinitiation of therapeutic dose should only be done after careful assessment of bleeding and thromboembolic risk. It is important to make decisions on reinitiation of anticoagulation in conjunction with the surgical team.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**Table 1 Continued: Bleeding Risk Assessment****Estimated Bleeding Risk of Various Dental Procedures¹**

Estimated Bleeding Risk	Low Bleeding Risk	Moderate Bleeding Risk	High Bleeding Risk
Procedures	<ul style="list-style-type: none"> • Supragingival scaling (standard cleaning) • Simple restorations • Local anesthetic injections 	<ul style="list-style-type: none"> • Subgingival scaling • Restorations with subgingival preparations • Standard root canal therapy • Simple extractions • Regional injection of local anesthetics 	<ul style="list-style-type: none"> • Extensive surgery • Apicoectomy (root removal) • Alveolar surgery (bone removal) • Multiple extractions
Suggestions	<ul style="list-style-type: none"> • Do not interrupt warfarin therapy • Use local measures to control bleeding² 	<ul style="list-style-type: none"> • Interruption of warfarin therapy is not necessary • Use local measures to prevent or control bleeding² • Consult with dentist to determine comfort with use of local measures to prevent bleeding when anticoagulation is not interrupted 	<ul style="list-style-type: none"> • May need to reduce INR or return to normal hemostasis • Use local methods to prevent or control bleeding²

¹©2015 University of Washington. Reproduced and used with permission from Suggestions for anticoagulation management before and after dental procedures. Available at: <http://depts.washington.edu/anticoag/home/content/suggestions-anticoagulation-management-and-after-dental-procedures>. Accessed July 15, 2015.

² For local measures to prevent or control bleeding, including the use of a hemostatic agent such as aminocaproic acid 5% mouth rinse, see <http://depts.washington.edu/anticoag/home/content/local-methods-prevent-or-control-bleeding>. Always discuss anticoagulation plan with the dentist/oral surgeon well in advance of the procedure.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Perioperative Management of Anticoagulation and Antithrombotic Therapy

Table 2: Perioperative Anticoagulation Management Guideline¹

TE Risk Category	Bleeding Risk Category		
	Very Low	Low or Moderate	High or Very High
Low risk	<ul style="list-style-type: none"> • Continue warfarin therapy through hospitalization and/or procedure: <ul style="list-style-type: none"> ▪ Adjust dose based on target INR • Continue dabigatran, rivaroxaban, apixaban, or fondaparinux through hospitalization and/or procedure 	<ul style="list-style-type: none"> • Stop warfarin therapy 5–7 days before procedure <ul style="list-style-type: none"> ▪ If INR is >1.5 1–2 days before surgery, give vitamin K 1 mg orally • Stop fondaparinux 5 days before surgery • Stop dabigatran 2 days (CrCl >80 mL/min), 3 days (CrCl 50–79 mL/min), or 4–5 days (CrCl 31–49 mL/min) prior • Stop rivaroxaban 3 days (CrCl >30 mL/min) prior • Stop apixaban 3 days (CrCl ≥50 mL/min) or 4 days (CrCl 30–49 mL/min) prior <p>Post procedure:</p> <ul style="list-style-type: none"> • VTE prophylaxis doses with UFH/LMWH can be considered at 12–24 hours • Restart warfarin once normal diet resumes but no sooner than 24–48 hours post-operation • Restart dabigatran, rivaroxaban, apixaban, or fondaparinux no sooner than 7 days post-operation 	<ul style="list-style-type: none"> • Stop warfarin therapy 5–7 days before procedure <ul style="list-style-type: none"> ▪ If INR is >1.5 1–2 days before surgery, give vitamin K 1 mg orally • Stop fondaparinux 5 days pre-operation • Stop dabigatran 2 days (CrCl >80 mL/min), 3 days (CrCl 50–79 mL/min), or 4–5 days (CrCl 31–49 mL/min) prior • Stop rivaroxaban 3 days (CrCl >30 mL/min) prior • Stop apixaban 3 days (CrCl ≥50 mL/min) or 4 days (CrCl 30–49 mL/min) prior <p>Post procedure:</p> <ul style="list-style-type: none"> • VTE prophylaxis doses with parenteral anticoagulants can be considered at 24 hours • Restart warfarin once normal diet resumes but no sooner than 48–72 hours post-operation • Restart dabigatran, rivaroxaban, apixaban, or fondaparinux no sooner than 7 days post-operation

[Continued](#)

¹These guidelines are meant to supplement but should not supercede clinical judgment. Careful attention to each patient’s individual clinical situation is the best guide to management. In general, we prefer to postpone re-initiation of irreversible anticoagulants such as dabigatran and rivaroxaban until after tolerance with more reversible forms of anticoagulation has been established.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016

Perioperative Management of Anticoagulation and Antithrombotic Therapy

Table 2 Continued: Perioperative Anticoagulation Management Guideline¹

TE Risk Category	Bleeding Risk Category		
	Very Low	Low or Moderate	High or Very High
Moderate risk	<ul style="list-style-type: none"> Continue warfarin therapy through hospitalization and/or procedure: <ul style="list-style-type: none"> Adjust dose based on target INR Continue dabigatran, rivaroxaban, apixaban, or fondaparinux through hospitalization and/or procedure: 	<ul style="list-style-type: none"> Stop warfarin 5 days before surgery Begin “bridge” therapy 2 days after discontinuing warfarin <ul style="list-style-type: none"> Prophylactic or therapeutic dose LMWH (therapeutic preferred for valves and atrial fibrillation) Stop LMWH dosing 24 hours before surgery; <ul style="list-style-type: none"> If once-daily LMWH used, the last dose should be half the total daily dose Stop fondaparinux 5 days pre-operation Stop dabigatran 2 days (CrCl >80 mL/min), 3 days (CrCl 50–79 mL/min), or 4–5 days (CrCl 31–49 mL/min) prior Stop rivaroxaban 3 days (CrCl >30 mL/min) prior Stop apixaban 3 days (CrCl >50 mL/min) or 4 days (CrCl 30–49 mL/min) prior <p>Post procedure:</p> <ul style="list-style-type: none"> VTE prophylaxis doses with UFH/LMWH can be considered at 12–24 hours. Can escalate to therapeutic dose anticoagulation no sooner than 48 hours if prophylactic dose AC tolerated. Restart warfarin once normal diet resumes no sooner than 24–48 hours after the procedure Restart dabigatran, rivaroxaban, apixaban, or fondaparinux no sooner than 7 days post-operation 	<ul style="list-style-type: none"> Stop warfarin 5–7 days before surgery Begin “bridge” therapy 2 days after discontinuing warfarin <ul style="list-style-type: none"> Prophylactic- or therapeutic-dose LMWH (therapeutic preferred for valves and atrial fibrillation) Stop LMWH dosing 24 hours before surgery, <ul style="list-style-type: none"> If once-daily LMWH is used, the last dose should be half the total daily dose Stop fondaparinux 5 days pre-operation Stop dabigatran 2 days (CrCl >80 mL/min), 3 days (CrCl 50–79 mL/min), or 4–5 days (CrCl 31–49 mL/min) prior Stop rivaroxaban 3 days (CrCl >30 mL/min) prior Stop apixaban 3 days (CrCl >50 mL/min) or 4 days (CrCl 30–49 mL/min) prior <p>Post procedure:</p> <ul style="list-style-type: none"> VTE prophylaxis doses with UFH/LMWH can be considered at 24 hours. Can escalate to therapeutic dose anticoagulation no sooner than 72 hours if prophylactic-dose AC tolerated. Restart warfarin once normal diet resumes no sooner than 48–72 hours post-procedure Restart dabigatran, rivaroxaban, apixaban, or fondaparinux no sooner than 7 days post-operation

¹These guidelines are meant to supplement but should not supercede clinical judgment. Careful attention to each patient’s individual clinical situation is the best guide to management. In general, we prefer to postpone re-initiation of irreversible anticoagulants such as dabigatran and rivaroxaban until after tolerance with more reversible forms of anticoagulation has been established.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)



NCCN Guidelines Version 1.2016

Perioperative Management of Anticoagulation and Antithrombotic Therapy

Table 2 Continued: Perioperative Anticoagulation Management Guideline¹

TE Risk Category	Bleeding Risk Category		
	Very Low	Low or Moderate	High or Very High
High risk	<ul style="list-style-type: none"> Continue warfarin therapy through hospitalization and/or procedure: <ul style="list-style-type: none"> Adjust dose based on target INR Continue dabigatran, rivaroxaban, apixaban, or fondaparinux through hospitalization and/or procedure: 	<ul style="list-style-type: none"> Stop warfarin 5 days before surgery Begin “bridge” therapy 2 days after warfarin discontinued: <ul style="list-style-type: none"> Therapeutic-dose LMWH Therapeutic-adjusted IV UFH Stop LMWH dosing 24 hours before surgery <ul style="list-style-type: none"> If once-daily LMWH used, the last dose should be half the total daily dose Discontinue IV UFH 4–6 hours before surgery Stop fondaparinux 5 days pre-operation (consider LMWH bridge as outlined above) Stop dabigatran 2 days (CrCl >80 mL/min), 3 days (CrCl 50–79 mL/min), or 4–5 days (CrCl 31–49 mL/min) prior Stop rivaroxaban 3 days (CrCl >30 mL/min) prior Stop apixaban 3 days (CrCl >50 mL/min) or 4 days (CrCl 30–49 mL/min) prior <p>Post procedure:</p> <ul style="list-style-type: none"> VTE prophylaxis doses with UFH/LMWH can be considered at 12–24 hours. Can escalate to therapeutic-dose anticoagulation within 48 hours if prophylactic-dose AC tolerated. Restart warfarin once normal diet resumes no sooner than 24–48 hours after the procedure Restart dabigatran, rivaroxaban, apixaban, or fondaparinux no sooner than 7 days post-operation 	<ul style="list-style-type: none"> Stop warfarin 5–7 days before surgery Begin “bridge” therapy 2 days after warfarin discontinued: <ul style="list-style-type: none"> Therapeutic-dose LMWH Therapeutic-adjusted IV UFH Stop LMWH dosing 24 hours before surgery <ul style="list-style-type: none"> If once-daily LMWH used, the last dose should be half the total daily dose Discontinue IV UFH 4–6 hours before surgery Stop fondaparinux 5 days pre-operation (consider LMWH bridge as outlined above) Stop dabigatran 2 days (CrCl >80 mL/min), 3 days (CrCl 50–79 mL/min), or 4–5 days (CrCl 31–49 mL/min) prior Stop rivaroxaban 3 days (CrCl >30 mL/min) prior Stop apixaban 3 days (CrCl >50 mL/min) or 4 days (CrCl 30–49 mL/min) prior <p>Post procedure:</p> <ul style="list-style-type: none"> VTE prophylaxis doses with UFH/LMWH can be considered at 24 hours. Can escalate to therapeutic-dose anticoagulation at 72 hours if prophylactic-dose AC tolerated. Restart warfarin once normal diet resumes no sooner than 48–72 hours after the procedure Restart dabigatran, rivaroxaban, apixaban, or fondaparinux no sooner than 7 days post-operation

¹These guidelines are meant to supplement but should not supercede clinical judgment. Careful attention to each patient’s individual clinical situation is the best guide to management. In general, we prefer to postpone re-initiation of irreversible anticoagulants such as dabigatran and rivaroxaban until after tolerance with more reversible forms of anticoagulation has been established.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



Table 3: Thromboembolic Risk Assessment for Arterial and Venous Thromboembolism*

Indication for Anticoagulation or Antiplatelet Therapy					
Arterial Thrombosis			Venous Thrombosis		
TE Risk Category	Event Rate	Mechanical Heart Mechanical Valves	Atrial Fibrillation	Venous Thromboembolism	Thrombophilia (in the context of previous TE)
High risk	>10% per year	<ul style="list-style-type: none"> Mitral valve prosthesis Caged ball (Starr-Edwards) or tilting disc (Bjork-Shiley) aortic valve prosthesis, Stroke or TIA within 6 months 	<ul style="list-style-type: none"> CHADS₂ score ≥5 Stroke or TIA within 3 months Rheumatic valvular heart disease 	<ul style="list-style-type: none"> DVT or PE within 3 months 	<ul style="list-style-type: none"> Protein C deficiency Protein S deficiency Antithrombin deficiency Homozygous factor V Leiden gene mutation Homozygous prothrombin gene mutation Antiphospholipid syndrome
Moderate risk	5%–10% per year	<ul style="list-style-type: none"> Bileaflet aortic valve prosthesis plus: <ul style="list-style-type: none"> Atrial fibrillation Prior stroke Prior TIA Hypertension Diabetes Congestive heart failure (CHF) Age ≥75 years 	<ul style="list-style-type: none"> CHADS₂ score 3 or 4 	<ul style="list-style-type: none"> DVT or PE within 3 to 12 months Recurrent DVT or PE Active cancer or cancer treatment within 6 months 	<ul style="list-style-type: none"> Heterozygous factor V Leiden gene mutation Prothrombin gene homozygous mutation
Low risk	<5% per year	<ul style="list-style-type: none"> Bileaflet aortic valve prosthesis and no other risk factors for stroke 	<ul style="list-style-type: none"> CHADS₂ score of 0 to 2 (and no prior stroke or TIA) 	<ul style="list-style-type: none"> Single VTE event >12 months prior and no other risk factors 	<ul style="list-style-type: none"> Hyperhomocysteinemia elevated Factors VIII, IX, XI

CHADS₂ scoring: congestive heart failure = 1 point; hypertension = 1 point; age greater than 75 = 1 point; diabetes mellitus = 1 point; prior stroke = 2 points

Abbreviations: DVT = deep vein thrombosis; PE = pulmonary embolism; TIA = transient ischemic attack; TE = thromboembolism

*Modified with permission from Douketis JD, Spyropoulos AC, Spencer FA, et al. Perioperative management of antithrombotic therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:e326S-350S.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2016 Cancer-Associated Venous Thromboembolic Disease

Discussion

This discussion is being updated to correspond with the newly updated algorithm. Last updated 10/17/14

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Table of Contents

Overview	MS-2
VTE Risk Assessment in Patients with Cancer	MS-3
Patient-related Factors	MS-3
Cancer-related Factors	MS-4
Treatment-related Factors	MS-4
Diagnosis and Evaluation of VTE in Cancer Patients	MS-6
Diagnosis and Evaluation of DVT	MS-6
Diagnosis and Evaluation of Superficial Vein Thrombosis (SVT)	MS-7
Diagnosis and Evaluation of Splanchnic Vein Thrombosis (SPVT)	MS-8
Diagnosis and Evaluation of PE	MS-10
Anticoagulation in Cancer Patients: Contraindications and Risks	MS-12
Contraindications to Anticoagulation	MS-12
Risks Associated with Anticoagulation Therapy	MS-13

Therapies for Prophylaxis or Treatment of VTE in Cancer Patients **MS-13**

Anticoagulants	MS-14
Aspirin	MS-17
Mechanical Devices	MS-18
VTE Prophylaxis	MS-19
Prophylactic Anticoagulation Therapy	MS-19
Mechanical Prophylaxis	MS-23
VTE Treatment	MS-24
Immediate VTE Treatment	MS-26
Chronic VTE Treatment	MS-26
Treatment of Catheter-related DVT	MS-28
Treatment of SVT	MS-29
Treatment of SPVT	MS-30
Treatment of PE	MS-32
VTE Therapies: Response Assessment	MS-34
Unfractionated Heparin	MS-34
LMWHs and Fondaparinux	MS-34
Direct Thrombin Inhibitors	MS-35
Warfarin	MS-36
Related Issues in VTE Prophylaxis and Treatment	MS-36
Reversal of Anticoagulant Activity	MS-36
Failure of Anticoagulation Therapy	MS-38
Perioperative Management of Anticoagulation and Antithrombotic Therapy	MS-39
Diagnosis and Management of HIT	MS-40
Anticoagulants for the Treatment of HIT	MS-42
Withholding Anticoagulation Therapy: Elements to Consider in the Decision Not to Treat	MS-44
Summary	MS-44
Patient Resources for VTE Management	MS-45
References	MS-46



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Overview

Venous thromboembolism (VTE) is a common and life-threatening condition in cancer patients.^{1,2} Results from a retrospective study of hospitalized adult cancer patients with neutropenia (n=66,106) showed that approximately 3% to 12% of these patients, depending on the type of malignancy, experienced VTE during their first hospitalization.¹ In a recent health claims database analysis of patients undergoing chemotherapy for solid tumors in the ambulatory setting (n=17,284), VTE occurred in 12.6% of patients during the 12-month period from initiation of chemotherapy.³ The incidence ranged from 8% to 19% depending on the tumor type. VTE incidence was 1.4% among age- and gender-matched control cohort without cancer.³ The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for VTE specifically outline strategies to prevent and treat VTE in adult patients with either a diagnosis of cancer or for whom cancer is clinically suspected. These guidelines are characterized by iterative evaluations of the therapeutic advantages of implementing pharmacologic anticoagulation measures based on both the perceived risk of bleeding (ie, contraindications to anticoagulation) and the cancer status of the patient.

In the NCCN Guidelines for VTE, we define VTE broadly to include deep venous thrombosis (DVT), pulmonary embolism (PE), superficial vein thrombosis (SVT), and thrombosis in other vascular territories (eg, portal vein, mesenteric vein, inferior vena cava [IVC] and superior vena cava [SVC], pelvis). DVT management is divided into categories that include the upper extremity and the SVC; the lower extremity including the IVC, pelvis, iliac, femoral, and popliteal veins; the distal lower extremity (eg, calf); the splanchnic vasculature; and catheter-related DVT.

The association of VTE with underlying malignancy was first reported by Armand Trousseau in 1865 and is supported by the results of more recent studies.^{4,5} Pathophysiologic explanations of the etiology of VTE in cancer include known hypercoagulability (eg, pro-coagulants such as tissue factor expressed by cancer cells), vessel wall damage, and vessel stasis from direct compression.⁶⁻⁸ The incidence of cancer-associated VTE is further increased by the presence of additional risk factors, such as acquired or congenital thrombophilia (eg, antiphospholipid syndrome, factor V Leiden), prolonged immobilization, surgical procedures, and chemotherapeutic regimens.^{7,9}

The occurrence of VTE has been reported to increase the likelihood of death for cancer patients by 2- to 6-fold.¹⁰⁻¹⁴ For example, gynecologic cancer patients with PE were found to have a 6-fold increased risk for death at 2 years compared with similar patients without PE.¹³ Furthermore, VTE has been reported to be the most common cause of death at 30-day follow-up among cancer patients undergoing surgery.¹⁵

The critical need for the development of clinical practice guidelines focusing specifically on VTE in cancer patients is further underscored by the results of practice surveys of VTE prophylaxis. The Fundamental Research in Oncology and Thrombosis (FRONTLINE) survey noted that only 50% of surgical oncologists and 5% of medical oncologists routinely used VTE prophylaxis in their cancer patients.¹⁶ Similar results were documented in the multinational IMPROVE and ENDORSE registries of hospitalized medically ill patients in which only 45% of cancer patients received any form of VTE prophylaxis.^{17,18} These results are of particular concern when juxtaposed with a recent review of postmortem reports that showed that approximately 80% of cases of fatal PE occur in nonsurgical patients.¹⁹



VTE Risk Assessment in Patients with Cancer

Many of the risk factors for development of VTE are common to patients with cancer.^{20,21} VTE risk factors in cancer patients can be grouped into 3 general categories: intrinsic and extrinsic patient-related factors, cancer-related factors, and treatment-related factors. The VTE risk factors in the individual cancer patient are likely to be represented by all 3 risk factor categories, and the VTE risk conferred by a single risk factor cannot be evaluated in isolation from the others.

Patient-related Factors

More advanced age, a common characteristic of many cancer patients, was shown to be associated with an increased risk for VTE in some clinical settings.^{1,15,22,23} In addition, obesity has been identified as a risk factor for VTE.^{22,24-26} There is also evidence that pre-chemotherapy thrombocytosis,²⁶⁻²⁸ leukocytosis,²⁶ and hemoglobin level <10 g/dL^{26,27} are predictive of VTE in patients receiving chemotherapy, although the association of anemia with VTE may be complicated by use of erythropoietic stimulating agents (ESAs). Acquired risk factors for VTE include a history of VTE and certain hypercoagulable conditions, such as pregnancy. A history of prior VTE has been identified in a number of studies as an independent risk factor for developing a subsequent VTE.^{15,25,28-31} Moreover, recurrent VTE was found to be more common among patients with cancer; for example, 12-month cumulative incidences of recurrent VTE of 20.7% and 6.8% were reported for patients with and without cancer, respectively, undergoing anticoagulant treatment.³² Although factor V Leiden and prothrombin mutations were identified in 3.7% and 2.6%, respectively, of patients with breast or colon cancer receiving adjuvant chemotherapy in a recent prospective observational study, these inherited risk factors were not associated with an increased risk for VTE among cancer patients.²⁸

A number of other patient-related VTE risk factors, although not exclusive to cancer patients, are commonly found. These risk factors include hospitalization, other medical comorbidities, such as infection, poor performance status, and prolonged immobilization.⁴ In the latest report from the U.S. Centers for Disease Control and Prevention (CDC),²³ VTE events were found to occur at a high rate among hospitalized patients. Among hospitalized adults, VTE was reported in more than 547,000 patients annually (annual rate of 239 per 100,000 persons hospitalized), with more than 28,700 deaths annually in these patients.²³ The risk for VTE increased with age in hospitalized patients. This report confirms that hospitalization is an important risk factor for VTE, and emphasizes the need for greater awareness of VTE risks and appropriate implementation of preventive measures in this setting. Infection has also been identified as an important risk factor for VTE, including in patients with cancer.^{33,34} A recently published case-crossover study in individuals (≥51 years of age) hospitalized for VTE (n=399 among n=16,781 participating in the Health and Retirement Study) reported that infections, use of ESAs, blood transfusions, major surgeries, fractures, immobility, and chemotherapy were significant risk factors for VTE hospitalization.³³ In the subgroup of patients with cancer from this study, the major predictors of VTE hospitalization were infections, blood transfusions, and insertion of a central catheter.³³ In a recent population-based case-control study in patients with hospital-diagnosed VTE (n=15,009), the estimated incidence rate for VTE was increased by 3-fold among patients within the first 3 months after infection, compared with those without an infectious event during the year before VTE (incidence rate ratio=3.3 after adjustment for other VTE risk factors).³⁴

Cancer-related Factors

Several VTE risk factors are exclusive to cancer patients, including the presence of malignancy, chemotherapy, and extrinsic vascular compression due to cancer-associated regional bulky lymphadenopathy. Results from 2 population-based case-control studies showed that the presence of cancer increased the risk for VTE by 4- and 7-fold.^{35,36} An increased risk for VTE in patients with cancer has also been supported by the results of other studies.^{29,37} Furthermore, researchers have reported cancer as the cause of approximately 20% of the VTE cases seen in the community,⁴ and a recent cancer diagnosis and the occurrence of advanced malignancies and distant metastases also increase VTE risk.^{2,25,35,38,39} For example, Blom et al³⁵ reported an adjusted odds ratio of 19.8 for VTE risk in solid tumor cancer patients with distant metastases compared with patients without. In addition, tumor histology has been shown to influence the risk for VTE in patients. Several studies have evaluated the association between different types of cancer and the risk for developing a VTE.^{1-3,10,35,37,40} For example, pancreatic cancer^{1-3,10,37,38,40} and brain tumors^{1,2,35,41-43} were associated with a high risk for VTE in a number of the studies. Adenocarcinomas appear to be associated with a higher risk than squamous cell cancers.³⁷ Although differences in study designs make it difficult to compare VTE rates according to a specific type of malignancy, other cancers that have been associated with an increased risk for VTE include cancers of the stomach, kidney, uterus, lung, ovary, bladder, and testis.^{1,3,22,35,44} In addition, an increased risk for VTE has been observed in certain hematologic malignancies, such as lymphoma, acute leukemia, and multiple myeloma.^{1,45,46} Patients with high-grade lymphoma and acute promyelocytic leukemia appear to be at higher risk than patients with other forms of lymphoma or leukemia.⁴⁵ In a study of patients with high-grade non-Hodgkin's lymphoma,

disease-related venous compression was shown to be the most common cause of VTE in that population.⁴⁷

Several factors associated with an increased risk for VTE in myeloma patients include the diagnosis of multiple myeloma itself, hyperviscosity, and treatment with thalidomide- or lenalidomide-based combination regimens (combined with high-dose dexamethasone, doxorubicin, or multiagent chemotherapy).⁴⁶ Further validation of the influence of these risk factors on VTE rates in patients with myeloma is warranted. In contrast, breast cancer was associated with a relatively low VTE risk in some studies.^{1,11,48} Nevertheless, because of the relatively high prevalence of breast cancer, the occurrence of VTE in a patient with breast cancer is not uncommon.⁴² Furthermore, the risk for VTE was shown to increase by 6-fold when patients with metastatic breast cancer were compared with patients with localized disease.¹¹

Treatment-related Factors

Treatment-related risk factors include surgery, the presence of a central venous access device (CVAD, also known as a catheter), and administration of chemotherapy and other systemic treatments. For example, Heit et al³⁶ reported nearly 22-fold and 8-fold increases in risks for the development of VTE in patients hospitalized or confined to a nursing home with and without recent surgery, respectively, compared with non-institutionalized patients who had not undergone recent surgery.

A number of specific agents used in cancer treatment are associated with an increased risk for developing VTE. A detailed listing of these agents is not provided here; rather, the NCCN Guidelines describe some of the evidence for the association of 3 representative classes of cancer drugs (cytotoxic chemotherapy regimens, hormone therapy with



estrogenic compounds, and antiangiogenic agents) with increased VTE risk.

The association of cytotoxic chemotherapy with the development of VTE in cancer patients has been shown in several studies.^{26,27,49} For example, in one population-based case-control study, odds ratios of 6.5 and 4.1 for development of VTE were determined when cancer patients receiving chemotherapy and cancer patients not receiving chemotherapy, respectively, were compared with patients without a malignant neoplasm.³⁶ In another retrospective study, the annual incidence of VTE was 15% in patients with colorectal cancer treated with chemotherapeutic regimens.⁹ Khorana et al have published a risk assessment model to estimate the risk for VTE in ambulatory cancer patients receiving chemotherapy.²⁶ This risk assessment model has been recently validated and extended by Ay and colleagues,⁵⁰ who identified D dimer and P selectin as additional discriminatory risk factors for VTE in ambulatory cancer patients. However, these laboratory tests are not routinely measured in cancer patients, so their inclusion in routine thrombotic risk assessment should be predicated upon their validation in future studies. The risk factors identified by Khorana et al,²⁶ which formed the basis for the risk assessment models, set the stage for prospective, confirmatory randomized clinical trials evaluating the risks and benefits of risk-targeted VTE prophylaxis in ambulatory cancer patients receiving chemotherapy.

Increased VTE risk was shown to be associated with the use of exogenous hormonal compounds, such as selective estrogen receptor modulators (eg, tamoxifen, raloxifene) for the prevention and treatment of certain estrogen-receptor positive cancers.⁵¹⁻⁵⁵ Use of hormonal compounds, such as hormone replacement therapy^{56,57} or oral contraceptive agents,⁵⁸⁻⁶¹ has also been associated with increased risk for developing VTE. Recent case-control studies and meta-analysis

suggest that for combined oral contraceptives, VTE risks may be different between formulations, depending on the type of progestogen used.^{60,62,63} Diethylstilbestrol phosphate used in combination with doxorubicin for the treatment of hormone-refractory prostate cancer was reported to increase VTE risk when compared with use of doxorubicin alone.⁶⁴ Evidence has been presented to support the association of immunomodulating agents that have antiangiogenic properties (eg, thalidomide in combination with doxorubicin and/or dexamethasone; lenalidomide in combination with dexamethasone) with an increased incidence of VTE when used in the treatment of multiple myeloma (see Guidelines section *Outpatient Prophylactic Therapy in Ambulatory Cancer Patients*).^{46,65-70} Other agents used in supportive cancer care, such as ESAs, have also been associated with the development of VTE.^{3,27,44,71} Concomitant use of erythropoietin with cancer therapies associated with the development of VTE, such as lenalidomide, may further increase VTE risk.⁶⁸

Results from numerous studies have identified the presence of a CVAD as a risk factor for development of an upper-extremity DVT (UEDVT),^{36,72-74} although discrepancies exist concerning the incidence of catheter-related DVT.^{74,75} The association between catheter/device placement and the development of DVT may be the result of venous stasis and vessel injury after insertion of the CVAD^{74,76,77} or infections occurring as a result of catheter placement.^{77,78} Possible reasons for the reported discrepancies in the incidence of catheter-related DVT may include recent improvements in catheter materials and design and the different diagnostic strategies used in some of the studies (ie, clinical, which identifies symptomatic events, versus radiologic surveillance, which identifies symptomatic and asymptomatic events).^{74,75}

Diagnosis and Evaluation of VTE in Cancer Patients

Clinical prediction models, such as the Wells criteria in combination with D-dimer testing, have proven useful in the diagnosis of VTE with comparable results to conventional radiologic imaging strategies.

However, cancer patients comprised a minority of the subjects in these studies.^{79,80} It is therefore unclear whether this strategy is as safe or effective in cancer patients. Although one study employing the Wells criteria and D-dimer testing in the diagnosis of VTE noted the performance of this strategy was comparable in patients with and without cancer, the number of cancer patients (in whom VTE had been excluded by testing) with symptomatic VTE during follow-up was 4-fold higher (2% vs. 0.5%). In addition, the number of false-positive D-dimer assays was 3-fold higher in cancer patients compared with non-cancer patients,⁸¹ and results of a large prospective study of patients with suspected DVT that had been excluded on radiologic testing showed that high D-dimer levels were present in a large percentage of patients with cancer.⁸² D-dimer testing is not recommended for the diagnosis of VTE in cancer patients, and further investigation/validation of D-dimer testing and clinical prediction models is warranted before these strategies are incorporated into the diagnostic evaluation of VTE in cancer patients.

In addition to the imaging described below, the initial diagnostic evaluation of all patients with suspected VTE should include the following components: comprehensive medical history and physical examination; complete blood count (CBC) with platelet count and differential; prothrombin time (PT); activated partial thromboplastin time (aPTT); and comprehensive metabolic panel including liver and kidney function tests (see Guidelines sections on *DVT/SVT: Diagnosis* and *PE: Diagnosis*). In patients with a high suspicion of DVT or PE and without any known contraindications to anticoagulation, initiation of

anticoagulation should be considered while awaiting results from imaging studies.

Diagnosis and Evaluation of DVT

Classic clinical symptoms are not present in all cases of acute DVT. These symptoms may include pain, unilateral edema and heaviness in the extremity distal to the site of the venous thrombosis, or edema in the face, neck, or supraclavicular space, or unexplained persistent cramping. In the prospective, multicenter MASTER registry of patients with VTE, the most common presenting symptoms of DVT were extremity edema, pain, and erythema observed in 80%, 75%, and 26% of patients with DVT, respectively.⁸³ Diagnosis of DVT in adults with cancer should be tempered by an increased level of clinical suspicion on presentation of any clinically overt signs/symptoms that could represent an acute DVT. As mentioned previously, in patients with a high suspicion of DVT and without contraindications to anticoagulation, early initiation of anticoagulation should be considered while awaiting results from imaging studies.

Duplex venous ultrasonography is recommended as the preferred venous imaging method for initial diagnosis of DVT. Duplex ultrasonography allows for both an analysis of venous compressibility and Doppler imaging of venous blood flow,⁸⁴ although assessment of venous compressibility is considered to be more definitive^{73,85}. Other advantageous characteristics of ultrasonography include accuracy for diagnosing symptomatic DVT in femoral and popliteal veins; noninvasive methodology; the lack of need for intravenous contrast agents; ability to be performed at the bedside; and lower cost.^{84,86} It has been reported that 2 normal ultrasound examinations obtained 1 week apart exclude progressive lower-extremity DVT,⁸⁵ although these types of studies have not been performed in cancer patients. Disadvantages



of ultrasonography include difficulties associated with imaging more central veins, such as large pelvic and iliac veins, the proximal subclavian vein, the IVC, and the SVC^{87,88}; a lower sensitivity for diagnosing distal lower-extremity DVT and asymptomatic DVT⁸⁹; limitations associated with bandages, casts, or pain; and results that are more operator dependent.⁹⁰

In cases of negative or indeterminate ultrasound results following repeat venous imaging and a continued high clinical suspicion of DVT, other imaging modalities (listed in order of preference) are recommended: 1) Contrast-enhanced CT, also known as indirect CT venography, is reportedly as accurate as ultrasonography in diagnosing femoro-popliteal DVT and provides accurate imaging of the large pelvic and iliac veins, the IVC, subclavian veins, and the SVC.^{86,91} However, this method requires relatively high concentrations of contrast agent; 2) MRI provides a sensitive and specific evaluation of the pelvic and iliac veins and vena cava without the need for nephrotoxic contrast agents.^{86,92,93} Drawbacks to this method include higher cost, longer imaging times, and limited availability in some practice settings⁹²; and 3) Standard invasive venography, once considered the gold standard for DVT diagnosis, has largely been replaced by less invasive methods.⁹²

Few studies of UEDVT have been performed. Although UEDVT is frequently related to the presence of a CVAD^{72,73,76,94} and associated with device malfunction,⁷⁵ neither a clot within a catheter nor a simple fibrin sheath around a catheter represents a DVT. Ultrasonography has been reported to accurately detect a DVT in peripheral UEDVT involving the brachial, distal subclavian, and axillary veins.⁷³ However, in one study, only 50% of isolated flow abnormalities in the upper extremity were related to the presence of DVT.⁹⁵ A CT venogram may provide a more accurate assessment in cases of isolated flow abnormalities associated with an upper extremity.⁹⁵ CT venography or MR

angiography may be needed to diagnose UEDVT located in the proximal subclavian vein, brachiocephalic vein, or the SVC.^{76,77} Invasive venography for the detection of UEDVT should be performed through a peripheral vessel in the extremity, although venous access may be limited by edema.

The panel recommends that patients diagnosed with calf and UEDVT who have contraindications to anticoagulation therapy be re-evaluated for clot progression (eg, at 1 week for patients with calf DVT) after initial diagnosis. Similarly, patients with catheter-related DVT and central/proximal DVT should undergo follow-up imaging as clinically indicated. Reassessments of contraindications to anticoagulant therapy should accompany imaging evaluations.

The effectiveness of anticoagulation therapy in patients with established DVT should be monitored clinically during and after anticoagulant treatment. Follow-up examinations and imaging evaluations allow physicians to detect clot progression in patients undergoing anticoagulation therapy and DVT recurrence after successful treatment and to identify chronic injury to the venous system. These studies should be performed in response to symptoms.

Diagnosis and Evaluation of Superficial Vein Thrombosis (SVT)

A SVT is distinct from a DVT and generally does not have the same implications for morbidity and mortality as a DVT.^{96,97} Nevertheless, SVT and DVT can occur simultaneously and each predisposes the patient to the other condition.⁹⁷ Few data are available on the incidence of SVT in patients with cancer; it has been estimated that the majority of SVT in the lower extremities occurs in the greater saphenous vein.^{96,97} Although the clinical sequelae of SVT is generally less severe than for DVT, it is important to note that an extensive SVT in the saphenous vein can progress to involve the deep venous system at the saphenofemoral



junction. Such clots can precipitate PE. Therefore, the location and extent of SVT should be evaluated by venous ultrasound if the possibility of proximal deep vein involvement exists.⁹⁷

Diagnosis of SVT is made primarily on the basis of clinical symptoms (tenderness, erythema, and/or an indurated cord associated with a superficial vein) and a negative ultrasound finding for DVT. Progression of symptoms should be accompanied by follow-up imaging. SVT is more likely than DVT to be symptomatic, especially if occurring in the lower extremities. Peripheral catheter-related SVT, sometimes referred to as infusion thrombophlebitis, is often associated with a palpable tender cord along the course of the affected vein.⁹⁷ A key decision point in the treatment algorithm of SVT is the determination of the location of non-catheter-related SVT.

Trousseau's Syndrome

The presence of migratory thrombophlebitis in the presence of cancer should increase clinical suspicion for the presence of a relatively rare condition called Trousseau's syndrome. The clinical characteristics of Trousseau's syndrome can include warfarin resistance, thrombocytopenia, chronic disseminated intravascular coagulation, non-bacterial thrombotic (verrucous) endocarditis, and arterial emboli.^{98,99} Effective treatment of thrombosis in Trousseau's syndrome requires the use of unfractionated or low-molecular-weight heparin (LMWH) or fondaparinux.

Diagnosis and Evaluation of Splanchnic Vein Thrombosis (SPVT)

SPVT refers to a relatively rare group of VTE within the splanchnic vasculature comprising the hepatic (characteristic of Budd-Chiari syndrome), portal, mesenteric, and splenic venous segments.^{100,101}

Thrombotic events may occur in multiple segments (approximately 38%–50% of SPVT cases) or in isolated segments within the splanchnic

vasculature, with isolated portal vein thrombosis (approximately 34%–40% of SPVT cases) being the most common amongst the latter.^{101,102} Limited data are available to assess the relative prognosis of patients with SPVT according to the venous segment affected. In a large single-center retrospective analysis of patients with SPVT (n=832), the 10-year survival rate was significantly decreased among patients with thrombosis in multiple segments compared with those with thrombosis in a single/isolated segment (48% vs. 68%; $P < .001$); the 10-year survival rate for the entire cohort was 60%.¹⁰¹ Moreover, the 10-year survival rate was highest among patients with isolated hepatic vein thrombosis (82%), while the lowest survival rate (63%) was reported in those with isolated portal vein thrombosis ($P = .045$ for comparison of Kaplan-Meier survival estimates across subgroups of isolated SPVT). The investigators attributed the lower survival rate of patients with portal vein thrombosis to the relatively high incidence of malignancies present in this group; in this retrospective study, the presence of malignancy was significantly associated with decreased survival for patients with SPVT, both in univariate and multivariate analyses.¹⁰¹ In a separate retrospective study in patients with extrahepatic portal vein thrombosis (n=172), a concurrent diagnosis of mesenteric vein thrombosis was significantly predictive of decreased survival based on multivariate analysis; presence of cancer was also a significant independent predictor of mortality.¹⁰³ Several smaller retrospective studies have also reported on adverse outcomes for patients with mesenteric vein thrombosis, with a 30-day mortality rate of 20%.^{104,105} Thromboses in the mesenteric vein can lead to intestinal infarction, which is frequently life-threatening.^{104,105} In one study, intestinal infarction was present in 45% of patients diagnosed with mesenteric vein thrombosis, of which 19% were fatal.¹⁰²

Various risk factors have been identified in the development of SPVT, including inherited thrombophilic states (ie, antithrombin deficiency, protein C deficiency, protein S deficiency, Factor V Leiden mutation, prothrombin *G20210A* mutation) and acquired risk factors such as malignancies, myeloproliferative disorders (eg, polycythemia vera, essential thrombocythemia), *JAK2V617F* mutation with or without overt myeloproliferative disorders, paroxysmal nocturnal hemoglobinuria (PNH), abdominal surgery (eg, splenectomy), pancreatitis, and cirrhosis.^{102,103,106-108} In addition, the use of exogenous estrogen, such as oral contraceptives or hormone replacement therapy, has also been linked to SPVT.^{103,106,108} Patients with SPVT may have multiple risk factors, whether inherited and/or acquired. The presence of cancer itself, especially abdominal malignancies, is both a common risk factor for SPVT and a frequent cause of death in cancer patients with SPVT.^{101,103,108} Several retrospective studies have reported cancer to be a significant independent predictor of mortality in patients with SPVT.^{101,103-105} Moreover, among patients with cancer, the presence of SPVT has been associated with decreased survival. Portal vein thrombosis has been reported in about 20% to 30% of patients with hepatocellular carcinoma at the time of diagnosis.¹⁰⁹⁻¹¹¹ In a retrospective study of patients with hepatocellular carcinoma treated at a referral center in Germany (n=389), patients with portal vein thrombosis had significantly decreased median survival (6 months) compared with patients without portal vein thrombosis (16 months); based on multivariate analysis, presence of portal vein thrombosis was a significant independent predictor of 5-year survival in this population.¹¹⁰ The poor prognosis associated with SPVT in patients with hepatocellular carcinoma was demonstrated in another retrospective study (n=194), which also showed significantly decreased median survival in patients with portal vein thrombosis (2.3 months vs. 17.6 months in patients without; $P = .004$).¹⁰⁹ In a recent meta-analysis of 30

randomized controlled trials in patients with previously untreated hepatocellular carcinoma receiving palliative treatments, the presence of portal vein thrombosis was identified as one of the independent predictors of decreased survival.¹¹²

Clinical manifestations of acute SPVT typically include abdominal pain, ascites, hepatomegaly, nausea, vomiting, anorexia, and diarrhea.^{108,113-}

¹¹⁸ SPVT may also be an incidental finding. Among patients with acute thrombosis in the mesenteric vein, intestinal infarction has been reported in 30% to 45% of patients at the time of diagnosis.^{102,104}

Abdominal pain associated with mesenteric vein thrombosis has been described as a mid-abdominal, colicky pain.¹⁰⁸ Fever, guarding, and rebound tenderness may also be present, which may be indicative of progression to bowel infarction.¹⁰⁸ Chronic SPVT may often be asymptomatic due to formation of collateral veins,^{108,113,115,119} although abdominal pain, nausea, vomiting, anorexia, lower-extremity edema, and splenomegaly have been reported with chronic presentations.^{115,118} Weight loss, abdominal distension, and postprandial abdominal pain may also be associated with chronic mesenteric vein thrombosis.¹¹⁹ Presence of splenomegaly and/or esophageal varices is a sign of portal hypertension associated with chronic SPVT, and complications may arise due to bleeding from varices.^{114,115,119}

The diagnostic evaluation includes both imaging and laboratory testing. Diagnosis is confirmed by the absence of blood flow or presence of a thrombus in the splanchnic veins based on noninvasive imaging by duplex ultrasonography, CT angiography (CTA) and/or MR venography (MRV) of the abdomen. Acute SPVT is associated with presenting signs or symptoms of ≤ 8 weeks duration, with no portal cavernoma (cavernous transformation showing a network of collaterals around the portal vein) and no signs of portal hypertension.¹¹⁶ The presence of portal cavernoma on imaging is indicative of chronic thrombosis.^{114,116,120}

For suspected cases of SPVT involving the hepatic and/or portal veins, duplex ultrasonography is considered the initial choice of imaging.^{106,113,114,120} CTA or MRV may be useful in evaluating vascular structure, venous patency, presence of ascites, potential impairment of the bowel and other adjacent organs, and for identifying complications such as bowel ischemia.^{106,119,120} For cases of SPVT involving the mesenteric veins, use of duplex ultrasonography frequently may be limited by overlying bowel gas; for suspected mesenteric vein thrombosis, CTA is the preferred method of diagnostic imaging.^{106,108,119} Once a diagnosis of SPVT has been established, considerations may be given to evaluate the patient for thrombophilia or to test for PNH or the *JAK2* gene mutation. PNH is a rare acquired hematopoietic disorder resulting in chronic hemolysis, and has been associated with a high propensity for venous thrombosis particularly in the splanchnic vasculature.^{121,122} PNH is an important acquired risk factor for SPVT^{106,108}; in a recent *post hoc* analysis (n=77) from a study of patients with Budd-Chiari syndrome, patients who had underlying PNH more frequently presented with additional SPVT (ie, portal, mesenteric, or splenic vein thrombosis) at baseline compared with patients without PNH (47% vs. 10%; *P* = .002).¹²³ The *JAK2V617F* mutation is detected in a high proportion of patients with polycythemia vera, essential thrombocythemia, and primary myelofibrosis, and now constitutes a part of both diagnostic and prognostic assessment of these myeloproliferative disorders.¹²⁴⁻¹²⁷ The presence of myeloproliferative disorders or having *JAK2V617F* mutation, with or without myeloproliferative disorders, is the most common acquired risk factor for SPVT.¹⁰⁶ In the absence of overt myeloproliferative disorders, *JAK2V617F* has been detected in approximately 20% to 40% of patients with SPVT.^{106,128-130} Mutations in exon 12 of *JAK2* may also be associated with SPVT in patients without *JAK2V617F*.¹³¹

Diagnosis and Evaluation of PE

Diagnosis of PE in adults with cancer should include an increased level of clinical suspicion on presentation of any clinically overt signs or symptoms that could represent acute PE. Classic clinical signs and/or symptoms, including unexplained shortness of breath, chest pain—particularly pleuritic chest pain—tachycardia, apprehension, tachypnea, syncope, and hypoxia, are not present in all cases of acute PE. The clinical presentation of PE can range from stable hemodynamics to cardiogenic shock.¹³² In the prospective multicenter MASTER registry, the most common presenting symptoms of PE were dyspnea, pain, and tachypnea, which were present in 85%, 40%, and 29% of patients with PE, respectively.⁸³

Radiographic evidence of DVT is found in up to 50% to 70% of patients presenting with symptomatic PE and vice versa.^{83,133,134} Asymptomatic patients with incidental radiographic findings of PE should be treated similarly to patients with symptomatic PE, as many have subtle clinical symptoms of active disease on further evaluation.¹³⁵ They should undergo additional workup to evaluate for PE; however, repeat imaging is not routinely needed for these patients. As mentioned previously, in patients with a high clinical suspicion of PE and without contraindications to anticoagulation, early initiation of anticoagulation should be considered while awaiting results from imaging studies.

Neither a chest radiograph nor an electrocardiogram (EKG) in a patient with suspected PE is sensitive or specific enough to diagnose PE. However, a chest radiograph facilitates the diagnosis of comorbidities and conditions with clinically similar presentations and is useful in the interpretation of a ventilation-perfusion (VQ) lung scan.¹³⁶ The EKG provides information about existing cardiac disease and PE-related changes. Furthermore, EKG patterns characteristic of right ventricular



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

(RV) strain have been associated with PE,¹³⁷ and inverted T waves in precordial leads may be evident in cases of massive PE.¹³⁸

The NCCN Panel recommends CTA, which allows for indirect evaluation of pulmonary vessels, as the preferred imaging technique for the initial diagnosis of PE in most patients. Advantages of this method include accurate imaging of mediastinal and parenchymal structures; accurate visualization of emboli in many regions of the pulmonary vasculature; the capability to be performed in conjunction with indirect CT venography, which can detect DVT^{86,139} (since the most common cause of PE is DVT in lower extremities or pelvis¹⁴⁰); and the ability to detect signs of RV enlargement, which can be used in assessing the patient's risk for adverse clinical outcomes.¹⁴¹ Disadvantages of CTA include the associated radiation exposure and the need for large amounts of IV contrast, particularly when CTA is followed by indirect CT venography.⁸⁶

Alternative imaging modalities used for the diagnosis for PE include: 1) VQ lung scan; and 2) conventional pulmonary angiography. A VQ scan is associated with less fetal radiation exposure than CTA, so it is useful for pregnant patients and patients with renal insufficiency or untreatable contrast allergies in whom IV contrast is not feasible. It is also less invasive than conventional pulmonary angiography. A normal VQ scan result essentially excludes PE.¹⁴² In a recent non-inferiority study, 1417 patients determined to have a high risk for PE according to the Wells criteria were randomized to undergo CTA or VQ scanning. CTA identified significantly more PE than VQ scans (19.2% vs. 14.2%; 95% CI, 1.1%–8.9%).¹⁴² Elderly patients are more likely than younger patients to be diagnosed with an intermediate probability VQ scan result.¹⁴³ Both intermediate and low-probability VQ scan results lack diagnostic utility and should be considered indeterminate. Further diagnostic testing should be performed if clinically indicated. In a patient

clinically suspected to have a PE, a high-probability VQ scan is diagnostic. Conventional pulmonary angiography (direct pulmonary angiography), often considered to be the gold standard for PE diagnosis, is infrequently used today because of its invasive nature. Rarely, this method is combined with catheter-directed thrombectomy or thrombolysis. These measures should be planned before and executed simultaneously with conventional pulmonary angiography.

Fatality due to PE primarily occurs through RV heart failure and cardiogenic shock.¹³² Since the 3-month mortality rate of patients with PE has been reported to be 15%, outpatient management should be limited to individuals at low-risk for adverse outcomes.¹⁴⁴ The panel recommends that patients with PE be risk-stratified.^{145,146} CTA or echocardiography can be used to assess PE patients for RV enlargement/dysfunction, which is associated with an increased risk for adverse clinical outcomes.^{132,141,144,147-150} Elevated serum troponin levels, which are released due to endomyocardial damage, have also been associated with adverse clinical outcomes^{132,147,151,152} as has the presence of residual DVT on lower-extremity duplex imaging.¹⁵³ A recent study demonstrated that combining the results from at least 2 of the above tests (ie, serum troponin measurement, echocardiography for detecting RV dysfunction, lower-extremity ultrasonography for detecting DVT) improved the specificity and positive predictive value compared with the use of individual tests alone in identifying patients at high risk for PE-related mortality.¹⁴⁵

A clinical risk assessment tool—the Pulmonary Embolism Severity Index (PESI)—has also been used to assess the advisability of outpatient management and intensity of initial follow-up and treatment. The PESI score is a validated patient assessment rule that includes age, sex, a history of heart or lung disease, a history of cancer, and physiologic signs associated with PE that can be used to determine a patient's risk

for an adverse outcome associated with PE.^{154,155} Another stratification tool, known as the RIETE (Computerized Registry of Patients with Venous Thromboembolism) Cancer Score, has been developed to identify individuals at low-risk of mortality from PE and validated in the cancer population.¹⁵⁶ The NCCN Panel recommends that upon diagnosis, all cancer patients with PE be considered for risk stratification with a combination of imaging modalities (CTA or transthoracic echocardiogram to assess RV enlargement or dysfunction) plus serum troponin measurement.^{145,146} The PESI or RIETE score can be included as an adjunctive risk assessment tool, but should not be substituted for the above risk-stratification procedures until validation studies are conducted in patients with cancer.

Anticoagulation in Cancer Patients: Contraindications and Risks

Contraindications to Anticoagulation

Contraindications to anticoagulation can be relative or absolute, and temporary or permanent. Consideration of the degree of contraindication to anticoagulation and its duration are essential when evaluating the risks and benefits of anticoagulation in the individual patient (see Guidelines section *Contraindications to Prophylactic or Therapeutic Anticoagulation Treatment*). Absolute contraindications to anticoagulation include recent central nervous system bleeding, the presence of intracranial or spinal lesions at high risk of bleeding, and major active bleeding requiring >2 units of blood transfusions in 24 hours. Relative contraindications, for which the risks and benefits of anticoagulation must be considered on an individual basis, include: 1) chronic, clinically significant bleeding (for >48 hours); 2) recent major surgery associated with a high risk of bleeding; 3) high risk for falls and/or head trauma; 4) thrombocytopenia (platelets <50,000/mcL); 5) severe platelet dysfunction (eg, due to uremia, medications, dysplastic

hematopoiesis); 6) underlying hemorrhagic coagulopathy; and 7) neuraxial anesthesia or lumbar puncture. Timing is of concern when using LMWH in the setting of neuraxial procedures. Placement or removal of a neuraxial catheter should be delayed for at least 12 hours after administration of prophylactic doses of anticoagulation. Longer delays of up to 24 hours are appropriate to consider for patients receiving therapeutic doses of LMWH. A postprocedure dose of LMWH should usually be given no sooner than 4 hours after catheter removal. The panel recommends frequent re-evaluation of these contraindications and of the risks and benefits of anticoagulation therapy for any cancer patient considered to be at increased risk of bleeding.

Patients with a recent history of bleeding associated with the central nervous system or a spinal lesion are at increased risk of anticoagulant-associated bleeding. Package inserts for all 3 of the LMWHs and fondaparinux include boxed warnings specifying that the risk for spinal or epidural hematoma resulting in long-term paralysis is increased when these anticoagulants are administered to patients receiving epidural or spinal anesthesia or those undergoing spinal puncture.¹⁵⁷⁻¹⁶⁰ Unfractionated heparin (UFH) should also be used with extreme caution in patients receiving spinal anesthesia or undergoing spinal puncture.¹⁶¹ Other factors, such as a patient's risk of falling, should also be considered before anticoagulation therapy is ordered.

A prolonged aPTT is not considered a contraindication to anticoagulation therapy in patients with a lupus inhibitor or lupus anticoagulant, such as those diagnosed with antiphospholipid syndrome. Antiphospholipid antibodies prolong the aPTT by interfering with the interaction of coagulation factors in the patient plasma sample with the phospholipids provided in the aPTT test reagent. Antiphospholipid antibodies have been associated with an increased

risk for venous and arterial thromboembolism and adverse pregnancy outcomes.¹⁶²⁻¹⁶⁴ Any patient who has experienced a thrombotic event and fulfills diagnostic criteria for antiphospholipid syndrome should be considered for indefinite anticoagulation therapy.¹⁶³

Risks Associated with Anticoagulation Therapy

The use of anticoagulants in cancer patients is complicated by the fact that these patients have higher risks of both recurrent VTE and bleeding.^{32,165,166} In one prospective follow-up study of patients undergoing anticoagulation therapy for VTE, the 12-month cumulative incidence of major bleeding was 12.4% and 4.9% in patients with and without cancer, respectively (hazard ratio, 2.2; 95% CI, 1.2–4.1).³² In this study, one-third of all cases of major bleeding occurred during the initial 5 to 10 days of heparinization, and the risk of bleeding increased with the extent of cancer. In contrast to patients without cancer, cancer patients remain at increased risk of bleeding during vitamin K antagonist therapy regardless of International Normalized Ratio (INR) level.^{32,165,166} These findings suggest that factors other than the intensity of anticoagulation, such as thrombocytopenia and organ or vascular invasion by tumors, are responsible for increased bleeding in cancer patients. Subsequent randomized controlled studies of LMWHs and vitamin K antagonists in the chronic treatment of VTE in cancer patients have demonstrated that LMWH is associated with a similar incidence of bleeding events, including major bleeding¹⁶⁷⁻¹⁶⁹; however, in one study, fatal bleeding within the 3-month treatment period was reported in 8% of patients receiving vitamin K antagonists compared with none receiving LMWH.¹⁶⁹ Other risks associated with chronic use of anticoagulants include osteoporosis and heparin-induced thrombocytopenia (HIT) for patients receiving heparins, and drug and food interactions for patients receiving oral anticoagulants. For example, in patients who underwent chronic anticoagulant therapy for 3 to 24 months with an oral

anticoagulant or enoxaparin, decreases in bone mineral density of 1.8% and 3.1% at 1-year follow-up, and 2.6% and 4.8% at 2-year follow-up, respectively, were seen.¹⁷⁰

Warfarin has a very narrow therapeutic window, and its activity is known to be affected by the administration of many other drugs. For example, a number of antibiotics and antifungal therapies, including trimethoprim-sulfamethoxazole, ciprofloxacin, metronidazole, and fluconazole, potentiate the effect of warfarin, whereas other antibiotics such as rifampin and dicloxacillin antagonize the effect of warfarin.^{171,172}

Furthermore, certain chemotherapeutic agents, such as the fluoropyrimidines (5-fluorouracil and capecitabine), are known to increase the INR in patients undergoing warfarin anticoagulation,^{173,174} and drug interactions between warfarin and certain selective estrogen receptor modulators (tamoxifen and raloxifene) have also been reported.¹⁷⁵ Dietary intake of vitamin K and certain dietary supplements can also influence the effects of warfarin.^{176,177} Finally, acetaminophen, found in many medications, can increase the therapeutic effects of warfarin when taken in daily doses exceeding 2 g.¹⁷⁸

Therapies for Prophylaxis or Treatment of VTE in Cancer Patients

The only placebo-controlled, randomized clinical trial on the use of anticoagulants to treat VTE was performed in 1960.^{179,180} Results from this study showed that treatment with heparin followed by warfarin dramatically reduced VTE recurrence and associated mortality in patients with symptoms of acute PE. Although most of the subsequent clinical trials evaluating the use of anticoagulation therapy in the prevention and treatment of VTE have not been placebo-controlled, the evidence supporting the effectiveness of such therapies is strong.¹⁸¹⁻¹⁸³ Clinical evidence for the safety and efficacy of anticoagulation therapy in cancer patients is described later. It is the directive of NCCN that all

adult, hospitalized patients with cancer receive anticoagulation therapy in the absence of contraindications (category 1).

Anticoagulants

Anticoagulation agents used in the prophylaxis and/or treatment of VTE are listed and described according to guideline recommendations (see Guidelines sections *Inpatient/Outpatient Prophylactic Anticoagulation Treatment*, *Therapeutic Anticoagulation Treatment for Venous Thromboembolism*, and *Therapeutic Options for Heparin-Induced Thrombocytopenia [HIT]*). U.S. Food and Drug Administration (FDA) indications and NCCN recommendations for use of each of these therapies are listed in the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Venous Thromboembolic Disease (for the latest version of the NCCN Compendium, please visit www.nccn.org). The panel recommends that agent selection be based on criteria such as the presence of renal insufficiency, FDA approval, cost, ease of administration, need for therapeutic monitoring, and ease of reversibility. Suggested dosing schedules included within this guideline were established according to the NCCN VTE Guidelines Panel consensus and follow, with several exceptions, manufacturer recommendations. To avoid potential conflicts, users can also consult dosing schedules listed in specific institutional standard operating procedure (SOP) documents. Recommendations of the American College of Chest Physicians (ACCP) provide another legitimate source for anticoagulant dosing schedules.¹⁸¹⁻¹⁸³

Low-Molecular-Weight Heparins

LMWHs, such as dalteparin and enoxaparin, are attractive agents for VTE treatment and prevention because they facilitate outpatient treatment and eliminate the need for therapeutic monitoring in most patients. Another LMWH, tinzaparin, has been discontinued in the United States. Although the 2 LMWHs are commonly considered

therapeutically equivalent and are often used interchangeably, few clinical studies have tested whether the clinical effects of these agents are comparable. Furthermore, the agents differ pharmacologically with respect to mean molecular weight, half-life, and ability to inhibit thrombin and factor Xa. Enoxaparin¹⁵⁸ is approved by the FDA for both prophylaxis and immediate treatment of VTE, and dalteparin¹⁶⁰ is approved for VTE and extended treatment of symptomatic VTE in patients with cancer.

NCCN-recommended dosing regimens for dalteparin in immediate VTE treatment are based on the results of clinical studies and panel consensus (see Guidelines section on *Therapeutic Anticoagulation Treatment for Venous Thromboembolism*).^{168,184-188} Extended or chronic anticoagulation therapy with a LMWH may require dosage reduction after an initial period. For example, in the CLOT study, the dalteparin dosing was lowered from 200 units/kg every day to 150 units/kg every day after 1 month.¹⁶⁸ In addition, the European Society for Medical Oncology (ESMO) clinical recommendations for management of VTE in cancer patients specifies using 75% to 80% of the initial dose of LMWH for extended anticoagulation therapy.¹⁸⁹ Only limited evidence exists concerning the safety and efficacy of LMWHs in special populations, such as patients with renal insufficiency, patients with a body mass index (BMI) more than 30 kg/m², patients weighing less than 50 kg, patients aged 70 years or older, and patients with cancer.¹⁹⁰⁻¹⁹² Of the 3 LMWHs, specific dosing recommendations for patients with severe renal insufficiency (creatinine clearance [C_{cr}] <30 mL/min) are available only for enoxaparin.^{158,193} Manufacturer recommendations specify 30 mg enoxaparin subcutaneously daily for VTE prophylaxis and 1 mg/kg subcutaneously every 24 hours for VTE treatment of patients with C_{cr} <30 mL/min. These recommendations are supported by results of a meta-analysis showing enoxaparin to be associated with a 2- to 3-fold

increase in risk of bleeding when administered in standard, unadjusted therapeutic doses to patients with severe renal insufficiency, compared with patients without severe renal insufficiency.¹⁹⁴ In another study, renal clearance of enoxaparin was shown to be reduced by 31% and 44% in patients with moderate (30–60 mL/min) and severe renal impairment (<30 mL/min), respectively, leading the authors to suggest dose reductions for patients with C_{cr} values <50 mL/min.¹⁹⁵ Furthermore, some evidence supports downward dose adjustments of enoxaparin in the management of patients with C_{cr} of 30 to 60 mL/min.¹⁹⁶

Some data are available with respect to the safety of dalteparin in patients with renal insufficiency. In a small study of patients (n=22) treated with dalteparin, mean anti-Xa activity was similar between patients with renal impairment (mean C_{cr} 26 mL/min; range, 16–38) and those with normal renal function (C_{cr} >80 mL/min).¹⁹⁷ In a more recent study of prophylactic dalteparin in critically ill patients (n=138 evaluable) with severe renal impairment (C_{cr} <30 mL/min), no bioaccumulation was detected after a median of 7 days of prophylactic dose dalteparin (5000 IU daily), and treatment was not associated with excessive anticoagulation; peak anti-Xa levels were between 0.29 and 0.34 IU/mL.¹⁹⁸ For cancer patients with C_{cr} <30 mL/min receiving dalteparin for extended treatment of acute VTE, the manufacturer recommends monitoring of peak anti-Xa levels to achieve a target range of 0.5 to 1.5 IU/mL; it is suggested that sampling for anti-Xa levels be taken 4 to 6 hours after dosing, and only after the patient has received 3 to 4 doses of dalteparin.¹⁶⁰

The panel currently recommends using caution when administering LMWH to patients with severe renal insufficiency and following manufacturer specifications when administering enoxaparin to these patients.¹⁵⁸ The panel also recognizes current evidence suggesting caution should be used when administering LMWHs to patients with C_{cr}

less than 50 mL/min. Additional studies are needed to determine the safety of LMWH in patients with compromised renal function, including patients with cancer. Concerns also exist with respect to maintaining and monitoring therapeutic concentrations of anticoagulants in obese patients. In one study, thromboprophylaxis with 5000 units of dalteparin per day was ineffective in reducing the incidence of symptomatic VTE and asymptomatic DVT in patients with a BMI of 40 kg/m² or greater.¹⁹⁹ Hospitalization of morbidly obese cancer patients with administration of UFH should be considered. The panel suggests that each institution prepare a LMWH dosing algorithm tailored for obese patients. Because only limited data are available for the use of LMWHs in patients weighing less than 50 kg,^{157,158,160} the panel also recommends caution when using these agents in patients with low body weight and in elderly patients. LMWHs are contraindicated in patients with HIT, and should only be used with caution in patients with a history of HIT. In this situation, a direct thrombin inhibitor (DTI) or fondaparinux represent safer alternatives (see Discussion section *Related Issues in VTE Prophylaxis and Treatment*). Later sections summarize the clinical evidence for the safety and efficacy of LMWHs in cancer patients (see Discussion sections *VTE Prophylaxis* and *VTE Treatment*).

Factor Xa Inhibitors

Fondaparinux is a specific Factor Xa inhibitor approved by the FDA for the prophylaxis of DVT in patients undergoing hip fracture surgery, hip or knee replacement surgery, or abdominal surgery, and treatment of DVT or acute PE when administered in conjunction with warfarin.¹⁵⁹ Advantages of fondaparinux in the treatment of VTE include specific neutralization of factor Xa, elimination of the need to monitor anticoagulant response in most patients, and the lack of cross reactivity with the antibody associated with HIT.^{159,200-202} However, the use of fondaparinux in patient populations with renal insufficiency, obesity, or HIT has not been well defined,^{192,202} although there is some evidence to

support its safe and effective use for VTE prophylaxis for older patients with a broad range of body weights.²⁰³ Pharmacologic characteristics of fondaparinux include renal elimination and a very long half-life of 17 to 21 hours.¹⁵⁹ Prescribing information for fondaparinux provided by the manufacturer specifies that the drug is contraindicated in patients with severe renal insufficiency ($C_{cr} < 30$ mL/min) and for thromboprophylaxis in patients weighing less than 50 kg undergoing orthopedic or abdominal surgery.¹⁵⁹ It should be used with caution in elderly patients²⁰³ and individuals with moderate renal insufficiency ($C_{cr} < 50$ mL/min).¹⁵⁹ The NCCN Panel recommends against the use of fondaparinux in patients with severe renal insufficiency and advises caution when using fondaparinux in all patients weighing less than 50 kg, patients with renal dysfunction (C_{cr} 30–50 mL/min), and elderly patients (>75 years of age).

Rivaroxaban is an orally administered direct Factor Xa inhibitor approved by the FDA for the prevention of DVT, which may lead to PE, in patients undergoing hip or knee replacement surgery; it is also approved for the treatment of DVT and PE and prevention of stroke and systemic embolism in patients with nonvalvular atrial fibrillation.^{204–206} The drug is primarily eliminated via the kidneys (66% renal excretion) with a lesser proportion cleared by hepatic metabolism (CYP450 3A4 dependent and independent mechanisms). Rivaroxaban is considered a low-clearance drug, as protein binding in plasma is high (92%–95%).²⁰⁴ The half-life is 5 to 9 hours in healthy individuals aged 20 to 45 years and extends to 11 to 13 hours in elderly patients. The prescribing information for rivaroxaban provided by the manufacturer specifies that the drug should be avoided in patients with severe renal impairment ($C_{cr} < 30$ mL/min) and should be used with caution in those with moderate impairment (C_{cr} 30–50 mL/min).²⁰⁴ In randomized clinical trials, rivaroxaban has been evaluated for thromboprophylaxis in hospitalized

acutely ill medical patients²⁰⁷ and for chronic anticoagulation therapy for prevention of recurrent VTE in patients who experienced an initial VTE event (PE with or without DVT),²⁰⁶ in comparison with the LMWH enoxaparin. Although results showed non-inferiority of rivaroxaban compared with enoxaparin, the proportion of enrolled patients with active cancer were very low (5%–6%) in these studies. Until further data become available in cancer patients, this agent is currently not recommended by the NCCN Guidelines Panel for prophylactic or therapeutic anticoagulation in patients with cancer.

Apixaban is another orally administered direct Factor Xa inhibitor recently approved by the FDA. It is currently approved for prevention of thromboembolism in patients with nonvalvular atrial fibrillation, the prevention of VTE after hip and knee arthroplasty, and the treatment of VTE.^{208–211} Apixaban is primarily metabolized via the liver (CYP450 3A4 dependent); renal elimination accounts for about 27% of total drug clearance. The apparent half-life after oral administration of the drug is about 12 hours.²⁰⁸ The prescribing information for apixaban provided by the manufacturer specifies that the drug should be avoided in patients with severe hepatic impairment. Recent randomized clinical trials have evaluated the potential role of apixaban for thromboprophylaxis in hospitalized acutely ill medical patients (in comparison with the LMWH enoxaparin),²¹² and for extended anticoagulation therapy in patients who completed initial anticoagulation for VTE (in comparison with placebo).²¹³ Apixaban (2.5 mg twice daily for 30 days) was not superior to a standard course of enoxaparin (40 mg once daily for 6–14 days) in preventing VTE in acutely ill patients, and was associated with an increased risk for major bleeding events.²¹² In the randomized study involving patients who received 6 to 12 months of anticoagulation for chronic VTE management, extended treatment with apixaban was associated with a significantly decreased risk for recurrent VTE

compared with placebo.²¹³ However, only a small percentage of patients with active cancer (1.7%) were included in this study (see Discussion section *Chronic VTE Treatment*). As in the case with rivaroxaban mentioned above, the NCCN Panel currently does not recommend apixaban for thromboprophylaxis or for the treatment of VTE due to the lack of sufficient clinical data in patients with cancer.

Unfractionated Heparin

UFH is generally administered subcutaneously for VTE prophylaxis (low-dose heparin) and by intravenous infusion for treatment of VTE.²¹⁴ Low-dose UFH (5000 units) administered 3 times/day (every 8 hours) was shown to be more effective than low-dose UFH administered twice a day in preventing DVT in general surgery patients²¹⁵ and is the regimen recommended by the panel for VTE prophylaxis in cancer patients. However, no difference in the overall rate of VTE based on the dosing of prophylactic UFH (5000 units 2 times/day vs. 3 times/day) was observed in a meta-analysis of clinical trials conducted in general medical patients, although a decrease was seen in the combined endpoint of proximal DVT and PE ($P = .05$) and the risk of major bleeding was significantly higher when UFH was administered 3 times daily ($P < .001$).^{216,217}

Initial dosing of UFH in the treatment of VTE is weight-based, with a recommended regimen of 80 units/kg bolus followed by 18 units/kg per hour infusion.¹⁹¹ The safety and efficacy of fixed dose, unmonitored, subcutaneous UFH has been reported to be comparable to LMWH in the treatment of patients with acute VTE,²¹⁸ but further investigation is warranted before this regimen can be routinely used in cancer patients. Patients receiving intravenous UFH must be hospitalized and monitored for anticoagulant response. The panel recommends UFH as the agent of choice in patients with $C_{cr} < 30$ mL/min, because the liver is a main site of heparin biotransformation.^{161,200} Some exceptions include patients

with severe renal dysfunction but without intravenous access and those with a new diagnosis of VTE despite therapeutic doses of UFH. UFH is contraindicated in patients with HIT and should only be used with extreme caution in patients with a history of HIT. In this situation, a DTI or fondaparinux is a better alternative (see Discussion section *Related Issues in VTE Prophylaxis and Treatment*).

Warfarin

Warfarin is an option for long-term treatment of VTE in cancer patients. If warfarin is to be used for chronic therapy, it should be administered concomitantly with UFH, LMWH, or fondaparinux for at least 5 days and until an INR of 2 or more is achieved before discontinuing the parenteral anticoagulant agent. When treating patients with HIT, warfarin should not be initiated until the platelet count has recovered and then it should be overlapped with a DTI or fondaparinux for at least 5 days and until the INR is 2 or more (see Discussion section *Related Issues in VTE Prophylaxis and Treatment*). During the transition to warfarin monotherapy, the INR should be measured at least twice weekly and then at least once every week once the patient has begun receiving warfarin monotherapy. Warfarin can be safely administered to patients with renal insufficiency, although the response to warfarin is accentuated in patients with hepatic insufficiency.²¹⁹

Direct Thrombin Inhibitors

DTIs are discussed in a later section (see Discussion sections *VTE Therapies: Response Assessment* and *Diagnosis and Management of HIT*).

Aspirin

Aspirin (81–325 mg daily) is an option for VTE prophylaxis in only a select group of multiple myeloma patients with one or fewer individual or multiple myeloma-specific risk factors. Aspirin is not considered to be

effective VTE prophylaxis in other settings. In the Women's Health study, a 10-year study of healthy women randomly assigned to aspirin (100 mg) or placebo on alternate days, no significant differences in the incidence of VTE were observed between the 2 arms.²²⁰ Thus, aspirin provided no benefit for initially healthy women who had no or very few risk factors for VTE. A recent double-blind, randomized, controlled study compared the efficacy and safety of aspirin (100 mg daily; n=205) versus placebo (n=197) in patients with a first unprovoked VTE who had completed 6–12 months of oral anticoagulation therapy prior to study initiation.^{221,222} Study treatment was administered for at least 2 years. During the study period (median 24.6 months), VTE recurrence occurred in 14% and 22% of patients who received aspirin and placebo, respectively; this translated to a significant reduction in risk for VTE recurrence with aspirin (6.6% vs. 11.2% per year; hazard ratio, 0.58; 95% CI, 0.36–0.93).²²² The incidence of clinically relevant bleeding events was similar between study arms; major bleeding occurred in 1 patient in each study arm. Similar results were identified in the placebo-controlled randomized controlled ASPIRE study which found that low dose aspirin reduced the incidence of VTE by from 6.5% per year to 4.8% per year (HR with aspirin, 0.74; 95% CI, 0.52 to 1.05; $P = .09$). Although these studies suggested that extended therapy with aspirin following initial oral anticoagulation was beneficial in preventing VTE recurrence, these data cannot be extrapolated to cancer patients with VTE who were excluded from participation in the studies.

Mechanical Devices

Intermittent Pneumatic Compression (IPC) Device

One of the main advantages of an IPC device is the absence of an associated bleeding risk. However, disadvantages include the potential for interference with ambulation and the need to keep the devices in place nearly continuously until patients are fully ambulatory. Graduated

compression stockings (GCS) can be used in conjunction with an IPC device as a method of mechanical prophylaxis.

Vena Cava Filters

Vena cava filters are indicated for prevention of PE in patients who cannot be anticoagulated due to an absolute contraindication to therapeutic anticoagulation or complications from anticoagulation.²²³⁻²²⁷ However, placement of an IVC filter does not prevent DVT and has been associated with an increased risk for recurrent DVT.^{223,228,229} A randomized controlled trial has assessed the efficacy and safety of IVC filters in conjunction with anticoagulation compared with anticoagulant therapy alone in the treatment of acute VTE. However, this pivotal trial did not test the efficacy of IVC filters in the usual clinical scenario in which they are used, in patients without concomitant anticoagulation.^{223,228} It is unclear if IVC filter placement is beneficial in the absence of ilio-popliteal lower-extremity IVC or pelvic DVT.

IVC filters are available as either a retrievable (“optional”) or permanent filter; however, the time period for recovery of a retrievable filter is limited.^{230,231} Results from a retrospective cohort study of 702 patients with IVC filter placement showed that filter retrieval was attempted for only 15.5% of patients who received a retrievable filter, and only 70% of those attempts were successful.²³² No significant differences in PE protection or complication rates were observed between the 2 filter types, although mean follow-up time was limited to 11.5 months. A recent case series of patients who received a Bard G2 or Recovery filters noted filter strut fracture in up to 25% of recipients after a mean follow-up of 24 and 50 months, respectively.²³³ It remains unclear whether the frequency of this complication is device-specific or a characteristic of all filters. Until further data are available, this experience emphasizes the importance of placing filters only in patients



in whom the benefits outweigh the risks, and retrieving filters whenever possible.

VTE Prophylaxis

Prophylactic Anticoagulation Therapy

Inpatient Prophylactic Therapy

Hospitalized patients with cancer are at high risk for VTE.^{1,234} The panel recommends prophylactic anticoagulation therapy for all inpatients with a diagnosis of active cancer or clinical suspicion of cancer and without contraindication to such therapy (category 1). This recommendation is based on an assumption that ambulation in hospitalized cancer patients is inadequate to reduce VTE risk. Recommended anticoagulant options for VTE prophylaxis of cancer inpatients are listed within the Guidelines section *Inpatient/Outpatient Prophylactic Anticoagulation Treatment*.

The LMWHs, fondaparinux, and subcutaneous UFH (5000 units 3 times/day) are category 1 options for inpatient prophylactic therapy. Anticoagulation therapy should be administered throughout the duration of hospitalization. Adult inpatients with cancer should undergo the following evaluation prior to the initiation of thromboprophylaxis: comprehensive medical history and physical examination; CBC with platelet count and differential; PT; aPTT; and liver and kidney function tests.

Studies comparing different anticoagulant regimens for the prevention of VTE in cancer patients have not clearly identified a particular regimen to have superior efficacy. In a randomized multicenter clinical trial, no difference in VTE and bleeding rates were seen for cancer patients receiving perioperative enoxaparin (40 mg) once daily versus low-dose UFH 3 times a day to prevent VTE after major elective abdominal or pelvic surgery.²³⁵ Furthermore, results from a meta-analysis of randomized clinical studies of general surgery patients found LMWHs to be as safe and effective as UFH in the prevention of VTE.²³⁶ However, results from a nonrandomized historically controlled study comparing

the effectiveness of the LMWH dalteparin (5000 units once daily) to low-dose UFH (5000 units 3 times/day) as VTE prophylaxis in high-risk women undergoing surgery for gynecologic cancer indicated that the dalteparin dosing regimen may not be optimal in these patients.²³⁷ More recently, a meta-analysis comparing outcomes of perioperative VTE prophylaxis with LMWH versus UFH in cancer patients showed no difference in rates of mortality, suspected DVT, PE, or bleeding events.²³⁸

For prevention of catheter-related VTE, randomized controlled studies have not established the efficacy of prophylactic doses of LMWH or low-dose warfarin (1 mg daily).²³⁹⁻²⁴¹ A recent randomized trial (n=944) showed that dose-adjusted warfarin of INR 1.5 to 2.0 was significantly more effective than fixed-dose warfarin of 1 mg daily in prevention of catheter-related VTE at a cost of a trend toward more bleeding. However, a separate comparison of warfarin between fixed 1 mg dose and adjusted dose of INR 1.5 to 2 with placebo did not demonstrate a statistically significant reduction in VTE.²⁴² These data suggest that therapeutic or near-therapeutic doses of anticoagulation will likely be necessary for successful prevention of catheter-related VTE. Until additional data are available, the panel does not recommend VTE prophylaxis for cancer patients with a CVAD.

Outpatient Prophylactic Therapy in Ambulatory Cancer Patients

Certain groups of cancer patients are known to remain at risk for VTE after discharge from the hospital. In a retrospective observational study based on data from a large cohort of cancer patients (n=17,874) identified in a health care claims database, VTE (DVT or PE) occurred in nearly 6% of patients during the 12-month index period.²⁴³ A significantly higher proportion of VTE events was diagnosed in the outpatient setting compared with the inpatient setting (78% vs. 22%; $P < .0001$). Moreover, among patients who had a VTE in the outpatient



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

setting, 21% had been hospitalized within 30 days of the VTE event.²⁴³ This observational study suggests that a high proportion of VTE occurs in the cancer outpatient setting, and underscores the need to better identify patients who may benefit from outpatient thromboprophylaxis. The risk for VTE is sufficiently high in some surgical and medical oncology patients that VTE prophylaxis should be considered in the outpatient setting. Cancer patients undergoing abdominal or pelvic surgery should be considered for outpatient prophylaxis.²⁴⁴ Features that identify surgical oncology patients at higher risk for VTE include a previous episode of VTE, anesthesia times longer than 2 hours, advanced-stage disease, perioperative bed rest of 4 days or more, and patient aged 60 years or older.¹⁵ Extended prophylaxis out to 4 weeks post-surgery was associated with a more than 50% reduction in venographic VTE in patients undergoing major abdominal surgery.^{245,246} Since thromboembolic postoperative complications greatly exceeded hemorrhagic complications as a cause of death in the @RISTOS observational cohort study of cancer surgery patients,¹⁵ extended VTE prophylaxis of up to 4 weeks is recommended for cancer surgery patients, particularly the high-risk patients undergoing abdominal or pelvic surgery.

Although there is a lack of consistent evidence to support extended outpatient prophylaxis in most populations of ambulatory medical oncology patients,²⁴⁷ it is recommended for multiple myeloma patients receiving highly thrombogenic regimens. Immunomodulating agents with antiangiogenic properties, such as thalidomide or lenalidomide, have been associated with an increased incidence of VTE in patients with multiple myeloma in the absence of prophylaxis, although the reported rates of VTE vary widely across studies.^{46,65,66,69,247,248} It appears that a number of factors contribute to thrombosis associated with thalidomide or its derivatives,²⁴⁸ and VTE rates are especially high when

thalidomide or lenalidomide is combined with high-dose dexamethasone of 480 mg per month, or doxorubicin or multi-agent chemotherapy regimens.^{46,65,68-70} In a retrospective case-control study of thalidomide or lenalidomide combined with dexamethasone in newly diagnosed patients with multiple myeloma (n=411), the incidence of VTE among the subgroup of patients who received the combination with high-dose dexamethasone (480 mg per 28-day cycle) was 19% with thalidomide and 11% with lenalidomide.²⁴⁹ Data regarding the use of routine thromboprophylaxis were not provided. In an open-label, randomized, non-inferiority trial comparing lenalidomide combined with high-dose dexamethasone (480 mg per 28-day cycle) versus with low-dose dexamethasone (160 mg per 28-day cycle) in previously untreated patients with multiple myeloma (n=445), the incidence of DVT was significantly higher among the patients receiving the combination with high-dose dexamethasone (26% vs. 12%; $P = .0003$).²⁵⁰ Mandatory thromboprophylaxis was added to the study protocol after enrollment of approximately 60% of the patients. The package inserts for thalidomide and lenalidomide include “black box” warnings regarding the VTE risks associated with the administration of these agents.^{251,252}

For patients with multiple myeloma, the panel recommends a prophylaxis strategy based on a risk-assessment model published by the International Myeloma Working Group.⁴⁶ In their publication, VTE prophylaxis with either LMWH (eg, enoxaparin 40 mg daily) or dose-adjusted warfarin (INR 2–3) is recommended for patients with multiple myeloma who are receiving lenalidomide- or thalidomide-based combination regimens associated with a high thrombotic risk or in patients with two or more individual or disease-related risk factors (see Guidelines section on *VTE Risk Factors in Cancer Patients*, VTE-A 2 of 3). Aspirin prophylaxis (81–325 mg daily) is an option for multiple



myeloma patients receiving thalidomide or lenalidomide with one or fewer individual or multiple myeloma-specific risk factors.⁴⁶

In a recent phase III, open-label, multicenter, randomized trial in patients with previously untreated multiple myeloma (n=667) receiving thalidomide-containing regimens, both aspirin (100 mg daily) and fixed-dose warfarin (1.25 mg daily; dose adjustment allowed to maintain INR <3) were similarly effective in reducing thromboembolic events compared with LMWH (enoxaparin 40 mg daily).²⁵³ The primary endpoint was a composite measure including symptomatic DVT, PE, arterial thrombosis, acute cardiovascular events, or sudden otherwise unexplained death during the first 6 months from randomization. The incidence of the composite endpoint was 6.4%, 8.2%, and 5% in the aspirin, warfarin, and LMWH groups, respectively.²⁵³ The absolute risk for the composite endpoint was not statistically different when comparing aspirin with LMWH (absolute difference +1.3%; $P = .544$) or when comparing warfarin with LMWH (absolute difference +3.2%; $P = .183$). Although not statistically significant, LMWH was associated with trends for decreased risks for grade 3 to 4 thromboembolic events and major bleeding events when compared with aspirin. However, LMWH was associated with a significantly decreased risk for grade 3 to 4 thromboembolic events when compared with warfarin (absolute difference +5% for warfarin vs LMWH; $P = .024$). Moreover, among the subgroup of patients aged 65 years or older receiving combination therapy with bortezomib, melphalan, prednisone, and thalidomide, LMWH significantly reduced the risk for the composite endpoint compared with warfarin (absolute difference +11.3 for warfarin vs. LMWH; $P = .006$).²⁵³ It should be noted that this study was conducted in myeloma patients with "standard risk" for thromboembolism, who had no clinical indication for anticoagulation or antiplatelet therapy.

As part of a substudy of a phase III, open-label, randomized trial, thromboprophylaxis with aspirin (100 mg daily) was compared with LMWH (enoxaparin 40 mg daily) in patients with multiple myeloma (n=342) receiving lenalidomide-containing induction (combined with low-dose dexamethasone) and consolidation (combined with melphalan and prednisone).²⁵⁴ The primary endpoint was a composite measure including symptomatic DVT or PE, arterial thrombosis, acute cardiovascular events, or otherwise unexplained sudden death during the first 6 months after randomization. The incidence of the composite endpoint was not statistically different, with 2.3% in the aspirin arm and 1.2% in the LMWH arm. The incidence of DVT was 1.1% and 1.2%, respectively, and the incidence of PE was 1.7% and 0%, respectively. No patients in either treatment arm experienced arterial thrombosis, acute cardiovascular events, or sudden deaths.²⁵⁴ No major bleeding events occurred in either treatment arm; minor bleeding (involving the GI) was reported in 1 patient (<1%) in the LMWH arm. As in the case with the aforementioned phase III study of thromboprophylaxis in patients treated with thalidomide-containing regimens, the current study only included patients who had standard risk for VTE, who had no clear indication or contraindications for antiplatelet or anticoagulation therapy.²⁵⁴ Nevertheless, LMWH appeared to be more effective in preventing PE in this patient population. The investigators from this trial suggested that LMWH was preferred for thromboprophylaxis in patients at high risk for VTE during induction therapy with lenalidomide-containing regimens; in patients with no or only 1 risk factor for VTE, aspirin may be an alternative option. In addition, the investigators concluded that aspirin may also be a feasible thromboprophylaxis option during consolidation or maintenance therapy with lenalidomide.²⁵⁴

In light of the published data from the phase III randomized trials above, the NCCN Panel recommends prophylactic aspirin in multiple myeloma

patients receiving thalidomide or lenalidomide (excluding high-risk combinations) who have no other risk factors for VTE.

With respect to other ambulatory cancer patients, the NCCN Panel suggests risks/benefits conversations regarding the option of thromboprophylaxis in individuals considered to be at high risk for VTE based on an assessment of VTE risk factors (see Guidelines section on *VTE Risk Factors in Cancer Patients*, VTE-A). Some cancer patients undergoing chemotherapy are at increased risk of developing VTE. A predictive model for chemotherapy-associated VTE has been developed²⁶ and independently validated in several studies. The Khorana model considers the following parameters to determine the overall risk for VTE in patients with cancer: site of primary cancer (“very high risk” for stomach or pancreatic cancer; “high risk” for lymphoma, lung, gynecologic, bladder, or testicular cancer), increased pre-chemotherapy platelet count ($\geq 350 \times 10^9/L$), decreased hemoglobin level (< 10 g/dL) or use of ESAs, increased pre-chemotherapy leukocyte count ($> 11 \times 10^9/L$), and high BMI (≥ 35 kg/m²).²⁶ Using a scoring system that assigns risk points to each of the above parameters, patients with 0 points (none of the above risk parameters) are categorized as low risk, those with a total of 1 or 2 points are categorized as intermediate risk, and those with a total score of 3 or higher are considered high risk of developing VTE (see Guidelines section on *VTE Risk Factors in Cancer Patients*, VTE-A 3 of 3). In the original Khorana et al study, the rate of symptomatic VTE in the derivation cohort was 0.8%, 1.8%, and 7.1% for the low-, intermediate-, and high-risk categories, respectively. In the validation cohort, the rates were 0.3%, 2%, and 6.7%, respectively.²⁶ Subsequent independent studies evaluated the utility of the Khorana scoring system in patients with cancer. Retrospective studies in patients with solid tumors and malignant lymphomas reported symptomatic VTE rates of 5% in low-risk, 16% in intermediate-risk, and 27% to 41% in

high-risk patient categories.^{255,256} In a more recent prospective study in patients with cancer (n=819), the rates of symptomatic VTE based on the Khorana scores were 3.8% for low-risk, 9.6% for intermediate-risk, and 17.7% for high-risk patient groups.⁵⁰

Data from a randomized, placebo-controlled, double-blind trial of patients with advanced cancer undergoing treatment with chemotherapy (PROTECHT trial) showed a statistically significant decrease in thromboembolic events (composite endpoint of venous and arterial) in the group receiving prophylactic LMWH (ie, nadroparin) compared with the placebo arm.²⁵⁷ Further, in the randomized CONKO-004 trial, the symptomatic VTE rate of pancreatic cancer patients receiving chemotherapy was significantly reduced at 3 and 12 months with enoxaparin thromboprophylaxis (1mg/kg daily for 3 months followed by 40 mg daily for 3 months) compared with no LMWH.²⁵⁸ Most recently, in a large phase III, randomized, placebo-controlled trial (SAVE-ONCO) in patients with advanced cancer receiving chemotherapy (n=3212), thromboprophylaxis with the investigational ultra-LMWH semuloparin 20 mg daily was compared with placebo.²⁵⁹ The primary efficacy endpoint of this study was a composite endpoint comprising symptomatic DVT, nonfatal or fatal PE, and other death related to VTE. The main safety endpoint was clinically relevant bleeding events. The most common primary cancer sites were lung (37%) and colorectal (29%). Thromboprophylaxis was associated with a significant decrease in the primary endpoint compared with placebo (1.2% vs. 3.4%; hazard ratio, 0.36; 95% CI, 0.21–0.60; $P < .001$).²⁵⁹ The benefit of thromboprophylaxis was observed for both symptomatic DVT (0.7% vs. 2.1%; hazard ratio, 0.32) and nonfatal or fatal PE (0.6% vs. 1.5%; hazard ratio, 0.41). Clinically relevant bleeding (2.8% vs. 2%) and major bleeding events (1.2% vs. 1.1%) with semuloparin versus placebo were not different. Survival outcomes were not significantly different between

study arms, with deaths occurring in 43% and 44.5% of patients in the semuloparin and placebo arms, respectively.²⁵⁹ It should be noted that semuloparin is an investigational agent and has not been approved by the FDA for any indication.

Patients with cancer at high risk for VTE (based on Khorana risk assessment score 3 or higher²⁶) could be considered for outpatient VTE prophylaxis on an individual basis. For these patients, the NCCN Guidelines Panel recommends discussions with patients/caregivers regarding the potential risks and benefits of administering VTE prophylaxis in the outpatient setting. However, thromboprophylaxis in the majority of cancer outpatients receiving chemotherapy is controversial and its broader application using the Khorana risk assessment model or the Vienna risk assessment model should await the results of randomized controlled trials evaluating the efficacy of risk-adjusted thromboprophylaxis based on these models.²⁶⁰

Mechanical Prophylaxis

Intermittent pneumatic compression (IPC) devices and GCS are mechanical prophylaxis options that are principally used in patients with contraindications to pharmacologic prophylaxis or in conjunction with pharmacologic agents in patients at very high risk for VTE. Mechanical prophylaxis should not be used in patients with an acute DVT or in the setting of severe atrial insufficiency (the latter pertains to GCS). In addition, consideration of risks and benefits should be weighed in the presence of large hematomas, thrombocytopenia (platelet count <20,000/mcL), skin ulceration or wounds (which may be more of a concern with GCS), mild arterial insufficiency (which pertains to GCS only), or peripheral neuropathy (which pertains to GCS only; see Guidelines section on *Contraindications to Mechanical Prophylaxis*,

VTE-B). Whenever mechanical prophylaxis is employed, steps should be taken to ensure its proper use and continuous application.

IPC devices have been less well-studied than the use of anticoagulation therapy in VTE prevention.¹⁸² Most of the data on the effectiveness of mechanical prophylaxis have come from surgical populations. For example, in a study comparing the VTE rate in gynecologic oncology surgery patients receiving either low-dose heparin 3 times a day (starting with the day before surgery and continuing for 7 days or longer after surgery) or IPC of the calf, no difference was seen between the 2 modalities.²⁶¹ A retrospective evaluation of high-risk colorectal surgery patients who had received mechanical prophylaxis without anticoagulant therapy indicated that IPC devices were effective in preventing postoperative VTE.²⁶² However, results from a retrospective study of 839 patients over a 2-year period who had undergone abdominal surgery for gynecologic cancers and received pneumatic compression and early ambulation for VTE prophylaxis found that the incidence of PE in cancer patients (4.1%) exceeded by 14-fold the incidence of PE in patients with benign disease (0.3%).²⁴⁴ Therefore, IPC devices should only be used alone for VTE prophylaxis in patients for whom anticoagulant prophylaxis is contraindicated.

GCS have been demonstrated to significantly reduce VTE in comparison to no prophylaxis and provide even greater protection when combined with other preventive therapies.²⁶³ However, many of these studies were conducted more than a decade ago and used fibrinogen uptake scans as a primary outcome measure—a now antiquated diagnostic method. In addition, very few of the patients were noted to have malignancies. Furthermore, a randomized controlled trial in patients undergoing hip surgery found that GCS did not provide significant additive protection against VTE in patients receiving fondaparinux 2.5 mg daily for 5 to 9 days, suggesting that GCS may not



have significant clinical benefits in patients able to receive more potent forms of VTE prophylaxis.²⁶⁴ Similarly, recent results from the CLOTS1 trial, which randomly assigned patients within 1 week of stroke to routine care with or without GCS, found that GCS did not reduce the incidence of DVT in these patients and was associated with a 4-fold increase in the frequency of skin ulcers and necrosis.²⁶⁵ However, the patient group studied in the CLOTS1 trial differs considerably from the patient population described in these guidelines. Furthermore, the long delay in the institution of prophylaxis and the prolonged duration of GCS use (up to 30 days in over 70%) indicate that the safety and efficacy of GCS may be different in different populations studied under different conditions. Therefore, further investigation is warranted.

Until data become available, GCS should not be relied on as the sole method of VTE prophylaxis in cancer patients. Furthermore, cancer patients prescribed GCS for VTE prophylaxis should be carefully monitored for skin complications.

VTE Treatment

Upon diagnosis of VTE, the panel recommends beginning immediate treatment with weight-based intravenous UFH, LMWH, or, in some cases, fondaparinux in cancer patients without contraindications to anticoagulation. Treatment should be at least 5 to 7 days in duration. Since chronic therapy with LMWH is associated with superior outcomes in cancer patients with VTE, its use in the acute phase of treatment may be preferable unless contraindications exist. In the event that warfarin will be used for chronic therapy, there should be a short-term transition phase of at least 5 days during which the acute parenteral anticoagulant (UFH, LMWH, or fondaparinux) is overlapped with warfarin until an INR of 2 or more is achieved. Cancer patients with a DVT or PE should be treated for a minimum duration of 3 months with either an LMWH or

warfarin.^{183,266} LMWH as monotherapy without warfarin is recommended for the first 6 months of chronic treatment of proximal DVT or PE, and for prevention of recurrent VTE in patients with advanced or metastatic cancer who do not have contraindications to anticoagulation (category 1). However, issues such as patient preference and cost should also be considered. Anticoagulation for an indefinite duration should be considered in patients with active cancer or persistent risk factors. Since the chronic treatment of VTE with LMWHs has not been evaluated in clinical trials of cancer patients for durations longer than 6 months, decisions to continue LMWH beyond this time frame or to switch to warfarin therapy for patients requiring longer durations of anticoagulation therapy should be based on clinical judgment.

IVC filter placement should be strongly considered for patients with acute proximal lower-extremity DVT or PE who have absolute contraindications to anticoagulation.^{183,266} However, the benefit of placing an IVC filter in the absence of a lower-extremity IVC or pelvic DVT is unclear. An IVC filter should also be considered in patients with PE where anticoagulation was ineffective (category 2B), patients who are non-adherent with prescribed anticoagulation (category 2B), those with baseline cardiac or pulmonary dysfunction severe enough to make any new or recurrent PE life-threatening (category 2B), and those with documented multiple PE and chronic thromboembolic pulmonary hypertension (category 2B).

In general, a retrievable IVC filter is preferred in most clinical situations; permanent filters should only be considered in rare situations in which patients have permanent contraindications to anticoagulation or chronic comorbidities that preclude the use of anticoagulants. When a retrievable filter is placed, it is imperative that patients be followed closely by their physicians so that the device can be removed in a timely fashion after the need for its placement is no longer present.

Improvements in technology and an increase in the number of available thrombolytic agents have increased the use of thrombolytic therapy for DVT. Anticoagulation prevents clot extension and recurrence, but does not actively dissolve clot. In contrast, thrombolytic agents promote clot dissolution, which may help to limit long-term complications such as post-thrombotic syndrome (PTS). PTS is a chronic complication of DVT that develops over months to years after the thrombotic episode. PTS is caused by chronic venous hypertension that results from impaired venous outflow from the affected limb due to vascular obstruction and venous valvular dysfunction. Thrombolytic agents theoretically may reduce the incidence of PTS by promoting rapid clot lysis, reducing venous outflow obstruction, and preventing venous valvular damage. Typical symptoms and signs of PTS include leg pain, heaviness, or swelling of the leg.²⁶⁷ The syndrome has been reported to occur in approximately 30% to 50% of patients within 5 to 8 years following symptomatic DVT, and can negatively affect a patient's quality of life.^{268,269} Severe forms of PTS can occur in up to 10% of patients, and may involve skin and subcutaneous tissue changes, such as skin ulcerations, hyperpigmentation, varicose eczema, and subcutaneous atrophy.²⁶⁷

Thrombolytics that have been used in the management of DVT include urokinase, streptokinase, and, more recently, recombinant plasminogen activators alteplase, reteplase, and tenecteplase given intravenously. In the past, thrombolytic agents were delivered systemically through an intravenous catheter, which likely reduced the efficacy of the therapy and increased the likelihood of bleeding complications. Nevertheless, thrombolysis was associated with increased rates of major or complete clot lysis and fewer post-thrombotic complications, compared with anticoagulation alone.²⁷⁰⁻²⁷⁵ In recent years, catheter-directed delivery of thrombolytic agents directly into the substance of the clot has allowed

more localized targeting of thrombolytic agents and the employment of catheter-based thrombectomy devices to accelerate clot removal. Catheter-directed thrombolysis (CDT) with or without mechanical thrombectomy is associated with significantly higher rates of complete clot lysis than conventional anticoagulation.²⁷⁶ Effective clot lysis in patients with DVT has been reported with CDT using urokinase, alteplase, reteplase, and tenecteplase,²⁷⁷⁻²⁷⁹ including a retrospective analysis that suggested potentially lower treatment costs associated with plasminogen activators (alteplase and reteplase) compared with urokinase.²⁷⁸

Initial results from an open-label, randomized, controlled trial comparing CDT with alteplase added to anticoagulation versus anticoagulation alone in patients with acute iliofemoral DVT (n=103) reported a higher rate of iliofemoral patency at 6 months with the addition of CDT (64% vs. 36%).²⁸⁰ Long-term follow-up from this study with larger patient numbers (n=209) confirmed the higher rate of iliofemoral patency at 6 months (66% vs. 47%) with the addition of CDT.²⁸¹ After completion of 24 months of follow-up, PTS was reported in significantly fewer patients in the CDT arm (41% vs. 56%; $P = .047$). In contrast, ECS did not prevent PTS compared to placebo in a randomized trial (SOX Trial) of patients who experienced a first proximal DVT.²⁸² Therefore, GCS should not be prescribed for prevention of PTS.

Retrospective patient series have demonstrated that cancer patients can benefit from catheter-directed pharmacomechanical thrombolysis.²⁸³ The 2012 ACCP Guidelines do not recommend routine use of CDT over anticoagulation alone, but suggest that patients with the following factors are most likely to benefit from CDT: iliofemoral DVT; symptom duration less than 14 days; good functional status; life expectancy of at least 1 year; and low risk of bleeding.¹⁸³ The NCCN Panel believes that CDT and thrombectomy can be considered a therapeutic option for

select patients with large symptomatic extremity DVT, particularly when they are not responding to conventional anticoagulation.²⁶⁶ Absolute contraindications to thrombolysis (administered locally or systemically) include history of hemorrhagic stroke (or stroke of unknown origin), intracranial tumor, ischemic stroke (in previous 3 months), history of major trauma, surgery or head injury in previous 3 weeks, low platelet count ($<100 \times 10^9/L$), active bleeding, and bleeding diathesis. Relative contraindications to thrombolysis include age greater than 75 years, pregnancy or first postpartum week, non-compressible puncture sites, traumatic resuscitation, refractory hypertension, advanced liver disease, infective endocarditis, recent GI bleeding within 3 months, and life expectancy of 1 year or less.¹⁸³ Selection of thrombolytic agents and thrombectomy devices should be made based on local expertise and experience. Broader use of CDT awaits the outcome of currently active clinical trials.

Treatment of patients with an incidental VTE following radiographic detection should be the same as for patients with symptomatic VTE.^{135,284,285}

Immediate VTE Treatment

In a recent meta-analysis of trials comparing outcomes with anticoagulants (UFH, LMWH, and fondaparinux) as initial treatment of VTE in cancer patients, LMWH was associated with a significant reduction in mortality rate at 3-month follow-up compared with UFH (relative risk, 0.71; 95% CI, 0.52–0.98).²⁸⁶ However, no significant difference was found in VTE recurrence between LMWH and UFH. No statistically significant differences were found between heparin and fondaparinux in terms of mortality, VTE recurrence, or bleeding events.²⁸⁶ In the absence of contraindications to their use, LMWHs are preferred for acute management of VTE in cancer patients because

they do not require hospitalization or monitoring, and are the preferred option for long term therapy.

Chronic VTE Treatment

Several studies comparing the efficacy and safety of LMWH and oral warfarin in the chronic treatment of VTE in patients with cancer have been performed. In a randomized open-label trial (CANTHANOX trial), the use of chronic (3 months) enoxaparin (1.5 mg/kg every 24 hours) versus chronic warfarin (INR 2–3) was evaluated after immediate treatment with either LMWH or UFH in 146 cancer patients with VTE.¹⁶⁹ The primary endpoint of this study was a combined outcome event including major bleeding and recurrent VTE within 3 months. In the groups receiving chronic enoxaparin and warfarin, 10.5% and 21.1% of patients, respectively, experienced either major bleeding or recurrent VTE ($P = .09$); fatal bleeding occurred in 0% and 8% of patients, respectively ($P = .03$). In another study, no significant differences in bleeding or recurrent VTE were observed when patients with active cancer and acute VTE were randomly assigned to receive either 6 months of enoxaparin (either 1.5 mg/kg or 1 mg/kg every 24 hours) or immediate enoxaparin therapy followed by warfarin to complete 6 months of therapy (ONCENOX trial).²⁸⁷

The CLOT trial compared the efficacy and safety of immediate dalteparin (200 units/kg daily for 5–7 days) followed by chronic (6 months) therapy with an oral coumarin derivative versus chronic dalteparin therapy (200 units/kg daily for one month followed by 150 units/kg for months 2–6) in patients with cancer (most had metastatic disease) after diagnosis of acute proximal DVT, PE, or both.¹⁶⁸ The Kaplan-Meier estimate for recurrence of VTE over the 6-month study period showed significantly decreased risks with dalteparin compared with oral anticoagulants (hazard ratio, 0.48; $P = .002$). This study

showed probabilities of recurrent VTE at 6 months of 9% and 17% in cancer patients receiving dalteparin or oral anticoagulants, respectively. No significant difference in bleeding rates was seen for the 2 groups.¹⁶⁸ The results of this study support use of LMWHs as chronic anticoagulation therapy in patients with metastatic disease who are diagnosed with acute VTE.

Some limitations of the CLOT study include the lack of patients with below-the-knee or catheter-related thrombosis; a study duration of only 6 months, for which the apparent efficacy difference was observed for development of recurrent DVT only (but not for PE, although the study was not designed to assess differences in outcomes according to type of VTE); and uncertainty as to whether these results can be extrapolated to LMWHs other than dalteparin. A Cochrane review of anticoagulation for the chronic treatment of VTE in patients with cancer combined the results of all these studies and found no significant differences in bleeding, thrombocytopenia, or survival outcomes with use of LMWH compared with oral vitamin K antagonists.²⁸⁸ However, the incidence of VTE was significantly lower for patients receiving LMWH (hazard ratio, 0.47; 95% CI, 0.32–0.71).

In a recent open-label, randomized, phase III trial in patients with acute PE with or without DVT (n=4832), treatment with the oral factor Xa inhibitor rivaroxaban (15 mg twice daily for the first 3 weeks, then 20 mg once daily thereafter) was compared with standard therapy with LMWH and vitamin K antagonist (enoxaparin 1 mg/kg twice daily for at least 5 days, and vitamin K antagonist adjusted to INR 2.0–3.0) for initial and chronic treatment (3, 6, or 12 months).²⁰⁶ This study was designed as a non-inferiority trial. The primary efficacy endpoint was symptomatic recurrent VTE, defined as a composite endpoint comprising fatal or nonfatal PE or DVT. The main safety endpoint was clinically relevant bleeding. Based on the primary endpoint of recurrent VTE, rivaroxaban

was found to be non-inferior to standard therapy (2.1% vs. 1.8%; hazard ratio, 1.12; 95% CI, 0.75–1.68).²⁰⁶ The incidence of clinically relevant bleeding events was not significantly different between rivaroxaban and standard therapy (10.3% vs. 11.4%; hazard ratio, 0.90; 95% CI, 0.76–1.07). However, rivaroxaban was associated with a significantly decreased incidence of major bleeding (1.1% vs. 2.2%; hazard ratio, 0.49; 95% CI, 0.31–0.79; $P = .003$).²⁰⁶ This randomized study suggested that initial and chronic treatment of PE with fixed-dose rivaroxaban was non-inferior to standard anticoagulation therapy. It should be noted that less than 5% of the study patients in this study had active cancer.

Apixaban is another oral factor Xa inhibitor that has also been evaluated for extended anticoagulation therapy in patients who had a VTE. In a recent double-blind, randomized, phase III trial in patients who had completed 6 to 12 months of anticoagulation for VTE (n=2486), extended treatment with apixaban (at a dose of 2.5 mg or 5 mg twice daily) was compared with placebo given for 12 months.²¹³ The primary outcome measure was a composite endpoint comprising symptomatic recurrent VTE (fatal or nonfatal PE or DVT) and death from any cause. During the 12-month treatment period, the primary endpoint occurred in 3.8% of patients in the 2.5 mg apixaban arm and 4.2% in 5 mg apixaban arm compared with 11.6% in the placebo arm ($P < .001$ for both comparisons). The rate of symptomatic recurrent VTE and death from VTE was also significantly reduced with apixaban 2.5 mg (1.7%) or apixaban 5 mg (1.7%) compared with placebo (8.8%; $P < .001$ for both comparisons).²¹³ The rate of major bleeding was 0.2% in the 2.5 mg apixaban arm, 0.1% in the 5 mg apixaban arm, and 0.5% in the placebo arm. This study showed that extended anticoagulation with apixaban reduced the risk for recurrent VTE without increasing the risk for major bleeding. It should be noted, however, that less than 2% of patients on this study had active cancer.²¹³ Thus, the use of apixaban



for acute and extended chronic treatment of VTE in patients with cancer remains to be investigated in future prospective trials.

Increased survival rates have been reported for subgroups of cancer patients receiving chronic treatment with LMWH versus other VTE therapies or placebo.^{289,290} For example, although no survival differences were seen in groups of patients with advanced cancer without VTE receiving either dalteparin or placebo in the FAMOUS study, results from a subgroup analysis of patients with better prognoses (more indolent disease and survival beyond 17 months post-randomization) suggested that 2- and 3-year survival rates were higher for patients receiving dalteparin compared with patients receiving placebo.²⁰⁰ A post hoc analysis of patients from the CLOT study also indicated that no differences in 1-year survival were seen between groups of patients with metastatic disease receiving either long-term dalteparin or oral coumarin derivatives, whereas 1-year survival rates were higher in the subgroup of patients without metastases receiving dalteparin when compared with patients in the same subgroup receiving an oral vitamin K antagonist.²⁹⁰ Results of other randomized studies have also provided evidence of improvement in median progression-free survival and/or overall survival of cancer patients receiving LMWHs.^{291,292} In addition, a Cochrane review assessing the antineoplastic properties of anticoagulants found that heparins appear to improve the survival of cancer patients with limited-stage disease and that further research is warranted to identify the most effective regimens and most responsive cancer patient populations.²⁹³ Additional evaluations of the putative anti-tumor effects of LMWHs are needed before recommendations pertaining to their use as antineoplastic agents can be made.

Treatment of Catheter-related DVT

The central tenant guiding the treatment of catheter-related DVT is based on the question of whether the device is required for continued treatment of the patient. Device removal is recommended in the case of catheter-related DVT when the device is no longer required, or when it is required but contraindications to anticoagulation exist. If device removal is planned, some have recommended a short period of anticoagulation of 5 to 7 days, if feasible, to reduce the chances of clot embolization with device removal. An assessment of the likelihood and consequences of clot embolization based on the size and position of the device-associated thrombus should be conducted prior to removal. Anticoagulation therapy is recommended while the catheter is in place (in the absence of contraindications) and for a total duration of therapy of at least 3 months, or for as long as the catheter remains in place, whichever is longer. Consider removal of the catheter if DVT symptoms persist or if the catheter is infected, dysfunctional, or no longer necessary. Patients with catheter-related DVT and contraindications to anticoagulation therapy should be followed for changes in these contraindications as clinically indicated; anticoagulation therapy is recommended after contraindications are no longer present.

No randomized controlled trials have been reported evaluating the effects of particular therapeutic strategies on outcomes of catheter-related VTE. A prospective study of 444 cancer patients with CVAD showed an incidence of symptomatic catheter-related thrombosis of 4.3%.⁷⁵ Of 19 patients with catheter-related thrombosis, 9 were treated with anticoagulation therapy only, 8 patients underwent anticoagulation therapy and catheter removal, 1 patient was treated with catheter removal only, and 1 patient did not receive any treatment. The duration of anticoagulation therapy was not specified, but evaluation of the 15 patients alive at 24 weeks after diagnosis of catheter-related thrombosis



revealed that residual symptoms were present in only 2 patients. A more recent pilot study of cancer patients with catheter-related, symptomatic UEDVT demonstrated that anticoagulation with dalteparin followed by warfarin (INR 2–3) was not associated with episodes of recurrent VTE and/or line removal as a consequence of thrombosis/infusion failure; major bleeding occurred in 3 patients (4%).²⁹⁴

Treatment of SVT

Anticoagulation therapy with intravenous UFH or a LMWH for at least 6 weeks is a category 2B recommendation for patients with a non-peripheral catheter-related SVT in close proximity to the deep venous system. Consider treating up to 12 weeks if SVT is in close proximity to the common femoral vein. Since migratory superficial thrombophlebitis is a characteristic presentation for Trousseau's syndrome, a heightened awareness of this cancer-associated hypercoagulable state is warranted as indefinite therapy with UFH or LMWH is essential for its treatment.

Catheter removal is recommended for a peripheral catheter-related SVT. Anti-inflammatory medications, warm compresses, and elevation of the affected limb should be employed as clinically indicated. These strategies are also recommended for the initial treatment of SVT that is not associated with a peripheral catheter. Aspirin and nonsteroidal anti-inflammatory drugs (NSAIDs) should be avoided in patients with platelet counts less than 20,000 to 50,000/mcL or with severe platelet dysfunction. Anti-inflammatory agents are recommended for the symptomatic treatment of certain types of SVT only, not for DVT prophylaxis. Only a limited number of studies have evaluated the clinical significance of SVT, its associated progression to VTE, and the effect of anticoagulant agents on its course.^{295,296} In a large observational study

of patients (n=844) with symptomatic SVT (at least 5 cm), 66% had SVT of the greater saphenous vein, and in 20% of these patients the median distance between the thrombus and the saphenofemoral junction was less than or equal to 3 cm.⁹⁶ In this study, 25% of patients had DVT or PE at inclusion, and 10% of the patients without VTE at study inclusion (isolated SVT only) who were available at 3-month follow-up subsequently developed thromboembolic complications (eg, PE, DVT, extension of SVT) despite the use of anticoagulation therapy in about 90% of these individuals⁹⁶; a possible limitation of this study is that all of these patients were evaluated in a specialist referral setting. In a prospective assessment of 60 consecutive patients with SVT of the greater saphenous vein, the combined incidence of DVT and SVT events over a 6-month follow-up period was lower in patients treated with twice-daily subcutaneous injections of high-dose heparin (12,500 IU for 1 week, followed by 10,000 IU for 4 weeks) when compared with patients receiving 4 weeks of low-dose (5000 IU) heparin (3% vs. 20%; $P = .05$).²⁹⁷ A pilot study evaluating the effects of once-daily administration of an LMWH, an NSAID, or placebo for 8 to 12 days on the clinical course of SVT showed no significant differences between treatment and placebo groups with respect to progression to DVT.²⁹⁸ However, all active treatments reduced the combined rate of DVT and SVT compared with placebo, although no significant differences were observed between active treatment groups.²⁹⁸ This finding possibly indicates that longer treatment durations may be required.

In a placebo-controlled, randomized, double-blind study (CALISTO), Decousus and colleagues randomized 3002 patients with acute symptomatic superficial thrombophlebitis of the lower extremities to fondaparinux 2.5 mg daily (n=1502) or placebo (n=1500) for 45 days. Fondaparinux was associated with a significant reduction in the primary outcome (all-cause mortality, symptomatic VTE, or extension of the



superficial thrombophlebitis to involve the saphenofemoral junction or symptomatic recurrence of SVT) compared with placebo (0.9% vs. 5.9%; relative risk reduction [RRR], 85%; 95% CI, 0.74–0.92; $P < .0001$).²⁹⁹ Fondaparinux 2.5 mg daily significantly reduced the incidence of DVT and/or PE (0.2% vs. 1.3%; RRR, 85%; 95% CI, 0.50–0.95; $P < .001$). The rate of major bleeding was similar (1 patient in each group). The trial establishes the efficacy of prophylactic dose fondaparinux in the treatment of SVT. However, it is important to note that patients with active cancer and SVT within 3 cm of the saphenofemoral junction were excluded.²⁹⁹

Treatment of SPVT

The management of patients with SPVT encompasses the use of anticoagulation therapy with or without invasive procedures, such as CDT, (TIPS), surgical shunting, or surgical resection of bowel, as well as other medical management, such as the use of a β -blocker. Management depends on the extent and location of the thrombus, presence of acute symptoms of intestinal infarction, and signs of portal cavernoma or portal hypertension. In the absence of contraindications, anticoagulation with unfractionated heparin or LMWH (preferred) should be initiated, followed by oral anticoagulation for at least 6 months in the case of triggered thrombotic events, such as in a postsurgical setting.^{106,108,114,115} The benefit of anticoagulation as initial and long-term therapy in patients with SPVT has been reported in several studies.^{102,117,300,301} In a long-term follow-up study of patients with SPVT ($n=95$; median follow-up of 41 months) primarily treated with anticoagulation (LMWH 200 IU/kg per day for 7–10 days followed by oral anticoagulation for 6 months), 45% of patients with acute SPVT ($n=21$) had complete recanalization with anticoagulants.¹⁰² Patients requiring resection for intestinal infarction or having incomplete recanalization of thrombus or having inherited thrombophilia were given

lifelong oral anticoagulation in this study. Recurrent VTE occurred in 18.5% of patients overall, and was significantly more frequent among patients with concurrent myeloproliferative disorders at presentation versus those without such disorders (70% vs. 13%; $P < .0001$). Moreover, recurrent VTE was only observed among patients who did not receive anticoagulation.¹⁰² Gastrointestinal (GI) bleeding occurred in 15% of patients and was significantly more frequent among patients with bleeding from esophageal varices at presentation compared with those without prior bleeding (57% vs. 5%; $P < .0001$). None of the patients receiving oral anticoagulation had bleeding events.¹⁰² In a recent prospective multicenter study in patients with acute portal vein thrombosis ($n=95$) treated with anticoagulation (initial therapy with heparin followed by oral anticoagulation targeting INR 2–3 for 6 months or long-term in patients with permanent prothrombotic disorders or obstruction of mesenteric vein), the 1-year recanalization rate in the portal vein was 38%.¹¹⁷ The 1-year recanalization rates in the mesenteric and splenic veins were 61% and 54%, respectively. GI bleeding occurred in 9% of patients, none of which were fatal events.¹¹⁷ Anticoagulation appears to lower the risk for recurrent thrombosis in patients with SPVT without increasing the risks for severe bleeding,^{102,117,300,301} including in patients with underlying prothrombotic states.³⁰⁰ However, a recent retrospective study in a large cohort of patients with SPVT ($n=832$) showed that the rate of recurrent VTE was not significantly improved with oral anticoagulation with warfarin (10-year recurrence-free survival rate was 89% vs. 77% in patients who did not receive anticoagulation; $P = .38$).¹⁰¹ Based on multivariate analysis, hormone therapy was the only independent predictor of recurrence. Major bleeding events were reported more frequently among patients receiving anticoagulation compared with those who did not (26% vs. 19%; $P < .05$). Moreover, based on multivariate analysis, presence of gastroesophageal varices and anticoagulation were independent



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

predictors for bleeding events.¹⁰¹ In chronic SPVT, the presence of portal hypertension may increase the risk of bleeding from esophageal varices and splenomegaly may lead to decreased platelet counts, which can further increase the risks of bleeding events in patients treated with anticoagulation.¹⁰⁰ Thus, in the absence of randomized controlled trials, the issue of long-term or lifelong anticoagulation remains somewhat controversial in patients with SPVT. An individual patient's risk factor(s) for SPVT should be taken into consideration when weighing the risks and benefits of long-term anticoagulation. The panel currently recommends lifelong anticoagulation in patients with active cancer, underlying thrombophilia, and/or idiopathic thrombosis.

In patients with acute SPVT with clinical deterioration or progression of thrombosis despite anticoagulation, more invasive approaches using CDT, TIPS, or surgical shunting may be required.^{106,108,115} Acute thrombosis involving the mesenteric veins is associated with high risks of intestinal infarction, which is life-threatening and requires immediate surgery to resect necrotic sections of the bowel.^{102,104,108,114} Catheter-directed thrombolytic therapy has been reported to have some success in acute SPVT in small retrospective studies.³⁰²⁻³⁰⁵ Thrombolytic therapy may be most suitable when administered locally for patients with recent thrombosis^{115,304}; however, this approach should be used with caution due to risks for major bleeding complications.^{114,115,302,305} The decision for administering thrombolytic therapy should be based on availability and expertise at the local institution, the location of the thrombus, and evaluation of risks of bleeding in individual patients. In addition, the selection of regimens should be based on institutional experience, with decisions made in conjunction with specialists in interventional radiology and vascular surgery. For patients with acute hepatic vein thrombosis with contraindications to anticoagulation or for patients with chronic hepatic vein thrombosis for whom medical management alone are

unsuccessful, TIPS or surgical shunts may be considered. TIPS is an interventional radiologic procedure that creates a portocaval shunt between the hepatic and portal veins, and may be appropriate for patients with an occluded IVC or a portacaval pressure gradient <10 mm Hg.^{115,306} TIPS may also be appropriate for patients with refractory ascites and progressive hepatic dysfunction despite medical management and/or interventions to achieve recanalization.^{306,307} This procedure is less invasive than surgical interventions, and has been successful in reducing portal hypertension, resolving ascites, and improving hepatic function in patients with Budd-Chiari syndrome.³⁰⁶⁻³¹¹ Although shunt dysfunction or stenosis is common during follow-up, TIPS is associated with promising long-term outcomes with 5-year transplant-free survival rates of 74% to 78% in recent studies.^{306,311} Surgical portosystemic shunts may be appropriate in patients without an occluded IVC, with a portacaval pressure gradient >10 mm Hg, and with preservation of hepatic function.^{115,312} The impact of surgical shunts versus other interventions on long-term outcomes is unknown³¹³; nevertheless, 5-year survival rates range from 75% to 87% in patients with Budd-Chiari syndrome undergoing successful surgical portosystemic shunts,³¹⁴⁻³¹⁶ and this procedure may improve survival outcomes in patients with intermediate-risk prognostic factors as defined by Darwish Murad et al.³¹⁷ Of note, surgical shunts appear to have largely been replaced with TIPS in recent years.³⁰⁷

Patients with chronic portal or mesenteric vein thrombosis frequently present with cavernous transformation and/or signs of portal hypertension, the latter of which can lead to complications such as variceal bleeding.¹¹⁶ Gastroesophageal varices may be seen in 35% to 50% of patients with portal vein thrombosis at presentation,^{101,114} and remain a significant independent risk factor for major bleeding in patients with SPVT.¹⁰¹ Thus, an important goal in the management of

patients with chronic portal or mesenteric thrombosis is risk reduction for and prevention of bleeding events.^{108,114} The use of β -blockers and endoscopic treatments has been evaluated in the primary and secondary prophylaxis for variceal bleeding in patients at high risk of bleeding events. In several prospective randomized studies comparing the use of variceal banding ligation versus propranolol for primary prophylaxis of variceal bleeding in cirrhotic patients presenting with high-risk gastroesophageal varices, the treatment methods were similarly effective in preventing variceal bleeding (which occurred in 12%–25% of patients with ligation versus 24%–29% receiving propranolol), with a similar overall mortality rate.³¹⁸⁻³²⁰ In one of the studies, patients (n=75) treated with variceal banding ligation had a significantly decreased incidence of esophageal variceal bleeding compared with patients receiving propranolol (5% vs. 25%; $P = .027$), but at the expense of a higher incidence of subcardial variceal bleeding (8% vs. 0%; $P = .027$).³¹⁸ In another prospective randomized trial comparing the effectiveness of primary prophylaxis using these methods in cirrhotic patients (n=60), ligation was reported to be more effective than propranolol in preventing variceal bleeding (which occurred in 7% vs. 30%; $P = .043$).³²¹ A large randomized study comparing variceal banding ligation with or without propranolol for primary prophylaxis of variceal bleeding in patients with high-risk varices (n=144) showed that the combined modality did not significantly reduce the risks of bleeding (actuarial probability, 7% vs. 11%; $P = .72$) or death (actuarial probability, 8% vs. 15%; $P = .37$) at 20 months compared with ligation alone.³²² The use of variceal banding ligation and propranolol has also been evaluated in the secondary prophylaxis setting in patients with noncirrhotic portal hypertension at risk of recurrent variceal bleeding. In a recent study (n=101), the incidence of recurrent variceal bleeding was found to be similar between patients receiving ligation versus propranolol (24% vs. 18%; $P = .625$) for

prevention of recurrent bleeding.³²³ However, a recent meta-analysis of randomized studies demonstrated that variceal banding ligation or sclerotherapy combined with β -blockers was significantly more effective than endoscopic treatment alone in preventing overall recurrent bleeding (OR, 2.20; 95% CI, 1.69–2.85; $P < .0001$) and in decreasing overall mortality (OR, 1.43; 95% CI, 1.03–1.98; $P = .03$), suggesting that combined modality may be preferred as secondary prophylaxis for esophageal variceal bleeding.³²⁴ The panel recommends initiation of β -blockers in patients with chronic portal or mesenteric thrombosis presenting with gastroesophageal varices with or without signs of portal hypertension. In patients who had prior variceal bleeding, it may be appropriate to consider variceal banding ligation or sclerotherapy in conjunction with β -blockers.

Treatment of PE

Once a diagnosis of PE is made, the panel recommends that patients be risk-stratified to determine the advisability of outpatient management and intensity of initial follow-up and treatment.^{145,146} Anticoagulation therapy is recommended for all patients with acute PE who do not have a contraindication to such therapy.¹⁸³ In patients with a contraindication to anticoagulation, an IVC filter should be strongly considered if PE is due to lower extremity, pelvic, or abdominal DVT and the patient should be closely followed for a change in clinical status that would allow anticoagulation to be instituted.^{183,266}

In patients with submassive PE and evidence of moderate or severe RV enlargement or dysfunction, thrombolytic therapy is a therapeutic consideration.^{132,266,325} In patients without contraindications to anticoagulation, immediate anticoagulation therapy should be started at PE diagnosis; evaluation of risk should be performed concurrently with PE diagnosis or as soon as relevant data are available. After

assessment of the cancer status of the high-risk patient with PE, the physician should consider the use of thrombolytic therapy and/or pulmonary embolectomy after weighing the severity of the patient's illness and his or her risk of bleeding. Although IVC filters are typically reserved for patients with a contraindication to anticoagulation, IVC filters are occasionally placed in patients with severely compromised cardiopulmonary status. In either case, a retrievable filter with a wide window of retrievability is preferred over a permanent one to allow subsequent filter retrieval once the patient's cardiopulmonary status has stabilized. Patients should be followed closely and their filters should be retrieved once they are stable on therapeutic anticoagulation. Permanent filters should only be considered for rare patients with chronic comorbidities or with permanent contraindication to anticoagulation.

In the randomized placebo-controlled MAPPET-3 trial of hemodynamically stable patients with submassive acute PE and pulmonary hypertension or evidence of RV dysfunction who received heparin in conjunction with thrombolysis with alteplase or heparin plus placebo, addition of thrombolysis was associated with significantly decreased incidence of in-hospital mortality and clinical deterioration requiring treatment escalation (primary endpoint; 11% vs. 25%; $P = .006$). This difference was due to a higher incidence of clinical instability in the placebo group, as in-hospital mortality rates were similar between treatment groups.³²⁶ The clinical endpoints and other aspects of the design of this trial have been criticized.^{327,328} The recently published PEITHO study found similar results. The investigators randomized 1005 patients with intermediate risk submassive PE to tenecteplase plus heparin or placebo plus heparin. Death or hemodynamic decompensation occurred in 13 of 506 patients (2.6%) in the tenecteplase group as compared with 28 of 499 (5.6%) in the placebo

group (odds ratio, 0.44; 95% confidence interval, 0.23 to 0.87; $P = .02$). There was no difference in death between the two groups (6 tenecteplase patients (1.2%) versus 9 placebo patients (1.8%)($P = .42$). Extracranial bleeding occurred in 32 patients (6.3%) in the tenecteplase group and 6 patients (1.2%) in the placebo group ($P < .001$). Stroke occurred in 12 patients (2.4%) in the tenecteplase group (hemorrhagic in 10) and 1 patient (0.2%) in the placebo group ($P = .003$). These data indicate that tenecteplase can reduce the incidence of hemodynamic decompensation in patients with intermediate risk PE but at the cost of increased extracranial bleeding and stroke and no improvement in mortality. Seventy-three patients (7.3%) on the PEITHO study had active cancer.

In a meta-analysis of 16 trials that randomized 2115 patients with PE to thrombolytic therapy or anticoagulation, use of thrombolytic therapy was associated with lower all-cause mortality (OR, 0.53; 95% CI, 0.32–0.88) and higher risk of major bleeding (OR, 2.73; 95% CI, 1.91–3.91).³²⁹ This analysis included 8 trials involving patients with intermediate-risk PE (hemodynamically stable patients with right ventricular dysfunction). Results from previous meta-analyses reported no significant benefit with thrombolytic therapy compared with heparin alone in terms of recurrent PE or death, particularly for patients with hemodynamically stable PE.³³⁰⁻³³⁴ Reports from several studies evaluating the use of pulmonary embolectomy in patients with acute PE provide support for the use of this procedure in patients with hemodynamically stable or unstable acute PE characterized by RV dysfunction.³³⁵⁻³³⁷ An important consideration for these guidelines is that none of these studies evaluating the use of thrombolytic therapy or surgical embolectomy to treat patients with acute PE specifically address treating cancer patients. However, no significant difference in bleeding risk was observed in a recent retrospective consecutive case series comparing

the safety of percutaneous CDT for upper- or lower-extremity acute symptomatic DVT in patients with or without cancer.²⁸³

The ACCP recommends against the use of thrombolytic therapy or pulmonary embolectomy in most patients with PE.¹⁸³ Use of thrombolytic therapy in selected patients, such as those with PE associated with hypotension or hemodynamic instability, and without a high risk of bleeding, is recommended.^{183,266} Absolute contraindications to thrombolysis (administered locally or systemically) include a history of hemorrhagic stroke (or stroke of unknown origin), intracranial tumor, or ischemic stroke in previous 3 months, a history of major trauma, surgery or head injury in the previous 3 weeks, thrombocytopenia (platelet count $<100 \times 10^9/L$), active bleeding, and a bleeding diathesis. Relative contraindications to thrombolysis include older age (>75 years), pregnancy or first postpartum week, non-compressible puncture sites, traumatic resuscitation, refractory hypertension, advanced liver disease, infective endocarditis, recent GI bleeding in the last 3 months, and a life expectancy of 1 year or less.¹⁸³ Catheter or surgical embolectomy may be considered in patients with massive PE who have contraindications to thrombolytic therapy or those who remain unstable following thrombolysis.^{183,266} Selection of thrombolytic agents and thrombectomy devices should be made based on local expertise and experience.

VTE Therapies: Response Assessment

Intensive monitoring of the antithrombotic effects of some anticoagulants is particularly important in patients with cancer.¹⁹² The recommendations on monitoring anticoagulant response included in the NCCN Guidelines for VTE may be superseded by written standard procedures specific to an institution.

Unfractionated Heparin

Heparins indirectly affect the coagulation system by potentiating antithrombin activity, thereby facilitating inhibition of thrombin, factor Xa, and, to a lesser extent, several other activated coagulation factors.^{181,338}

The aPTT measures the overall activity of the intrinsic and common coagulation pathways and is particularly sensitive to agents that inhibit thrombin.^{191,339} Therefore, UFH is most commonly monitored during the treatment of VTE with the aPTT and depends on the establishment of a therapeutic aPTT range.^{214,340} The aPTT therapeutic range should be established by each institution using regular calibration of the aPTT therapeutic range against UFH levels of 0.3 to 0.7 units/mL (as determined by factor Xa inhibition using a chromogenic assay) or 0.2 to 0.4 units/mL (as determined by protamine sulfate titration) as recommended by the College of American Pathologists (CAP) and ACCP.^{181,340,341} Such testing should be performed in the clinical laboratories at each institution according to an institutional SOP, and the aPTT therapeutic range should be printed on the laboratory report. In the event that this information is unavailable, a fixed aPTT therapeutic range of 2 to 2.5 times the baseline aPTT for the patient is recommended by the panel to monitor UFH dosing. Monitoring is generally not performed in patients receiving prophylactic doses of subcutaneous UFH.³³⁸

LMWHs and Fondaparinux

LMWHs act by potentiating the inhibitory activity of antithrombin against factor Xa and, to a lesser extent, thrombin.¹⁸¹ Fondaparinux is a synthetic indirect factor Xa inhibitor that also functions through potentiation of antithrombin activity.¹⁵⁹ Measurement of factor Xa inhibition, not the aPTT, is necessary to monitor the anticoagulant effect of LMWH or fondaparinux, because thrombin inhibition associated with LMWH or fondaparinux is weak or absent, respectively.^{159,181} However,



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

only limited data are available on the use of factor Xa inhibition to monitor and adjust LMWH or fondaparinux therapy, and monitoring of patients receiving LMWH or fondaparinux is generally not performed because of the more predictable dose response associated with these agents.^{181,200} As previously discussed, LMWH should be used with caution in patients with renal dysfunction; the prescribing information for specific agents should be followed for renal dysfunction and body-weight dosing. Dose adjustments should be considered in patients with severe renal dysfunction ($C_{cr} < 30$ mL/min). LMWH anti-Xa monitoring (peak and trough) has been recommended for patients with severe renal dysfunction, although only limited data are available to support the clinical relevance of anti-Xa levels.^{181,194,342} A meta-analysis of studies in patients treated with LMWH (enoxaparin in the large majority of studies) showed that anti-Xa levels were significantly elevated among patients with severe renal dysfunction ($C_{cr} \leq 30$ mL/min) compared with those with $C_{cr} > 30$ mL/min, and that risks of major bleeding were increased in the former subgroup.¹⁹⁴ These effects were found when therapeutic doses of enoxaparin were used, but not with prophylactic doses. If anti-Xa levels are monitored in LMWH-treated patients, it is recommended that chromogenic methods be used.³⁴² In general, the panel recommends limiting the use of LMWHs and fondaparinux in patients with renal insufficiency and those at extremes of body weight, rather than close monitoring. Panel opinions diverged on the utility of measuring factor Xa inhibition in certain cases, such as in patients with very high body weight (>150 kg) receiving LMWH for an extended period of time.

Direct Thrombin Inhibitors

Argatroban and bivalirudin are parenteral DTIs that do not require antithrombin for anticoagulant activity. Therefore, the anticoagulant effect of these agents can be measured using the aPTT, although

results can be affected by the specific DTI and the aPTT assay reagents used.³³⁸ Target aPTT ratio ranges of 1.5 to 3 times control and 1.5 to 2.5 times control are recommended when using argatroban and bivalirudin, respectively (see Guidelines section on Therapeutic Options for *Heparin-Induced Thrombocytopenia [HIT]*). As argatroban is metabolized in the liver,³⁴³ significant dose reductions are necessary in patients with impaired liver function; argatroban should be avoided in patients with severely impaired hepatic function.

A third thrombin inhibitor, desirudin, is currently unavailable in the United States.

Dabigatran is a DTI indicated for prevention of stroke and systemic embolism in patients with non-valvular atrial fibrillation.³⁴⁴ Dabigatran levels can be measured using the Hemoclot® thrombin inhibition test (a dilute thrombin time) or the ecarin clotting time, but monitoring is generally not necessary to guide therapy. The aPTT is relatively insensitive to dabigatran (approximately 2-fold prolongation at peak levels post-dosing and 1.5-fold prolonged at trough levels) and the sensitivity of the aPTT to dabigatran can vary depending on the reagents and coagulometers employed.³⁴⁵ However, the aPTT can be used to gain a rough idea of whether dabigatran is present. The half-life of dabigatran in healthy subjects is 12 to 17 hours.³⁴⁴ According to the manufacturer, the concomitant use of dabigatran with P-glycoprotein (P-gp) inducers (eg, rifampin) reduces exposure to dabigatran and should generally be avoided. P-gp inhibition and impaired renal function are the major independent factors that result in increased exposure to dabigatran. Concomitant use of P-gp inhibitors in patients with renal impairment is expected to produce increased exposure of dabigatran compared to that seen with either factor alone.³⁴⁴ For patients with $C_{cr} > 30$ mL/min, the manufacturer recommends a dose of 150 mg orally, twice daily.³⁴⁴ No dose adjustment of dabigatran is recommended in

patients with mild or moderate renal impairment. For patients with C_{cr} 15 to 30 mL/min, the recommended dosing is 75 mg orally, twice daily.³⁴⁴ Recent reports of bleeding complications in patients taking dabigatran have noted that reduced renal function (C_{cr} <50 mL/min), lower body weight (<60 kg), and older age (\geq 80 years) were associated with an increased risk of bleeding complications.^{346,347} Since there is no antidote to dabigatran should bleeding occur, it should be avoided in patients with reduced renal function, lower body weight, and advanced age. Because dabigatran has not been tested in cancer patients, the NCCN Panel currently does not recommend its use for therapeutic or prophylactic anticoagulation or for the management of HIT in patients with cancer.

Warfarin

Warfarin inhibits production of functional forms of vitamin K-dependent anticoagulation factors, such as factors II, VII, IX, and X as well as the endogenous anticoagulant proteins, protein C and protein S, by the liver.²¹⁹ Warfarin dose requirements are highly variable and influenced by a large number of factors, including individual genetic factors (polymorphisms of the vitamin K epoxide reductase and CYP2C9 genes), vitamin K intake, use of medications that influence warfarin and vitamin K metabolism, and liver function. Therefore, close monitoring of the INR (ratio of PT to the mean normal PT normalized for PT reagent sensitivity to warfarin-induced reductions in vitamin K-dependent coagulation factors) is required to determine the therapeutic warfarin dose for an individual patient.³³⁸ The panel recommends a target INR of 2 to 3 for VTE treatment, which is consistent with ACCP recommendations.¹⁸³ Initially, the INR should be checked at least twice weekly during the transition phase from concurrent therapy with a parenteral anticoagulant (UFH, LMWH, or fondaparinux) to warfarin monotherapy. Once stable INRs are achieved, monitoring can be

gradually decreased in frequency in a step-wise fashion from once weekly to once monthly. Dose changes, addition of new medications—particularly medications with the potential to interact with warfarin—or changes in clinical status should prompt more frequent monitoring. A recent multicenter randomized clinical trial demonstrated that computer-assisted dosing of warfarin was superior to dosing directed by experienced providers.³⁴⁸ Therefore, use of computer-assisted dosing should be considered in the management of patients on chronic warfarin therapy. Care should be used when making the transition from a DTI to warfarin in the management of HIT, because all of the DTIs prolong the INR to a varying degree (the strength of this effect is: argatroban > bivalirudin > lepirudin)^{181,338,349,350}, and the duration of this effect is extended in argatroban-treated patients with hepatic dysfunction³⁴³ (see Guidelines section *Therapeutic Options for Heparin-Induced Thrombocytopenia [HIT]*).

Related Issues in VTE Prophylaxis and Treatment

Reversal of Anticoagulant Activity

The anticoagulant effects of UFH are fully reversible with protamine sulfate, and the anti-Xa activity of LMWHs are partially reversed by protamine sulfate (up to 60%–75% depending on the LMWH).^{157,158,160,351} This agent must be used with caution because it can cause severe hypotension or anaphylactoid reactions, particularly if infused more rapidly than 5 mg per minute.^{157,158,160,161,181,351-353} Patients with fish allergies, previous exposure to protamine (eg, NPH insulin), or vasectomized or infertile men are at increased risk for allergic reactions^{181,352} (see Guidelines section *Reversal of Anticoagulation*). The management of patients with a supratherapeutic INR while treated with warfarin is a common clinical challenge. In many cases, the effects of warfarin therapy in the patient with an elevated INR who is not bleeding can be reversed by withholding or reducing the warfarin dose,



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

and, depending on the INR, the addition of small doses of oral vitamin K1 for patients who are thought to be at higher risk of bleeding.^{219,354-356} It should be noted that a randomized placebo-controlled trial of 1.25 mg oral vitamin K versus placebo for INR reversal in asymptomatic patients (with an INR of 4.5–10) did not demonstrate any reduction in bleeding or thromboembolic complications.³⁵⁷ Of note, only 8% of the participants had active cancer. Therefore, use of oral vitamin K should be considered on a case-by-case basis. Consistent with the 2012 ACCP Guidelines, the NCCN Panel recommends the use of oral vitamin K (1–2.5 mg) for patients with an INR greater than 10 on warfarin, and who have no evidence of bleeding.³⁵⁶ Patients with an INR ranging up to 10 on warfarin, and who have no evidence of bleeding, should have their warfarin dose held, but the routine use of vitamin K is not warranted in these cases.³⁵⁶ It is prudent to review potential drug and/or dietary interactions so that the effects of such interactions can be eliminated or taken into account for future warfarin dosing. After withholding/reducing the warfarin dose and administering vitamin K1 as needed, the INR should be monitored closely—daily for inpatients and every 1 to 2 days for outpatients. When INR approaches therapeutic levels (INR <4), warfarin may be restarted at reduced doses if a causal factor cannot be identified or eliminated. INR should then be rechecked after 4 to 7 days, and warfarin dose should be adjusted based on weekly INR measurements until stable therapeutic levels are obtained.

For patients requiring rapid warfarin reversal for urgent or emergent surgical procedures, IV administration of vitamin K1 may be preferred over oral vitamin K1. In a prospective randomized study that compared INR outcomes with IV (0.5 mg) or oral (2.5 mg) vitamin K1 in patients with baseline INR values of 6 to 10 on warfarin (47 episodes among n=44), a greater proportion of patients in the IV therapy arm achieved rapid therapeutic INR (2–4) at 6 hours (46% vs. 0%) and at 12 hours

(67% vs. 35%) compared with oral therapy.³⁵⁸ In a recent prospective study that evaluated vitamin K1 in patients requiring rapid warfarin reversal for elective surgery (n=178), IV vitamin K1 (3 mg; given 12–18 hours prior to procedure) resulted in INR ≤1.5 on the day of surgery in nearly all patients (94%).³⁵⁹ Thus, for patients on warfarin requiring reversal within 24 hours of surgery, IV vitamin K1 (1–2.5 mg over 1 hour) is recommended. INR assessment should be repeated prior to surgery to determine the need for supplemental reversal agents such as fresh frozen plasma (FFP). For patients requiring reversal within 48 hours of surgery, 2.5 mg of oral vitamin K1 can be given. For these cases, INR measurements should be repeated 24 hours prior and immediately prior to surgery to determine the need for additional vitamin K or FFP (see Guidelines section *Reversal of Anticoagulation*).

The patient with serious or life-threatening bleeding requires 10 mg of IV vitamin K1 and the 4-factor prothrombin complex concentrate (4-factor PCC).^{354,360} Close monitoring of INR is required. If 4-factor PCC is unavailable or the patient is allergic to heparin or had experienced HIT within a year, 3-factor PCC may be used. Other alternatives include FFP and rhFVIIa. Administration of IV vitamin K1 alone is insufficient in such a critical situation because it requires at least 4 to 6 hours to begin to reduce the INR.³⁶¹ Since warfarin acts by inhibiting production of functional vitamin K-dependent clotting factors (factors II, VII, IX, and X) as well as protein S and protein C, the administration of a 4-factor PCC containing therapeutic quantities of factors II, VII, IX, and X can facilitate the rapid reversal of its anticoagulation effect.³⁶⁰ FFP alone can be given in place of PCC plus FFP if PCC is not available, but a disadvantage of this approach is the time delay associated with the preparation, delivery, and infusion of FFP.^{351,362} RhVIIa can also be used to rapidly reverse warfarin in place of either FFP or PCC.^{354,363,364} There is a small risk for anaphylaxis (3 per 10,000) associated with the IV



administration of vitamin K1, especially when it is administered more rapidly than 1 mg per minute,^{351,365} and PCC and rhFVIIa have been associated with a low risk for thromboembolic events.^{366,367}

Specific agents to reverse many of the newer anticoagulants do not exist. There is less evidence to guide the management of patients treated with these drugs in need of anticoagulant reversal. Nevertheless, IV rhFVIIa, which rapidly induces thrombin generation, has been shown to reduce the anticoagulant effects of LMWHs, DTIs, and fondaparinux in laboratory tests.^{181,351,364,368-373} Although evidence from published studies are limited, available data from *in vitro* models and healthy volunteers support the use of rhFVIIa for management of severe bleeding events with fondaparinux.^{181,368-370} For DTIs, activated PCC (APCC) such as FEIBA has been evaluated as a potential option for reversing the effects of DTIs by improving hemostatic capacity.³⁷⁴⁻³⁷⁶ It is important to note that rhFVIIa and APCC have been associated with thromboembolic events,^{372,377} and therefore require caution when employing these agents as potential reversal agents. Other proposed strategies for reversal of DTIs include administration of FFP or cryoprecipitate; use of desmopressin acetate (DDAVP), which stimulates release of factor VIII and von Willebrand factor; and antifibrinolytic agents, which block plasmin activity (ie, the enzyme which breaks down fibrin clots).^{351,378-383} It is important to remember that DDAVP is effective only for 3 or 4 doses after which tachyphylaxis develops (see Guidelines section on *Reversal of Anticoagulation*).^{384,385} Although rare, DDAVP has also been associated with hyponatremia.³⁸⁴ The DTI dabigatran is primarily excreted in the urine and shows low plasma protein binding (approximately 35% bound to human plasma proteins). Thus, dabigatran can be dialyzed with the removal of about 60% of drug over 2 to 3 hours, although clinical data supporting this approach are limited.^{344,345,386,387} APCC and rhFVIIa have been

suggested as reversal agents for dabigatran, as with other DTIs.^{387,388} Activated charcoal may also be considered for reversal of dabigatran, especially within a few hours of overdose.^{345,389}

As with the newer anticoagulant agents mentioned above, no specific reversal agent currently exists for the direct factor Xa inhibitors rivaroxaban and apixaban.³⁹⁰ The prescribing information indicates that activated charcoal may be considered for reduction of drug absorption.^{204,208} Due to the high plasma protein binding, these agents are not expected to be dialyzable. Limited data from *in vivo* models and healthy volunteers suggest that PCC may at least partially reverse the anticoagulation effects.^{391,392} The use of rhFVIIa may also be considered, although data are presently unclear in terms of its benefits.^{388,389,393}

Failure of Anticoagulation Therapy

Anticoagulation failure is defined as extension of DVT or PE, or new DVT or PE, while on recommended anticoagulation therapy (see Guidelines section on *Therapeutic Anticoagulation Failure*).³⁹⁴ Although there are many potential causes of anticoagulation therapy failure, an initial determination of whether the INR or aPTT is within the therapeutic range is important for patients with recurrent VTE who are receiving warfarin or UFH, respectively. When INR or aPTT values are subtherapeutic, one obvious option is to increase the anticoagulant dose to a therapeutic level.

Although anticoagulation therapy failure for patients receiving warfarin, UFH, LMWH, or fondaparinux can result if the prescribed anticoagulant dose is inadequate, other factors to consider include patient adherence to self-administered medications, such as an oral vitamin K antagonists, or subcutaneously administered anticoagulants, and the dosing frequency for patients receiving LMWH.³⁹⁴ For example, an increased

risk for VTE recurrence was reported in one study of cancer patients receiving once-daily enoxaparin in the acute therapy setting.³⁹⁵ Thus, a twice-daily dosing schedule is an option for patients exhibiting recurrent VTE while receiving once-daily therapy with an LMWH. A dose increase can also be considered for patients exhibiting recurrent VTE while receiving anticoagulant therapies for which anticoagulant effects are not typically monitored in the laboratory (eg, LMWH and fondaparinux).³⁹⁶

INR or aPTT values may be subtherapeutic in situations where inadequate anticoagulant dosing is not the direct cause of recurrent VTE. For example, warfarin resistance (inability to achieve a therapeutic INR on warfarin doses typically used to treat VTE) can be due to genetic variability associated with the enzymatic metabolism of warfarin, or the concomitant administration of medications that interact with warfarin.^{397,398} An option for patients undergoing warfarin therapy and exhibiting a subtherapeutic INR is a switch to intravenous LMWH (preferred), intravenous UFH, or fondaparinux. A switch to LMWH in the setting of a subtherapeutic INR with warfarin therapy is supported by the results of one study in which a low VTE recurrence rate was reported for patients treated with LMWH following failure of warfarin therapy.³⁹⁹ Likewise, heparin resistance (inability to achieve therapeutic aPTT on heparin doses typically used to treat VTE), though rare, can occur as a result of pharmacokinetic or biophysical/physiologic limitations of heparin therapy.⁴⁰⁰

Anticoagulation failure of warfarin or UFH can also occur in the setting of a therapeutic INR or aPTT value. Causes include cancer-related hypercoagulability such as the Trousseau's syndrome, HIT, cancer-related anatomic causes such as vascular compression, and acquired and/or familial thrombophilia.^{394,400} Diagnostic testing to identify syndromes identified above, when present, is critical to the management of VTE in these patients.³⁹⁴ In patients with anatomic

compression due to congenital (eg, May-Thurner syndrome or iliac vein compression syndrome, thoracic outlet syndrome) or acquired causes (vascular compression by nodal or tumor masses), relief of anatomic compression is essential to preventing recurrent thrombosis. Clinical suspicion of HIT should be high when recurrent VTE is observed in a cancer patient receiving heparin-based therapy or in a patient who received such therapy in the recent past. Options for patients with VTE recurrence while receiving UFH characterized by a therapeutic aPTT level include a switch to intravenous LMWH or fondaparinux or an increase in the dose of intravenous UFH. Likewise, patients with recurrent VTE and a therapeutic INR while on warfarin therapy can be switched to intravenous heparin (LMWH preferred) or fondaparinux. A switch to heparin-based therapy is an option following failure of fondaparinux to prevent VTE recurrence and vice versa.

Placement of an IVC filter is an option for treating patients with PE or progression of central DVT despite therapeutic anticoagulation with UFH, LMWH, or fondaparinux, although filters should be avoided in the setting of HIT or migratory thrombophlebitis due to the systemic nature of these coagulopathies.^{98,99}

Perioperative Management of Anticoagulation and Antithrombotic Therapy

Management of surgery-associated bleeding in cancer patients is complicated by the need of anticoagulation in many of these patients due to their malignancy, cancer therapy, and/or comorbidities. A balance of the thrombotic risk and bleeding risk is important. An IVC filter should be considered if VTE occurred within 1 month of planned surgery. Bridging anticoagulation therapy refers to the use of short-acting anticoagulants (LMWH or UFH) for 10 to 20 days during the peri-procedural period.⁴⁰¹



Cancer patients on anticoagulants requiring emergent surgery should be managed according to the Guidelines section *Reversal of Anticoagulation in the Event of Life-threatening Bleeding or Emergent Surgery*.

If the surgical procedure is non-emergent, an assessment of the bleeding risk should be performed before the procedure. To date, a consensus has not been reached on the optimal preoperative screening strategy.⁴⁰¹ Guidelines on estimating bleeding risk based on the type of surgical procedure are listed in Table 1 of the Guidelines section *Perioperative Management of Anticoagulation and Antithrombotic Therapy*. Patients assessed to be at very low risk of bleeding can continue anticoagulation and undergo surgery. All other patients should then be assessed for thromboembolic risk according to Table 3 of the same Guidelines section. In general, anticoagulation therapy should be stopped before surgery for these patients. Bridging anticoagulation therapy should be administered for patients at high thrombotic risk and considered for patients at moderate thrombotic risk. Panel recommendations on anticoagulation before and after surgery based on both the bleeding and thrombotic risks are detailed in Table 2.

Diagnosis and Management of HIT

Specific guideline recommendations regarding HIT are available from the ACCP.^{202,402} HIT is caused by a relatively common immunologic reaction to heparin-based products. In one pharmacy-based surveillance study, 0.2% of patients receiving heparin therapy developed HIT, although the incidence of HIT was 1.2% in patients exposed to heparin for more than 4 days.⁴⁰³ In another study, 2.7% of patients treated with UFH developed HIT.⁴⁰⁴ HIT is triggered when UFH (or to a lesser extent LMWH) bind to platelet factor 4 (PF4) released by activated platelets and form an immunogenic PF4-heparin complex

leading to the development of antibodies. These antibodies increase platelet clearance and activate platelets, resulting in the release of procoagulant microparticles and increased thrombin generation.^{202,405} The end result is a consumptive thrombocytopenia and profound prothrombotic state that triggers symptomatic thromboembolism in as many as 75% of patients.^{202,405} Clinical evidence of HIT includes development of thrombocytopenia, formation of necrotic lesions at injection sites, arterial thromboembolic complications, and/or development of VTE.^{406,407} Most typically, HIT occurs after 5 to 10 days following initial exposure to heparin-based products. In rapid-onset HIT, HIT can occur within 1 day following administration of heparin in a patient with previous exposure to such agents within a period of 100 days.²⁰² Delayed-onset HIT is less common, and can occur days or weeks after heparin therapy has been discontinued.⁴⁰⁵

Some evidence indicates that cancer patients are at increased risk of developing HIT and HIT-related VTE,^{408,409} although this has not been firmly established. HIT has been associated with the use of both LMWHs and UFH. Increased rates of HIT have been observed in patients receiving heparin-based therapy who were previously exposed to such therapy.⁴¹⁰ Results of some studies have indicated that the frequency of HIT with LMWH and UFH is similar,^{410,411} whereas other studies suggest a lower incidence of HIT in patients receiving LMWH relative to those receiving UFH.^{404,412-414} It has been suggested that factors such as anticoagulant dose (lower with prophylactic doses, higher with treatment doses) and whether the patient is treated in the medical (lower-risk) or surgical (higher-risk) setting may account for these conflicting results, since a lower relative incidence of HIT with LMWH was primarily observed for surgical patients receiving prophylactic doses of anticoagulant therapy.⁴¹⁵

A diagnosis of HIT is based on both clinical and serologic evidence.²⁰² Hence, the presence of both clinical sequelae of HIT, including thrombosis and thrombocytopenia defined as a drop in platelet count by more than 50%, and anti-PF4/heparin antibodies are needed for a diagnosis. Furthermore, since most HIT antibodies do not activate platelets, a negative test result is more useful for excluding the diagnosis than a positive test result is for confirming it. As mentioned by Greinacher et al, “all HIT is caused by platelet activating antibodies, but not all PF4/heparin antibodies cause HIT.”⁴¹⁶ The specificity of functional platelet activation assays, such as the serotonin release assay (SRA), is higher than antigen assays, such as the heparin-PF4 ELISA, which detect the presence of HIT antibodies, but do not assess their ability to activate platelets.²⁰²

The diagnosis of HIT is complicated by the high frequency of heparin use in hospitals; the presence of HIT antibodies, which do not activate platelets; possible delays in obtaining serologic test results; and multiple causes of thrombocytopenia in patients receiving heparin-based products. In addition, there are increased bleeding risks associated with substitution of a DTI for heparin. Therefore, it is critically important that a high level of clinical suspicion is present before a patient is treated for HIT.⁴¹⁶

The 4Ts score is a simple, validated tool designed to assess the probability of HIT based on specific characteristics of 4 clinical parameters: **t**hrombocytopenia; the **t**iming of the onset of platelet fall; the presence of **t**hrombosis or other clinical sequelae; and evidence of **o**ther potential causes of thrombocytopenia (see Guidelines section *HIT Pre-test Probability Score Assessment*).⁴¹⁷⁻⁴¹⁹ Each of these 4 parameters is weighted by a score of 0 to 2 according to how likely it reflects a HIT diagnosis. A total score of 0 to 8 is possible. Total scores are grouped into 3 categories, which classify the patient as being at low-

(0–3), medium- (4–5), or high-risk (6–8) of HIT.⁴¹⁹ As described above for HIT antibody testing, evidence suggests that the negative predictive value of this assessment tool is considerably higher than its positive predictive value; hence, this tool is more likely to be useful in identifying patients at low risk for HIT.^{418,420} Cuker and colleagues⁴²¹ developed an alternative pre-test probability model based on broad expert opinion of HIT diagnosis known as the HIT Expert Probability score (HEP score). In a validation patient cohort, the HEP score demonstrated greater inter-observer reliability and correlation with laboratory test results and expert assessment of the probability of HIT diagnosis than the 4T score.⁴²¹ Neither HIT pre-test probability model has been assessed in an oncology-specific population.

The panel recommends platelet monitoring at baseline and then every 2 to 3 days for at least the first 14 days, and then every 2 weeks thereafter, or more frequently as clinically indicated, in patients receiving anticoagulation therapy with UFH or LMWH. If HIT is suspected, the patient should be evaluated using the 4Ts score. Recommendations for patients classified as being at low risk for HIT include the following: consider alternative causes of thrombocytopenia; weigh the risks/benefits of continued therapy with heparin versus a DTI or fondaparinux; consider maintaining anticoagulation with heparin; monitor their clinical status; and consider HIT antibody testing by ELISA in select patients based on clinical judgement. Patients classified as being at moderate/high risk for HIT on the basis of the 4Ts score should initially be managed as having a diagnosis of HIT. HIT antibody testing by ELISA should be ordered, although immediate discontinuation of heparin-based products and administration of an alternative anticoagulant, typically a DTI, is recommended. Warfarin should be discontinued and reversed with vitamin K. The safety of platelet transfusions in patients with HIT remains controversial. Platelet

transfusions may be considered for clinically significant bleeding or prior to invasive procedures in patients with a platelet count less than 50,000/mcL. Prophylactic platelet transfusions are otherwise not recommended because of the theoretic risk of triggering further thrombosis.

The results of HIT antibody testing by ELISA further direct management. For example, options for patients with a negative HIT antibody test result include a reassessment of anticoagulation therapy based on the 4Ts score, and consideration of repeat HIT antibody testing if the pre-test probability of HIT is moderate to high. A diagnosis of HIT can be ruled out in those patients with a negative HIT antibody test on repeat testing. The management of patients with a positive HIT antibody test on initial testing should be re-evaluated based on the 4Ts score pre-test probability. Patients with a moderate/high 4Ts score should be managed according to recommendations for patients with a diagnosis of HIT, whereas SRA testing should be considered in those with a low pre-test probability with test results directing further management. A 4-extremity duplex ultrasound may be considered in confirmed HIT case to identify sub-clinical DVT.

Anticoagulants for the Treatment of HIT

Direct Thrombin Inhibitors

DTIs available for the management of HIT include argatroban and bivalirudin.²⁰² Two prospective clinical trials evaluated the activity of argatroban in patients with clinically diagnosed HIT, with or without concurrent thrombosis.^{422,423} In the initial trial, argatroban significantly reduced the combined endpoint of death, limb amputation, and occurrence of new thrombotic events among patients with HIT without thrombosis (n=160), compared with historical controls (25.6% vs. 38.8%; $P = .014$). No significant differences in the combined endpoint were noted with argatroban versus control among patients with HIT and

thrombosis (n=144).⁴²² Similarly, results from the second trial of argatroban showed significantly decreased incidence of the combined endpoint with argatroban compared with historical controls in patients with HIT without thrombosis (n=189) (28.0% vs. 38.8%; $P = .04$), but not in patients with HIT and thrombosis (n=229) (41.5% vs. 56.5%; $P = .07$).⁴²³ In both trials, argatroban was shown to significantly decrease the incidence of death due to thrombosis, as well as the incidence of new thrombosis compared with controls ($P < .05$), in both groups of patients with HIT with or without concurrent thrombosis.^{422,423}

Argatroban is approved by the FDA for the immediate treatment of HIT.³⁴³ Argatroban is primarily metabolized by the liver, and prolonged clearance of this agent has been seen in patients with hepatic insufficiency.³⁴³ Therapeutic dosing regimens of many anticoagulants used in the treatment of critically ill patients with organ dysfunction and HIT are often lower than those recommended by the manufacturer and require frequent monitoring. The manufacturer-recommended dose for argatroban may be too high, especially for the treatment of HIT in critically ill patients.⁴²⁴⁻⁴²⁶ Argatroban administered at a reduced dose of 1 mcg/kg/min may be adequate to provide sufficient anticoagulation.⁴²⁶ Dose reductions have also been suggested for bivalirudin,⁴²⁷ another DTI, when used off-label in the treatment of HIT⁴²⁸ and in patients with HIT and hepatic and/or renal insufficiency or critically ill patients.^{350,429,430} Although some of the pharmacologic characteristics of bivalirudin, including HIT short half-life and enzymatic metabolism, are advantageous in the setting of HIT, data regarding its use in HIT are limited.³⁵⁰

Dabigatran is a DTI approved by the FDA for the reduction of risk for stroke and systemic embolism in patients with non-valvular atrial fibrillation.³⁴⁴ This agent is not approved for the treatment of HIT, and



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

the NCCN Guidelines Panel for VTE does not currently recommend the use of dabigatran in patients with cancer.

The panel recommends a DTI as the preferred treatment for the immediate management of HIT. No head-to-head trials comparing different DTIs in the treatment of HIT have been published. Clinician experience and comfort level with the agents used for the immediate treatment of HIT should be a consideration in the choice of therapy. Use of argatroban should be avoided in patients with hepatic failure and severe renal insufficiency, respectively.

Fondaparinux

The option of off-label use of fondaparinux as an alternative to parenteral DTIs in the treatment of a current episode of HIT without thrombosis is also included in the NCCN Guidelines for VTE.⁴³¹ Advantages to the use of fondaparinux in this setting, in addition to subcutaneous administration, include its lack of INR prolongation when administered concomitantly with warfarin. Although the long half-life of fondaparinux is a disadvantage in situations where anticoagulation reversal is necessary, a possible benefit may include a decreased risk for rebound hypercoagulability.⁴³² Furthermore, unlike DTIs, aPTT testing is not used to monitor treatment response of fondaparinux, thereby eliminating problems associated with warfarin prolongation of the aPTT when overlapped with a DTI. Fondaparinux has been used in small numbers of patients with HIT and generally appears to be safe.⁴³³⁻⁴³⁵ There have been rare reports of an association between fondaparinux use and development of HIT, although in most cases patients had prior exposure to UFH or LMWH.⁴³⁶⁻⁴³⁹ It has been suggested that use of fondaparinux for patients with HIT and without a contraindication to fondaparinux be restricted to those who have recovered from a recent episode of HIT without thrombosis and are ready to be discharged from the hospital, but who are not yet stable on

warfarin therapy.^{202,432} Fondaparinux is included in the guidelines as a category 2B option for the immediate management of HIT.

Warfarin

The panel recommends against the use of warfarin therapy in patients with a moderate or high pre-test probability of HIT by the 4T score. For patients receiving warfarin, it should be discontinued and reversed with vitamin K.²⁰² Warfarin should not be initiated in patients with HIT until after platelet count recovery because of the potential for skin necrosis and/or venous gangrene, which can result from warfarin-induced reductions in protein C levels in the setting of profound activated coagulation due to HIT.^{202,440} After platelet recovery ($\geq 150,000/\text{mL}$ or when platelets return to baseline), warfarin should be overlapped with a DTI or fondaparinux for at least 5 days; the DTI or fondaparinux should be discontinued only after the INR has reached the intended target range (INR 2–3) for 24 hours. Since both DTIs and warfarin reduce thrombin activity, co-administration of a DTI and warfarin produces a combined effect on the laboratory measurements of both aPTT and INR. However, concurrent therapy, compared with warfarin monotherapy, exerts no additional effect on vitamin K-dependent factor X activity. Therefore, the anticoagulation impact of warfarin may be underestimated in the presence of a DTI. Argatroban in particular, but also other DTIs, can prolong the INR during co-therapy with warfarin. Since argatroban has the lowest affinity for thrombin of the 3 DTIs, higher molar plasma concentrations of argatroban are needed to prolong the aPTT; hence, prolongation of INR is more pronounced with argatroban compared with the other DTIs.^{349,441} A higher target INR should therefore be achieved before argatroban is discontinued.^{202,343,441} Once DTI is discontinued, a repeat INR and aPTT should be obtained 2-6 hours later to determine whether the INR is therapeutic on warfarin monotherapy. Alternatively,



chromogenic factor X levels (which are not affected by DTIs) can be used to monitor warfarin activity during transition from co-therapy with argatroban.^{442,443} The duration of warfarin therapy is dependent on whether HIT is accompanied by thrombosis. In patients with HIT and thrombosis, the duration of therapy is dictated by the nature of the thrombotic event (3 months for DVT, 6 months for PE). In patients with HIT without thrombosis, at least 1 month of warfarin therapy is recommended.⁴⁰⁵

Withholding Anticoagulation Therapy: Elements to Consider in the Decision Not to Treat

The feasibility of invasive or aggressive intervention is not the only consideration for VTE prophylaxis and treatment in cancer patients.⁴⁴⁴ The risks and probability of success of the interventions should be considered as well. Factors to consider before implementing anticoagulation therapy include patient refusal; lack of therapeutic advantage; lack of palliative benefits; and whether anticoagulation is associated with an unreasonable burden. Likewise, careful consideration of these issues is also very important in deciding to withhold or withdraw VTE therapy.

Summary

Recognizing the increased risk for VTE in cancer patients is the first step in preventing the occurrence of VTE and promptly identifying VTE in these patients. The NCCN Guidelines Panel recommends VTE thromboprophylaxis for all hospitalized patients with cancer who do not have contraindications to such therapy, and the panel also emphasizes that an increased level of clinical suspicion of VTE should be maintained for cancer patients. Following hospital discharge, it is recommended that cancer patients in a high-risk setting for VTE continue to receive VTE prophylaxis, with the duration of

anticoagulation determined by the clinical situation. Careful evaluation of cancer patients in whom VTE is suspected and prompt treatment and follow-up for patients diagnosed with VTE is recommended after the cancer status of the patient is assessed and the risks and benefits of treatment are considered.

Discussion
update in
progress



Patient Resources for VTE Management

Websites:

- National Blood Clot Alliance - www.stoptheclot.org
- ClotCare Online Resource - <http://www.clotcare.com/>
- North American Thrombosis Forum - <http://natfonline.org/patients>
- Clot Connect - <http://www.clotconnect.org/patients>
- Living with Thrombophilia - <http://fvleiden.org/index.html>

Related Materials:

- Preventing deep vein thrombosis (American College of Obstetricians and Gynecologists) - <http://www.acog.org/~media/For%20Patients/faq174.pdf?dmc=1&ts=20140618T1232288986>
- Deep vein thrombosis (American Academy of Family Physicians) - <http://familydoctor.org/familydoctor/en/diseases-conditions/deep-vein-thrombosis.html>
- Deep vein thrombosis (National Heart, Lung, and Blood Institute) - <http://www.nhlbi.nih.gov/health/health-topics/topics/dvt/>
- Deep vein thrombosis (MedLinePlus) - <http://www.nlm.nih.gov/medlineplus/deepveinthrombosis.html>
- Deep vein thrombosis/pulmonary embolism (Centers for Disease Control and Prevention) - <http://www.cdc.gov/ncbddd/dvt/facts.html>
- Your guide to preventing and treating blood clots (Agency for Healthcare Research and Quality) - <http://www.ahrq.gov/patients-consumers/prevention/disease/bloodclots.html>

References

1. Khorana AA, Francis CW, Culakova E, et al. Thromboembolism in hospitalized neutropenic cancer patients. *J Clin Oncol* 2006;24:484-490. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16421425>.
2. Sallah S, Wan JY, Nguyen NP. Venous thrombosis in patients with solid tumors: determination of frequency and characteristics. *Thromb Haemost* 2002;87:575-579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12008937>.
3. Khorana AA, Dalal M, Lin J, Connolly GC. Incidence and predictors of venous thromboembolism (VTE) among ambulatory high-risk cancer patients undergoing chemotherapy in the United States. *Cancer* 2013;119:648-655. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22893596>.
4. Heit JA, O'Fallon WM, Petterson TM, et al. Relative impact of risk factors for deep vein thrombosis and pulmonary embolism: a population-based study. *Arch Intern Med* 2002;162:1245-1248. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12038942>.
5. Monreal M, Fernandez-Llamazares J, Perandreu J, et al. Occult cancer in patients with venous thromboembolism: which patients, which cancers. *Thromb Haemost* 1997;78:1316-1318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9408011>.
6. Bick RL. Cancer-associated thrombosis: focus on extended therapy with dalteparin. *J Support Oncol* 2006;4:115-120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16553136>.
7. Lopez JA, Kearon C, Lee AY. Deep venous thrombosis. *Hematology Am Soc Hematol Educ Program* 2004:439-456. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15561697>.
8. Prandoni P, Piccioli A, Girolami A. Cancer and venous thromboembolism: an overview. *Haematologica* 1999;84:437-445. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10329923>.
9. Otten HM, Mathijssen J, ten Cate H, et al. Symptomatic venous thromboembolism in cancer patients treated with chemotherapy: an underestimated phenomenon. *Arch Intern Med* 2004;164:190-194. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14744843>.
10. Chew HK, Wun T, Harvey D, et al. Incidence of venous thromboembolism and its effect on survival among patients with common cancers. *Arch Intern Med* 2006;166:458-464. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16505267>.
11. Chew HK, Wun T, Harvey DJ, et al. Incidence of venous thromboembolism and the impact on survival in breast cancer patients. *J Clin Oncol* 2007;25:70-76. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17194906>.
12. Kuderer NM, Francis CW, Culakova E, et al. Venous thromboembolism and all-cause mortality in cancer patients receiving chemotherapy (abstract). *J Clin Oncol* 2008;26(Supple 15):Abstract 9521. Available at: http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/9521.
13. Martino MA, Williamson E, Siegfried S, et al. Diagnosing pulmonary embolism: experience with spiral CT pulmonary angiography in gynecologic oncology. *Gynecol Oncol* 2005;98:289-293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15950268>.
14. Sorensen HT, Mellekjaer L, Olsen JH, Baron JA. Prognosis of cancers associated with venous thromboembolism. *N Engl J Med* 2000;343:1846-1850. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11117976>.
15. Agnelli G, Bolis G, Capussotti L, et al. A clinical outcome-based prospective study on venous thromboembolism after cancer surgery: the @RISTOS project. *Ann Surg* 2006;243:89-95. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16371741>.
16. Kakkar AK, Levine M, Pinedo HM, et al. Venous thrombosis in cancer patients: insights from the FRONTLINE survey. *Oncologist*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

2003;8:381-388. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12897335>.

17. Cohen AT, Tapson VF, Bergmann JF, et al. Venous thromboembolism risk and prophylaxis in the acute hospital care setting (ENDORSE study): a multinational cross-sectional study. *Lancet* 2008;371:387-394. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18242412>.

18. Tapson VF, Decousus H, Pini M, et al. Venous thromboembolism prophylaxis in acutely ill hospitalized medical patients: findings from the International Medical Prevention Registry on Venous Thromboembolism. *Chest* 2007;132:936-945. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17573514>.

19. Alikhan R, Peters F, Wilmott R, Cohen AT. Fatal pulmonary embolism in hospitalised patients: a necropsy review. *J Clin Pathol* 2004;57:1254-1257. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15563663>.

20. Cohen AT, Alikhan R, Arcelus JI, et al. Assessment of venous thromboembolism risk and the benefits of thromboprophylaxis in medical patients. *Thromb Haemost* 2005;94:750-759. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16270626>.

21. Khorana AA, Connolly GC. Assessing risk of venous thromboembolism in the patient with cancer. *J Clin Oncol* 2009;27:4839-4847. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19720906>.

22. Connolly GC, Khorana AA. Emerging risk stratification approaches to cancer-associated thrombosis: risk factors, biomarkers and a risk score. *Thromb Res* 2010;125 Suppl 2:1-7. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20433985>.

23. Venous thromboembolism in adult hospitalizations - United States, 2007-2009. *MMWR Morb Mortal Wkly Rep* 2012;61:401-404. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22672974>.

24. Ageno W, Squizzato A, Garcia D, Imberti D. Epidemiology and risk factors of venous thromboembolism. *Semin Thromb Hemost* 2006;32:651-658. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17024592>.

25. Hicks LK, Cheung MC, Ding K, et al. Venous thromboembolism and nonsmall cell lung cancer: a pooled analysis of National Cancer Institute of Canada Clinical Trials Group trials. *Cancer* 2009;115:5516-5525. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19711465>.

26. Khorana AA, Kuderer NM, Culakova E, et al. Development and validation of a predictive model for chemotherapy-associated thrombosis. *Blood* 2008;111:4902-4907. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18216292>.

27. Khorana AA, Francis CW, Culakova E, Lyman GH. Risk factors for chemotherapy-associated venous thromboembolism in a prospective observational study. *Cancer* 2005;104:2822-2829. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16284987>.

28. Mandalà M, Barni S, Prins M, et al. Acquired and inherited risk factors for developing venous thromboembolism in cancer patients receiving adjuvant chemotherapy: a prospective trial. *Ann Oncol* 2010;21:871-876. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19713246>.

29. Darze ES, Latado AL, Guimaraes AG, et al. Incidence and clinical predictors of pulmonary embolism in severe heart failure patients admitted to a coronary care unit. *Chest* 2005;128:2576-2580. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16236926>.

30. Kroger K, Weiland D, Ose C, et al. Risk factors for venous thromboembolic events in cancer patients. *Ann Oncol* 2006;17:297-303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16282243>.

31. Prandoni P, Lensing AWA, Prins MH, et al. Residual venous thrombosis as a predictive factor of recurrent venous thromboembolism.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Ann Intern Med 2002;137:955-960. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12484710>.

32. Prandoni P, Lensing AWA, Piccioli A, et al. Recurrent venous thromboembolism and bleeding complications during anticoagulant treatment in patients with cancer and venous thrombosis. Blood 2002;100:3484-3488. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12393647>.

33. Rogers MA, Levine DA, Blumberg N, et al. Triggers of hospitalization for venous thromboembolism. Circulation 2012;125:2092-2099. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22474264>.

34. Schmidt M, Horvath-Puho E, Thomsen RW, et al. Acute infections and venous thromboembolism. J Intern Med 2012;271:608-618. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22026462>.

35. Blom JW, Doggen CJM, Osanto S, Rosendaal FR. Malignancies, prothrombotic mutations, and the risk of venous thrombosis. JAMA 2005;293:715-722. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15701913>.

36. Heit JA, Silverstein MD, Mohr DN, et al. Risk factors for deep vein thrombosis and pulmonary embolism: a population-based case-control study. Arch Intern Med 2000;160:809-815. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10737280>.

37. Ogren M, Bergqvist D, Wahlander K, et al. Trousseau's syndrome - what is the evidence? A population-based autopsy study. Thromb Haemost 2006;95:541-545. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16525584>.

38. Hall IE, Andersen MS, Krumholz HM, Gross CP. Predictors of venous thromboembolism in patients with advanced common solid cancers. J Cancer Epidemiol 2009;2009:182521-182521. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20445797>.

39. Sandhu R, Pan C-X, Wun T, et al. The incidence of venous thromboembolism and its effect on survival among patients with primary bladder cancer. Cancer 2010;116:2596-2603. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20336780>.

40. Levitan N, Dowlati A, Remick SC, et al. Rates of initial and recurrent thromboembolic disease among patients with malignancy versus those without malignancy. Risk analysis using Medicare claims data. Medicine (Baltimore) 1999;78:285-291. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10499070>.

41. Gerber DE, Grossman SA, Streiff MB. Management of venous thromboembolism in patients with primary and metastatic brain tumors. J Clin Oncol 2006;24:1310-1318. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16525187>.

42. Lee AY, Levine MN. Venous thromboembolism and cancer: risks and outcomes. Circulation 2003;107:117-21. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12814981>.

43. Marras LC, Geerts WH, Perry JR. The risk of venous thromboembolism is increased throughout the course of malignant glioma: an evidence-based review. Cancer 2000;89:640-646. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10931464>.

44. Connolly GC, Dalal M, Lin J, Khorana AA. Incidence and predictors of venous thromboembolism (VTE) among ambulatory patients with lung cancer. Lung Cancer 2012;78:253-258. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23026639>.

45. Falanga A, Marchetti M. Venous thromboembolism in the hematologic malignancies. J Clin Oncol 2009;27:4848-4857. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19752334>.

46. Palumbo A, Rajkumar SV, Dimopoulos MA, et al. Prevention of thalidomide- and lenalidomide-associated thrombosis in myeloma. Leukemia 2008;22:414-423. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18094721>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

47. Ottinger H, Belka C, Kozole G, et al. Deep venous thrombosis and pulmonary artery embolism in high-grade non Hodgkin's lymphoma: incidence, causes and prognostic relevance. *Eur J Haematol* 1995;54:186-194. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7720839>.
48. Andtbacka RHI, Babiera G, Singletary SE, et al. Incidence and prevention of venous thromboembolism in patients undergoing breast cancer surgery and treated according to clinical pathways. *Ann Surg* 2006;243:96-9101. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16371742>.
49. Haddad TC, Greeno EW. Chemotherapy-induced thrombosis. *Thromb Res* 2006;118:555-568. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16388837>.
50. Ay C, Dunkler D, Marosi C, et al. Prediction of venous thromboembolism in cancer patients. *Blood* 2010;116:5377-5382. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20829374>.
51. Cummings SR, Eckert S, Krueger KA, et al. The effect of raloxifene on risk of breast cancer in postmenopausal women: results from the MORE randomized trial. Multiple Outcomes of Raloxifene Evaluation. *JAMA* 1999;281:2189-2197. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10376571>.
52. Decensi A, Maisonneuve P, Rotmensz N, et al. Effect of tamoxifen on venous thromboembolic events in a breast cancer prevention trial. *Circulation* 2005;111:650-656. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15699284>.
53. Fisher B, Costantino JP, Wickerham DL, et al. Tamoxifen for the prevention of breast cancer: current status of the National Surgical Adjuvant Breast and Bowel Project P-1 study. *J Natl Cancer Inst* 2005;97:1652-1662. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16288118>.
54. Fisher B, Costantino JP, Wickerham DL, et al. Tamoxifen for prevention of breast cancer: report of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer Inst* 1998;90:1371-1388. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9747868>.
55. Vogel VG, Costantino JP, Wickerham DL, et al. Update of the National Surgical Adjuvant Breast and Bowel Project Study of Tamoxifen and Raloxifene (STAR) P-2 Trial: Preventing breast cancer. *Cancer Prev Res (Phila)* 2010;3:696-706. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20404000>.
56. Anderson GL, Limacher M, Assaf AR, et al. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA* 2004;291:1701-1712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15082697>.
57. Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321-333. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12117397>.
58. Blanco-Molina A, Trujillo-Santos J, Tirado R, et al. Venous thromboembolism in women using hormonal contraceptives. Findings from the RIETE Registry. *Thromb Haemost* 2009;101:478-482. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19277408>.
59. Gomes MPV, Deitcher SR. Risk of venous thromboembolic disease associated with hormonal contraceptives and hormone replacement therapy: a clinical review. *Arch Intern Med* 2004;164:1965-1976. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15477430>.
60. Lidegaard O, Nielsen LH, Skovlund CW, et al. Risk of venous thromboembolism from use of oral contraceptives containing different progestogens and oestrogen doses: Danish cohort study, 2001-9. *BMJ* 2011;343:d6423. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22027398>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

61. Manzoli L, De Vito C, Marzuillo C, et al. Oral contraceptives and venous thromboembolism: a systematic review and meta-analysis. *Drug Saf* 2012;35:191-205. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22283630>.
62. Jick SS, Hernandez RK. Risk of non-fatal venous thromboembolism in women using oral contraceptives containing drospirenone compared with women using oral contraceptives containing levonorgestrel: case-control study using United States claims data. *BMJ* 2011;342:d2151. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21511805>.
63. Parkin L, Sharples K, Hernandez RK, Jick SS. Risk of venous thromboembolism in users of oral contraceptives containing drospirenone or levonorgestrel: nested case-control study based on UK General Practice Research Database. *BMJ* 2011;342:d2139. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21511804>.
64. Leaf AN, Propert K, Corcoran C, et al. Phase III study of combined chemohormonal therapy in metastatic prostate cancer (ECOG 3882): an Eastern Cooperative Oncology Group study. *Med Oncol* 2003;20:137-146. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12835516>.
65. Bennett CL, Angelotta C, Yarnold PR, et al. Thalidomide- and lenalidomide-associated thromboembolism among patients with cancer. *JAMA* 2006;296:2558-2560. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17148721>.
66. El Accaoui RN, Shamseddeen WA, Taher AT. Thalidomide and thrombosis. A meta-analysis. *Thromb Haemost* 2007;97:1031-1036. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17549307>.
67. Hussein MA. Thromboembolism risk reduction in multiple myeloma patients treated with immunomodulatory drug combinations. *Thromb Haemost* 2006;95:924-930. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16732369>.
68. Knight R, DeLap RJ, Zeldis JB. Lenalidomide and venous thrombosis in multiple myeloma. *N Engl J Med* 2006;354:2079-2080. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16687729>.
69. Weber DM, Chen C, Niesvizky R, et al. Lenalidomide plus dexamethasone for relapsed multiple myeloma in North America. *N Engl J Med* 2007;357:2133-2142. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18032763>.
70. Zangari M, Siegel E, Barlogie B, et al. Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy. *Blood* 2002;100:1168-1171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12149193>.
71. Bennett CL, Silver SM, Djulbegovic B, et al. Venous thromboembolism and mortality associated with recombinant erythropoietin and darbepoetin administration for the treatment of cancer-associated anemia. *JAMA* 2008;299:914-924. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18314434>.
72. Baarslag HJ, Koopman MMW, Hutten BA, et al. Long-term follow-up of patients with suspected deep vein thrombosis of the upper extremity: survival, risk factors and post-thrombotic syndrome. *Eur J Intern Med* 2004;15:503-507. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15668085>.
73. Prandoni P, Polistena P, Bernardi E, et al. Upper-extremity deep vein thrombosis. Risk factors, diagnosis, and complications. *Arch Intern Med* 1997;157:57-62. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8996041>.
74. Verso M, Agnelli G. Venous thromboembolism associated with long-term use of central venous catheters in cancer patients. *J Clin Oncol* 2003;21:3665-3675. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14512399>.
75. Lee AYY, Levine MN, Butler G, et al. Incidence, risk factors, and outcomes of catheter-related thrombosis in adult patients with cancer. *J*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Clin Oncol 2006;24:1404-1408. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16549834>.

76. Joffe HV, Goldhaber SZ. Upper-extremity deep vein thrombosis. Circulation 2002;106:1874-1880. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12356644>.

77. Linenberger ML. Catheter-related thrombosis: risks, diagnosis, and management. J Natl Compr Canc Netw 2006;4:889-901. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17020667>.

78. van Rooden CJ, Schippers EF, Barge RMY, et al. Infectious complications of central venous catheters increase the risk of catheter-related thrombosis in hematology patients: a prospective study. J Clin Oncol 2005;23:2655-2660. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15837979>.

79. Kearon C, Ginsberg JS, Douketis J, et al. An evaluation of D-dimer in the diagnosis of pulmonary embolism: a randomized trial. Ann Intern Med 2006;144:812-821. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16754923>.

80. Wells PS, Anderson DR, Rodger M, et al. Evaluation of D-dimer in the diagnosis of suspected deep-vein thrombosis. N Engl J Med 2003;349:1227-1235. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14507948>.

81. Sohne M, Kruij MJHA, Nijkeuter M, et al. Accuracy of clinical decision rule, D-dimer and spiral computed tomography in patients with malignancy, previous venous thromboembolism, COPD or heart failure and in older patients with suspected pulmonary embolism. J Thromb Haemost 2006;4:1042-1046. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16689757>.

82. Knowlson L, Bacchu S, Paneesha S, et al. Elevated D-dimers are also a marker of underlying malignancy and increased mortality in the absence of venous thromboembolism. J Clin Pathol 2010;63:818-822. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20671046>.

83. Agnelli G, Verso M, Ageno W, et al. The MASTER registry on venous thromboembolism: description of the study cohort. Thromb Res 2008;121:605-610. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17692901>.

84. Zierler BK. Ultrasonography and diagnosis of venous thromboembolism. Circulation 2004;109:19-14. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15051663>.

85. Birdwell BG, Raskob GE, Whitsett TL, et al. The clinical validity of normal compression ultrasonography in outpatients suspected of having deep venous thrombosis. Ann Intern Med 1998;128:1-7. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/9424975>.

86. Kanne JP, Lalani TA. Role of computed tomography and magnetic resonance imaging for deep venous thrombosis and pulmonary embolism. Circulation 2004;109:15-21. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15051664>.

87. Lim K-E, Hsu W-C, Hsu Y-Y, et al. Deep venous thrombosis: comparison of indirect multidetector CT venography and sonography of lower extremities in 26 patients. Clin Imaging 2004;28:439-444. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15531146>.

88. Male C, Chait P, Ginsberg JS, et al. Comparison of venography and ultrasound for the diagnosis of asymptomatic deep vein thrombosis in the upper body in children: results of the PARKAA study. Prophylactic Antithrombin Replacement in Kids with ALL treated with Asparaginase. Thromb Haemost 2002;87:593-598. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12008940>.

89. Segal JB, Eng J, Jenckes MW, et al. Diagnosis and treatment of deep venous thrombosis and pulmonary embolism. Evid Rep Technol Assess (Summ) 2003:1-6. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12674745>.

90. Gaitini D. Current approaches and controversial issues in the diagnosis of deep vein thrombosis via duplex Doppler ultrasound. J Clin



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Ultrasound 2006;34:289-297. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16788961>.

91. Taffoni MJ, Ravenel JG, Ackerman SJ. Prospective comparison of indirect CT venography versus venous sonography in ICU patients. AJR Am J Roentgenol 2005;185:457-462. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16037520>.

92. Fraser DGW, Moody AR, Davidson IR, et al. Deep venous thrombosis: diagnosis by using venous enhanced subtracted peak arterial MR venography versus conventional venography. Radiology 2003;226:812-820. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12601180>.

93. Sampson FC, Goodacre SW, Thomas SM, van Beek EJ. The accuracy of MRI in diagnosis of suspected deep vein thrombosis: systematic review and meta-analysis. Eur Radiol 2007;17:175-181. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16628439>.

94. Joffe HV, Kucher N, Tapson VF, Goldhaber SZ. Upper-extremity deep vein thrombosis: a prospective registry of 592 patients. Circulation 2004;110:1605-1611. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15353493>.

95. Baarslag H-J, van Beek EJ, Koopman MMW, Reekers JA. Prospective study of color duplex ultrasonography compared with contrast venography in patients suspected of having deep venous thrombosis of the upper extremities. Ann Intern Med 2002;136:865-872. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12069560>.

96. Decousus H, Quere I, Presles E, et al. Superficial venous thrombosis and venous thromboembolism: a large, prospective epidemiologic study. Ann Intern Med 2010;152:218-224. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20157136>.

97. Lee JT, Kalani MA. Treating superficial venous thrombophlebitis. J Natl Compr Canc Netw 2008;6:760-765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18926088>.

98. Sack GH, Levin J, Bell WR. Trousseau's syndrome and other manifestations of chronic disseminated coagulopathy in patients with neoplasms: clinical, pathophysiologic, and therapeutic features. Medicine (Baltimore) 1977;56:1-37. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/834136>.

99. Varki A. Trousseau's syndrome: multiple definitions and multiple mechanisms. Blood 2007;110:1723-1729. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17496204>.

100. Martinelli I, Franchini M, Mannucci PM. How I treat rare venous thromboses. Blood 2008;112:4818-4823. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18805965>.

101. Thatipelli MR, McBane RD, Hodge DO, Wysokinski WE. Survival and recurrence in patients with splanchnic vein thromboses. Clin Gastroenterol Hepatol 2010;8:200-205. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19782767>.

102. Amitrano L, Guardascione MA, Scaglione M, et al. Prognostic factors in noncirrhotic patients with splanchnic vein thromboses. Am J Gastroenterol 2007;102:2464-2470. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17958760>.

103. Janssen HL, Wijnhoud A, Haagsma EB, et al. Extrahepatic portal vein thrombosis: aetiology and determinants of survival. Gut 2001;49:720-724. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11600478>.

104. Acosta S, Alhadad A, Svensson P, Ekberg O. Epidemiology, risk and prognostic factors in mesenteric venous thrombosis. Br J Surg 2008;95:1245-1251. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18720461>.

105. Hedayati N, Riha GM, Kougiyas P, et al. Prognostic factors and treatment outcome in mesenteric vein thrombosis. Vasc Endovascular Surg 2008;42:217-224. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18332399>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

106. De Stefano V, Martinelli I. Splanchnic vein thrombosis: clinical presentation, risk factors and treatment. *Intern Emerg Med* 2010;5:487-494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20532730>.
107. Janssen HL, Meinardi JR, Vlegaar FP, et al. Factor V Leiden mutation, prothrombin gene mutation, and deficiencies in coagulation inhibitors associated with Budd-Chiari syndrome and portal vein thrombosis: results of a case-control study. *Blood* 2000;96:2364-2368. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11001884>.
108. Kumar S, Sarr MG, Kamath PS. Mesenteric venous thrombosis. *N Engl J Med* 2001;345:1683-1688. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11759648>.
109. Connolly GC, Chen R, Hyrien O, et al. Incidence, risk factors and consequences of portal vein and systemic thromboses in hepatocellular carcinoma. *Thromb Res* 2008;122:299-306. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18045666>.
110. Greten TF, Papendorf F, Bleck JS, et al. Survival rate in patients with hepatocellular carcinoma: a retrospective analysis of 389 patients. *Br J Cancer* 2005;92:1862-1868. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15870713>.
111. Rabe C, Pilz T, Klostermann C, et al. Clinical characteristics and outcome of a cohort of 101 patients with hepatocellular carcinoma. *World J Gastroenterol* 2001;7:208-215. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11819762>.
112. Cabibbo G, Enea M, Attanasio M, et al. A meta-analysis of survival rates of untreated patients in randomized clinical trials of hepatocellular carcinoma. *Hepatology* 2010;51:1274-1283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20112254>.
113. Hoekstra J, Janssen HL. Vascular liver disorders (I): diagnosis, treatment and prognosis of Budd-Chiari syndrome. *Neth J Med* 2008;66:334-339. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18809980>.
114. Hoekstra J, Janssen HL. Vascular liver disorders (II): portal vein thrombosis. *Neth J Med* 2009;67:46-53. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19299846>.
115. Menon KV, Shah V, Kamath PS. The Budd-Chiari syndrome. *N Engl J Med* 2004;350:578-585. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14762185>.
116. Parikh S, Shah R, Kapoor P. Portal vein thrombosis. *Am J Med* 2010;123:111-119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20103016>.
117. Plessier A, Darwish-Murad S, Hernandez-Guerra M, et al. Acute portal vein thrombosis unrelated to cirrhosis: a prospective multicenter follow-up study. *Hepatology* 2010;51:210-218. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19821530>.
118. Sogaard KK, Astrup LB, Vilstrup H, Gronbaek H. Portal vein thrombosis; risk factors, clinical presentation and treatment. *BMC Gastroenterol* 2007;7:34. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17697371>.
119. Bradbury MS, Kavanagh PV, Bechtold RE, et al. Mesenteric venous thrombosis: diagnosis and noninvasive imaging. *Radiographics* 2002;22:527-541. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12006685>.
120. Ponziani FR, Zocco MA, Campanale C, et al. Portal vein thrombosis: insight into physiopathology, diagnosis, and treatment. *World J Gastroenterol* 2010;16:143-155. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20066733>.
121. Hillmen P, Lewis SM, Bessler M, et al. Natural history of paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 1995;333:1253-1258. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7566002>.
122. Socie G, Mary JY, de Gramont A, et al. Paroxysmal nocturnal haemoglobinuria: long-term follow-up and prognostic factors. *French*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Society of Haematology. Lancet 1996;348:573-577. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/8774569>.

123. Hoekstra J, Leebeek FW, Plessier A, et al. Paroxysmal nocturnal hemoglobinuria in Budd-Chiari syndrome: findings from a cohort study. J Hepatol 2009;51:696-706. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19664836>.

124. Baxter EJ, Scott LM, Campbell PJ, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. Lancet 2005;365:1054-1061. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15781101>.

125. James C, Ugo V, Le Couedic JP, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. Nature 2005;434:1144-1148. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15793561>.

126. Kralovics R, Passamonti F, Buser AS, et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. N Engl J Med 2005;352:1779-1790. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15858187>.

127. Levine RL, Wadleigh M, Cools J, et al. Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. Cancer Cell 2005;7:387-397. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15837627>.

128. Colaizzo D, Amitrano L, Tiscia GL, et al. The JAK2 V617F mutation frequently occurs in patients with portal and mesenteric venous thrombosis. J Thromb Haemost 2007;5:55-61. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17059429>.

129. De Stefano V, Fiorini A, Rossi E, et al. Incidence of the JAK2 V617F mutation among patients with splanchnic or cerebral venous thrombosis and without overt chronic myeloproliferative disorders. J Thromb Haemost 2007;5:708-714. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17263783>.

130. Regina S, Herault O, D'Alteroche L, et al. JAK2 V617F is specifically associated with idiopathic splanchnic vein thrombosis. J Thromb Haemost 2007;5:859-861. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17403204>.

131. Colaizzo D, Amitrano L, Tiscia GL, et al. A new JAK2 gene mutation in patients with polycythemia vera and splanchnic vein thrombosis. Blood 2007;110:2768-2769. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17881638>.

132. Fanikos J, Goldhaber SZ. Risk factors for the assessment of patients with pulmonary embolism. J Natl Compr Canc Netw 2006;4:871-880. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17020665>.

133. Kearon C, Ginsberg JS, Hirsh J. The role of venous ultrasonography in the diagnosis of suspected deep venous thrombosis and pulmonary embolism. Ann Intern Med 1998;129:1044-1049. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9867760>.

134. Moser KM, Fedullo PF, LitleJohn JK, Crawford R. Frequent asymptomatic pulmonary embolism in patients with deep venous thrombosis. JAMA 1994;271:223-225. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/8277550>.

135. O'Connell CL, Boswell WD, Duddalwar V, et al. Unsuspected pulmonary emboli in cancer patients: clinical correlates and relevance. J Clin Oncol 2006;24:4928-4932. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17050877>.

136. Worsley DF, Alavi A, Aronchick JM, et al. Chest radiographic findings in patients with acute pulmonary embolism: observations from the PIOPED Study. Radiology 1993;189:133-136. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/8372182>.

137. Costantini M, Bossone E, Renna R, et al. Electrocardiographic features in critical pulmonary embolism. Results from baseline and



continuous electrocardiographic monitoring. *Ital Heart J* 2004;5:214-216. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15119504>.

138. Ferrari E, Imbert A, Chevalier T, et al. The ECG in pulmonary embolism. Predictive value of negative T waves in precordial leads--80 case reports. *Chest* 1997;111:537-543. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9118684>.

139. Schoepf UJ, Costello P. CT angiography for diagnosis of pulmonary embolism: state of the art. *Radiology* 2004;230:329-337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14752178>.

140. Czekajska-Chehab E, Drop A, Terlecka B, et al. Indirect CT venography of the abdominal cavity and lower limbs in patients with the suspicion of pulmonary embolism--indications, technique, diagnostic possibilities. *Ann Univ Mariae Curie Sklodowska Med* 2004;59:508-518. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16146139>.

141. Schoepf UJ, Kucher N, Kipfmüller F, et al. Right ventricular enlargement on chest computed tomography: a predictor of early death in acute pulmonary embolism. *Circulation* 2004;110:3276-3280. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15533868>.

142. Anderson DR, Kahn SR, Rodger MA, et al. Computed tomographic pulmonary angiography vs ventilation-perfusion lung scanning in patients with suspected pulmonary embolism: a randomized controlled trial. *JAMA* 2007;298:2743-2753. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18165667>.

143. Calvo-Romero JM, Lima-Rodriguez EM, Bureo-Dacal P, Perez-Miranda M. Predictors of an intermediate ventilation/perfusion lung scan in patients with suspected acute pulmonary embolism. *Eur J Emerg Med* 2005;12:129-131. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15891446>.

144. Goldhaber SZ, Visani L, De Rosa M. Acute pulmonary embolism: clinical outcomes in the International Cooperative Pulmonary Embolism

Registry (ICOPER). *Lancet* 1999;353:1386-1389. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10227218>.

145. Jimenez D, Aujesky D, Moores L, et al. Combinations of prognostic tools for identification of high-risk normotensive patients with acute symptomatic pulmonary embolism. *Thorax* 2011;66:75-81. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20978032>.

146. Jimenez D, Aujesky D, Yusen RD. Risk stratification of normotensive patients with acute symptomatic pulmonary embolism. *Br J Haematol* 2010;151:415-424. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20955409>.

147. Kucher N, Goldhaber SZ. Cardiac biomarkers for risk stratification of patients with acute pulmonary embolism. *Circulation* 2003;108:2191-2194. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14597581>.

148. Kucher N, Goldhaber SZ. Management of massive pulmonary embolism. *Circulation* 2005;112:e28-32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16009801>.

149. Pruszczyk P, Torbicki A, Kuch-Wociał A, et al. Diagnostic value of transthoracic echocardiography in suspected haemodynamically significant pulmonary embolism. *Heart* 2001;85:628-634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11359740>.

150. Ribeiro A, Lindmarker P, Juhlin-Dannfelt A, et al. Echocardiography Doppler in pulmonary embolism: right ventricular dysfunction as a predictor of mortality rate. *Am Heart J* 1997;134:479-487. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9327706>.

151. Becattini C, Vedovati MC, Agnelli G. Prognostic value of troponins in acute pulmonary embolism: a meta-analysis. *Circulation* 2007;116:427-433. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17606843>.

152. Konstantinides S, Geibel A, Olschewski M, et al. Importance of cardiac troponins I and T in risk stratification of patients with acute



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

pulmonary embolism. *Circulation* 2002;106:1263-1268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12208803>.

153. Jimenez D, Aujesky D, Diaz G, et al. Prognostic significance of deep vein thrombosis in patients presenting with acute symptomatic pulmonary embolism. *Am J Respir Crit Care Med* 2010;181:983-991. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20110556>.

154. Aujesky D, Roy P-M, Le Manach CP, et al. Validation of a model to predict adverse outcomes in patients with pulmonary embolism. *Eur Heart J* 2006;27:476-481. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16207738>.

155. Donze J, Le Gal G, Fine MJ, et al. Prospective validation of the Pulmonary Embolism Severity Index. A clinical prognostic model for pulmonary embolism. *Thromb Haemost* 2008;100:943-948. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18989542>.

156. den Exter PL, Gomez V, Jimenez D, et al. A clinical prognostic model for the identification of low-risk patients with acute symptomatic pulmonary embolism and active cancer. *Chest* 2013;143:138-145. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22814859>.

157. LEO Pharmaceutical Products. Prescribing information: Innohep (tinzaparin sodium injection) for subcutaneous (SC) use only. 2008. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2008/020484s011lbl.pdf. Accessed September 16, 2014.

158. sanofi-aventis U.S. LLC. Prescribing information: Lovenox (enoxaparin sodium injection). 2009. Available at: <http://products.sanofi-aventis.us/lovenox/lovenox.html>. Accessed September 16, 2014.

159. GlaxoSmithKline. Prescribing information: ARIXTRA (fondaparinux sodium) Solution for subcutaneous injection. 2011. Available at: http://us.gsk.com/products/assets/us_arixtra.pdf. Accessed September 16, 2014.

160. Eisai, Inc; Pfizer, Inc. Prescribing information: FRAGMIN (dalteparin sodium injection) for Subcutaneous Use Only. 2010. Available at:

http://www.accessdata.fda.gov/drugsatfda_docs/label/2010/020287s050lbl.pdf. Accessed September 16, 2014.

161. Hospira, Inc. Prescribing information: Heparin sodium injection, USP. 2008. Available at:

http://www.hospira.com/products_and_services/drugs/HEPARIN_SODIUM_INJECTION. Accessed September 16, 2014.

162. Cervera R, Khamashta MA, Shoenfeld Y, et al. Morbidity and mortality in the antiphospholipid syndrome during a 5-year period: a multicentre prospective study of 1000 patients. *Ann Rheum Dis* 2009;68:1428-1432. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18801761>.

163. Lim W, Crowther MA, Eikelboom JW. Management of antiphospholipid antibody syndrome: a systematic review. *JAMA* 2006;295:1050-1057. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16507806>.

164. Lim W. Antiphospholipid antibody syndrome. *Hematology Am Soc Hematol Educ Program* 2009:233-239. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20008203>.

165. Hutten BA, Prins MH, Gent M, et al. Incidence of recurrent thromboembolic and bleeding complications among patients with venous thromboembolism in relation to both malignancy and achieved international normalized ratio: a retrospective analysis. *J Clin Oncol* 2000;18:3078-3083. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10963635>.

166. Palareti G, Legnani C, Lee A, et al. A comparison of the safety and efficacy of oral anticoagulation for the treatment of venous thromboembolic disease in patients with or without malignancy. *Thromb Haemost* 2000;84:805-810. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11127860>.

167. Hull RD, Pineo GF, Brant RF, et al. Long-term low-molecular-weight heparin versus usual care in proximal-vein thrombosis patients with cancer. *Am J Med* 2006;119:1062-1072. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17145251>.

168. Lee AYY, Levine MN, Baker RI, et al. Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. *N Engl J Med* 2003;349:146-153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12853587>.

169. Meyer G, Marjanovic Z, Valcke J, et al. Comparison of low-molecular-weight heparin and warfarin for the secondary prevention of venous thromboembolism in patients with cancer: a randomized controlled study. *Arch Intern Med* 2002;162:1729-1735. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12153376>.

170. Wawrzynska L, Tomkowski WZ, Przedlacki J, et al. Changes in bone density during long-term administration of low-molecular-weight heparins or acenocoumarol for secondary prophylaxis of venous thromboembolism. *Pathophysiol Haemost Thromb* 2003;33:64-67. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14624046>.

171. Aronson J. Serious drug interactions. *Practitioner* 1993;237:789-791. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7903448>.

172. Lacey CS. Interaction of dicloxacillin with warfarin. *Ann Pharmacother* 2004;38:898-898. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15054148>.

173. Saif MW. An adverse interaction between warfarin and fluoropyrimidines revisited. *Clin Colorectal Cancer* 2005;5:175-180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16197620>.

174. Shah HR, Ledbetter L, Diasio R, Saif MW. A retrospective study of coagulation abnormalities in patients receiving concomitant capecitabine and warfarin. *Clin Colorectal Cancer* 2006;5:354-358. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16512995>.

175. Morello KC, Wurz GT, DeGregorio MW. Pharmacokinetics of selective estrogen receptor modulators. *Clin Pharmacokinet* 2003;42:361-372. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12648026>.

176. Nutescu EA, Shapiro NL, Ibrahim S, West P. Warfarin and its interactions with foods, herbs and other dietary supplements. *Expert Opin Drug Saf* 2006;5:433-451. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16610971>.

177. Sconce E, Khan T, Mason J, et al. Patients with unstable control have a poorer dietary intake of vitamin K compared to patients with stable control of anticoagulation. *Thromb Haemost* 2005;93:872-875. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15886802>.

178. Wittkowsky AK, Boccuzzi SJ, Wogen J, et al. Frequency of concurrent use of warfarin with potentially interacting drugs. *Pharmacotherapy* 2004;24:1668-1674. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15585436>.

179. Barritt DW, Jordan SC. Anticoagulant drugs in the treatment of pulmonary embolism. A controlled trial. *Lancet* 1960;1:1309-1312. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/13797091>.

180. Hirsh J, Bates SM. Clinical trials that have influenced the treatment of venous thromboembolism: a historical perspective. *Ann Intern Med* 2001;134:409-417. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11242501>.

181. Garcia DA, Baglin TP, Weitz JI, Samama MM. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012;141:e24S-43S. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22315264>.

182. Kahn SR, Lim W, Dunn AS, et al. Prevention of VTE in nonsurgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Guidelines. Chest 2012;141:e195S-226S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22315261>.

183. Kearon C, Akl EA, Comerota AJ, et al. Antithrombotic therapy for VTE disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:e419S-494S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22315268>.

184. Cheer SM, Dunn CJ, Foster R. Tinzaparin sodium: a review of its pharmacology and clinical use in the prophylaxis and treatment of thromboembolic disease. Drugs 2004;64:1479-1502. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15212562>.

185. Neely JL, Carlson SS, Lenhart SE. Tinzaparin sodium: a low-molecular-weight heparin. Am J Health Syst Pharm 2002;59:1426-1436.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12166042>.

186. Nutescu EA, Shapiro NL, Feinstein H, Rivers CW. Tinzaparin: considerations for use in clinical practice. Ann Pharmacother 2003;37:1831-1840. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/14632588>.

187. Planes A, Samama MM, Lensing AW, et al. Prevention of deep vein thrombosis after hip replacement--comparison between two low-molecular heparins, tinzaparin and enoxaparin. Thromb Haemost 1999;81:22-25. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10348714>.

188. Wells PS, Anderson DR, Rodger MA, et al. A randomized trial comparing 2 low-molecular-weight heparins for the outpatient treatment of deep vein thrombosis and pulmonary embolism. Arch Intern Med 2005;165:733-738. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15824291>.

189. Mandalà M, Falanga A, Roila F. Management of venous thromboembolism in cancer patients: ESMO clinical recommendations.

Ann Oncol 2008;19 Suppl 2:126-127. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18456750>.

190. Cook LM, Kahn SR, Goodwin J, Kovacs MJ. Frequency of renal impairment, advanced age, obesity and cancer in venous thromboembolism patients in clinical practice. J Thromb Haemost 2007;5:937-941. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17461927>.

191. Hirsh J, Bauer KA, Donati MB, et al. Parenteral anticoagulants: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). Chest 2008;133:141S-159S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18574264>.

192. Michota F, Merli G. Anticoagulation in special patient populations: are special dosing considerations required? Cleve Clin J Med 2005;72 Suppl 1:S37-42. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15853178>.

193. Sanderink G-JCM, Guimart CG, Ozoux M-L, et al. Pharmacokinetics and pharmacodynamics of the prophylactic dose of enoxaparin once daily over 4 days in patients with renal impairment. Thromb Res 2002;105:225-231. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11927128>.

194. Lim W, Dentali F, Eikelboom JW, Crowther MA. Meta-analysis: low-molecular-weight heparin and bleeding in patients with severe renal insufficiency. Ann Intern Med 2006;144:673-684. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16670137>.

195. Hulot J-S, Montalescot G, Lechat P, et al. Dosing strategy in patients with renal failure receiving enoxaparin for the treatment of non-ST-segment elevation acute coronary syndrome. Clin Pharmacol Ther 2005;77:542-552. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15961985>.

196. Kruse MW, Lee JJ. Retrospective evaluation of a pharmacokinetic program for adjusting enoxaparin in renal impairment. Am Heart J



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

2004;148:582-589. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15459586>.

197. Shprecher AR, Cheng-Lai A, Madsen EM, et al. Peak antifactor xa activity produced by dalteparin treatment in patients with renal impairment compared with controls. *Pharmacotherapy* 2005;25:817-822. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15927900>.

198. Douketis J, Cook D, Meade M, et al. Prophylaxis against deep vein thrombosis in critically ill patients with severe renal insufficiency with the low-molecular-weight heparin dalteparin: an assessment of safety and pharmacodynamics: the DIRECT study. *Arch Intern Med*

2008;168:1805-1812. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18779469>.

199. Kucher N, Leizorovicz A, Vaitkus PT, et al. Efficacy and safety of fixed low-dose dalteparin in preventing venous thromboembolism among obese or elderly hospitalized patients: a subgroup analysis of the PREVENT trial. *Arch Intern Med* 2005;165:341-345. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15710801>.

200. Prandoni P. How I treat venous thromboembolism in patients with cancer. *Blood* 2005;106:4027-4033. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16076870>.

201. Savi P, Chong BH, Greinacher A, et al. Effect of fondaparinux on platelet activation in the presence of heparin-dependent antibodies: a blinded comparative multicenter study with unfractionated heparin. *Blood* 2005;105:139-144. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15388575>.

202. Warkentin TE, Greinacher A, Koster A, Lincoff AM. Treatment and prevention of heparin-induced thrombocytopenia: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* 2008;133:340S-380S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18574270>.

203. Cohen AT, Davidson BL, Gallus AS, et al. Efficacy and safety of fondaparinux for the prevention of venous thromboembolism in older acute medical patients: randomised placebo controlled trial. *BMJ* 2006;332:325-329. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16439370>.

204. Janssen Pharmaceuticals, Inc. Prescribing Information: XARELTO (rivaroxaban) tablets. 2011. Available at:

http://www.xareltohcp.com/sites/default/files/pdf/xarelto_0.pdf#zoom=100. Accessed September 16, 2014.

205. Bauersachs R, Berkowitz SD, Brenner B, et al. Oral rivaroxaban for symptomatic venous thromboembolism. *N Engl J Med* 2010;363:2499-2510. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21128814>.

206. Buller HR, Prins MH, Lensin AW, et al. Oral rivaroxaban for the treatment of symptomatic pulmonary embolism. *N Engl J Med* 2012;366:1287-1297. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22449293>.

207. Cohen AT, Spiro TE, Buller HR, et al. Rivaroxaban for thromboprophylaxis in acutely ill medical patients. *N Engl J Med* 2013;368:513-523. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23388003>.

208. Bristol-Myers Squibb Company. Prescribing Information: ELIQUIS (apixaban) tablets for oral use. 2012. Available at:

http://packageinserts.bms.com/pi/pi_eliquis.pdf. Accessed September 16, 2014.

209. Agnelli G, Buller HR, Cohen A, et al. Oral Apixaban for the Treatment of Acute Venous Thromboembolism. *N Engl J Med* 2013. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23808982>.

210. Lassen MR, Gallus A, Raskob GE, et al. Apixaban versus enoxaparin for thromboprophylaxis after hip replacement. *N Engl J Med*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

2010;363:2487-2498. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21175312>.

211. Lassen MR, Raskob GE, Gallus A, et al. Apixaban versus enoxaparin for thromboprophylaxis after knee replacement (ADVANCE-2): a randomised double-blind trial. *Lancet* 2010;375:807-815. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20206776>.

212. Goldhaber SZ, Leizorovicz A, Kakkar AK, et al. Apixaban versus enoxaparin for thromboprophylaxis in medically ill patients. *N Engl J Med* 2011;365:2167-2177. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22077144>.

213. Agnelli G, Buller HR, Cohen A, et al. Apixaban for extended treatment of venous thromboembolism. *N Engl J Med* 2013;368:699-708. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23216615>.

214. Hirsh J, Anand SS, Halperin JL, Fuster V. Mechanism of action and pharmacology of unfractionated heparin. *Arterioscler Thromb Vasc Biol* 2001;21:1094-1096. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11451734>.

215. Clagett GP, Reisch JS. Prevention of venous thromboembolism in general surgical patients. Results of meta-analysis. *Ann Surg* 1988;208:227-240. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2456748>.

216. Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* 2008;133:453. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18574271>.

217. King CS, Holley AB, Jackson JL, et al. Twice vs three times daily heparin dosing for thromboembolism prophylaxis in the general medical population: A metaanalysis. *Chest* 2007;131:507-516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17296655>.

218. Kearon C, Ginsberg JS, Julian JA, et al. Comparison of fixed-dose weight-adjusted unfractionated heparin and low-molecular-weight heparin for acute treatment of venous thromboembolism. *JAMA* 2006;296:935-942. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16926353>.

219. Bristol-Myers Squibb. Prescribing information: COUMADIN® TABLETS (Warfarin Sodium Tablets, USP); COUMADIN® FOR INJECTION (Warfarin Sodium for Injection, USP). 2010. Available at: http://packageinserts.bms.com/pi/pi_coumadin.pdf. Accessed September 16, 2014.

220. Glynn RJ, Ridker PM, Goldhaber SZ, Buring JE. Effect of low-dose aspirin on the occurrence of venous thromboembolism: a randomized trial. *Ann Intern Med* 2007;147:525-533. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17938390>.

221. Becattini C, Agnelli G, Poggio R, et al. Aspirin After Oral Anticoagulants for Prevention of Recurrence in Patients with Unprovoked Venous Thromboembolism. the Warfasa STUDY [abstract]. *Blood* 2011;118:Abstract 543. Available at: <http://abstracts.hematologylibrary.org/cgi/content/abstract/ashmtg;118/2/1/543>.

222. Becattini C, Agnelli G, Schenone A, et al. Aspirin for preventing the recurrence of venous thromboembolism. *N Engl J Med* 2012;366:1959-1967. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22621626>.

223. Eight-year follow-up of patients with permanent vena cava filters in the prevention of pulmonary embolism: the PREPIC (Prevention du Risque d'Embolie Pulmonaire par Interruption Cave) randomized study. *Circulation* 2005;112:416-422. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16009794>.

224. Brender E. Use of emboli-blocking filters increases, but rigorous data are lacking. *JAMA* 2006;295:989-990. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16507791>.

225. Stein PD, Kayali F, Olson RE. Twenty-one-year trends in the use of inferior vena cava filters. *Arch Intern Med* 2004;164:1541-1545. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15277286>.

226. Streiff MB. Vena caval filters: a comprehensive review. *Blood* 2000;95:3669-3677. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10845895>.

227. Streiff MB. Vena caval filters: a review for intensive care specialists. *J Intensive Care Med* 2003;18:59-79. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15189653>.

228. Decousus H, Leizorovicz A, Parent F, et al. A clinical trial of vena caval filters in the prevention of pulmonary embolism in patients with proximal deep-vein thrombosis. Prevention du Risque d'Embolie Pulmonaire par Interruption Cave Study Group. *N Engl J Med* 1998;338:409-415. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9459643>.

229. Elting LS, Escalante CP, Cooksley C, et al. Outcomes and cost of deep venous thrombosis among patients with cancer. *Arch Intern Med* 2004;164:1653-1661. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15302635>.

230. Getzen TM, Rectenwald JE. Inferior vena cava filters in the cancer patient: current use and indications. *J Natl Compr Canc Netw* 2006;4:881-888. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17020666>.

231. Millward SF, Grassi CJ, Kinney TB, et al. Reporting standards for inferior vena caval filter placement and patient follow-up: supplement for temporary and retrievable/optional filters. *J Vasc Interv Radiol* 2005;16:441-443. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15802441>.

232. Kim HS, Young MJ, Narayan AK, et al. A comparison of clinical outcomes with retrievable and permanent inferior vena cava filters. *J*

Vasc Interv Radiol 2008;19:393-399. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18295699>.

233. Nicholson W, Nicholson WJ, Tolerico P, et al. Prevalence of fracture and fragment embolization of Bard retrievable vena cava filters and clinical implications including cardiac perforation and tamponade. *Arch Intern Med* 2010;170:1827-1831. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20696949>.

234. Francis CW. Prevention of venous thromboembolism in hospitalized patients with cancer. *J Clin Oncol* 2009;27:4874-4880. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19704060>.

235. Efficacy and safety of enoxaparin versus unfractionated heparin for prevention of deep vein thrombosis in elective cancer surgery: a double-blind randomized multicentre trial with venographic assessment. ENOXACAN Study Group. *Br J Surg* 1997;84:1099-1103. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9278651>.

236. Mismetti P, Laporte S, Darmon JY, et al. Meta-analysis of low molecular weight heparin in the prevention of venous thromboembolism in general surgery. *Br J Surg* 2001;88:913-930. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11442521>.

237. DeBernardo RL, Perkins RB, Littell RD, et al. Low-molecular-weight heparin (dalteparin) in women with gynecologic malignancy. *Obstet Gynecol* 2005;105:1006-1011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15863537>.

238. Akl EA, Terrenato I, Barba M, et al. Low-molecular-weight heparin vs unfractionated heparin for perioperative thromboprophylaxis in patients with cancer: a systematic review and meta-analysis. *Arch Intern Med* 2008;168:1261-1269. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18574082>.

239. Couban S, Goodyear M, Burnell M, et al. Randomized placebo-controlled study of low-dose warfarin for the prevention of central venous catheter-associated thrombosis in patients with cancer. *J Clin*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Oncol 2005;23:4063-4069. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15767639>.

240. Karthaus M, Kretschmar A, Kroning H, et al. Dalteparin for prevention of catheter-related complications in cancer patients with central venous catheters: final results of a double-blind, placebo-controlled phase III trial. *Ann Oncol* 2006;17:289-296. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16317012>.

241. Verso M, Agnelli G, Bertoglio S, et al. Enoxaparin for the prevention of venous thromboembolism associated with central vein catheter: a double-blind, placebo-controlled, randomized study in cancer patients. *J Clin Oncol* 2005;23:4057-4062. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15767643>.

242. Young AM, Billingham LJ, Begum G, et al. Warfarin thromboprophylaxis in cancer patients with central venous catheters (WARP): an open-label randomised trial. *Lancet* 2009;373:567-574. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19217991>.

243. Khorana AA, Dalal M, Tangirala K, Miao R. Higher Incidence of Venous Thromboembolism in the Outpatient Versus the Inpatient Setting Among U.S. Cancer Patients [abstract]. *Blood* 2011;118:Abstract 674. Available at:

<http://abstracts.hematologylibrary.org/cgi/content/abstract/118/21/674>.

244. Martino MA, Borges E, Williamson E, et al. Pulmonary embolism after major abdominal surgery in gynecologic oncology. *Obstet Gynecol* 2006;107:666-671. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16507939>.

245. Bergqvist D, Agnelli G, Cohen AT, et al. Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *N Engl J Med* 2002;346:975-980. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11919306>.

246. Rasmussen MS, Jorgensen LN, Wille-Jorgensen P, et al. Prolonged prophylaxis with dalteparin to prevent late thromboembolic

complications in patients undergoing major abdominal surgery: a multicenter randomized open-label study. *J Thromb Haemost* 2006;4:2384-2390. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16881934>.

247. Rana P, Levine MN. Prevention of thrombosis in ambulatory patients with cancer. *J Clin Oncol* 2009;27:4885-4888. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19752331>.

248. Zangari M, Fink LM, Elice F, et al. Thrombotic events in patients with cancer receiving antiangiogenesis agents. *J Clin Oncol* 2009;27:4865-4873. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19704059>.

249. Gay F, Hayman SR, Lacy MQ, et al. Lenalidomide plus dexamethasone versus thalidomide plus dexamethasone in newly diagnosed multiple myeloma: a comparative analysis of 411 patients. *Blood* 2010;115:1343-1350. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20008302>.

250. Rajkumar SV, Jacobus S, Callander NS, et al. Lenalidomide plus high-dose dexamethasone versus lenalidomide plus low-dose dexamethasone as initial therapy for newly diagnosed multiple myeloma: an open-label randomised controlled trial. *Lancet Oncol* 2010;11:29-37. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19853510>.

251. Celgene Corporation. Prescribing information: REVLIMID (lenalidomide) capsules. 2010. Available at:

<http://www.revlimid.com/wp-content/uploads/2013/11/PI.pdf>. Accessed September 16, 2014.

252. Celgene Corporation. Prescribing information: THALOMID® (thalidomide) Capsules 50 mg, 100 mg, 150 mg & 200 mg. 2010. Available at:

http://www.thalomid.com/pdf/Thalomid_PI.pdf. Accessed September 16, 2014.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

253. Palumbo A, Cavo M, Brinthen S, et al. Aspirin, Warfarin, or Enoxaparin Thromboprophylaxis in Patients With Multiple Myeloma Treated With Thalidomide: A Phase III, Open-Label, Randomized Trial. *J Clin Oncol* 2011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21282540>.

254. Larocca A, Cavallo F, Brinthen S, et al. Aspirin or enoxaparin thromboprophylaxis for patients with newly diagnosed multiple myeloma treated with lenalidomide. *Blood* 2012;119:933-939; quiz 1093. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21835953>.

255. Kearney JC, Rossi S, Glinert K, Henry DH. Venous Thromboembolism (VTE) and Survival in a Cancer Chemotherapy Outpatient Clinic: A Retrospective Chart Review Validation of a VTE Predictive Model [abstract]. *Blood* 2009;114:Abstract 2503. Available at: <http://abstracts.hematologylibrary.org/cgi/content/abstract/ashmtg;114/2/2503>.

256. Price LH, Nguyen MB, Picozzi VJ, Kozarek RA. Portal vein thrombosis in pancreatic cancer: Natural history, risk factors, and implications for patient management [abstract]. *Proceedings of the Gastrointestinal Cancers Symposium 2010*:Abstract 143. Available at: <https://meetinglibrary.asco.org/content/1359-72>.

257. Agnelli G, Gussoni G, Bianchini C, et al. Nadroparin for the prevention of thromboembolic events in ambulatory patients with metastatic or locally advanced solid cancer receiving chemotherapy: a randomised, placebo-controlled, double-blind study. *Lancet Oncol* 2009;10:943-949. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19726226>.

258. Riess H, Pelzer U, Opitz B, et al. A prospective, randomized trial of simultaneous pancreatic cancer treatment with enoxaparin and chemotherapy: Final results of the CONKO-004 trial [abstract]. *J Clin Oncol* 2010;28(Suppl 15):Abstract 4033. Available at: http://meeting.ascopubs.org/cgi/content/abstract/28/15_suppl/4033.

259. Agnelli G, George DJ, Kakkar AK, et al. Semuloparin for thromboprophylaxis in patients receiving chemotherapy for cancer. *N Engl J Med* 2012;366:601-609. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22335737>.

260. Cohen AT, Nandini B, Wills JO, Ota S. VTE prophylaxis for the medical patient: where do we stand? - a focus on cancer patients. *Thromb Res* 2010;125 Suppl 2:S21-29. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20434000>.

261. Clarke-Pearson DL, Synan IS, Dodge R, et al. A randomized trial of low-dose heparin and intermittent pneumatic calf compression for the prevention of deep venous thrombosis after gynecologic oncology surgery. *Am J Obstet Gynecol* 1993;168:1146-1153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8475960>.

262. Ramirez JI, Vassiliu P, Gonzalez-Ruiz C, et al. Sequential compression devices as prophylaxis for venous thromboembolism in high-risk colorectal surgery patients: reconsidering American Society of Colorectal Surgeons parameters. *Am Surg* 2003;69:941-945. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14627252>.

263. Amaragiri SV, Lees TA. Elastic compression stockings for prevention of deep vein thrombosis. *Cochrane Database Syst Rev* 2000:CD001484. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10908501>.

264. Cohen AT, Skinner JA, Warwick D, Brenkel I. The use of graduated compression stockings in association with fondaparinux in surgery of the hip. A multicentre, multinational, randomised, open-label, parallel-group comparative study. *J Bone Joint Surg Br* 2007;89:887-892. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17673580>.

265. Dennis M, Sandercock PA, Reid J, et al. Effectiveness of thigh-length graduated compression stockings to reduce the risk of deep vein thrombosis after stroke (CLOTS trial 1): a multicentre, randomised controlled trial. *Lancet* 2009;373:1958-1965. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19477503>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

[NCCN Guidelines Index](#)
[VTE Table of Contents](#)
[Discussion](#)

266. Jaff MR, McMurtry MS, Archer SL, et al. Management of Massive and Submassive Pulmonary Embolism, Iliofemoral Deep Vein Thrombosis, and Chronic Thromboembolic Pulmonary Hypertension: A Scientific Statement From the American Heart Association. *Circulation* 2011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21422387>.

267. Kearon C. Natural history of venous thromboembolism. *Circulation* 2003;107:122-30. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12814982>.

268. Kahn SR, Hirsch A, Shrier I. Effect of postthrombotic syndrome on health-related quality of life after deep venous thrombosis. *Arch Intern Med* 2002;162:1144-1148. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12020185>.

269. Prandoni P, Lensing AW, Cogo A, et al. The long-term clinical course of acute deep venous thrombosis. *Ann Intern Med* 1996;125:1-7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8644983>.

270. Comerota AJ, Aldridge SC. Thrombolytic therapy for deep venous thrombosis: a clinical review. *Can J Surg* 1993;36:359-364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8370018>.

271. Goldhaber SZ, Meyerovitz MF, Green D, et al. Randomized controlled trial of tissue plasminogen activator in proximal deep venous thrombosis. *Am J Med* 1990;88:235-240. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2106783>.

272. Schweizer J, Kirch W, Koch R, et al. Short- and long-term results after thrombolytic treatment of deep venous thrombosis. *J Am Coll Cardiol* 2000;36:1336-1343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11028492>.

273. Turpie AG, Levine MN, Hirsh J, et al. Tissue plasminogen activator (rt-PA) vs heparin in deep vein thrombosis. Results of a randomized trial. *Chest* 1990;97:172S-175S. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2108855>.

274. Verhaeghe R, Besse P, Bounameaux H, Marbet GA. Multicenter pilot study of the efficacy and safety of systemic rt-PA administration in the treatment of deep vein thrombosis of the lower extremities and/or pelvis. *Thromb Res* 1989;55:5-11. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2506661>.

275. Watson LI, Armon MP. Thrombolysis for acute deep vein thrombosis. *Cochrane Database Syst Rev* 2004;CD002783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15495034>.

276. Mewissen MW, Seabrook GR, Meissner MH, et al. Catheter-directed thrombolysis for lower extremity deep venous thrombosis: report of a national multicenter registry. *Radiology* 1999;211:39-49. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10189452>.

277. Castaneda F, Li R, Young K, et al. Catheter-directed thrombolysis in deep venous thrombosis with use of reteplase: immediate results and complications from a pilot study. *J Vasc Interv Radiol* 2002;13:577-580. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12050297>.

278. Grunwald MR, Hofmann LV. Comparison of urokinase, alteplase, and reteplase for catheter-directed thrombolysis of deep venous thrombosis. *J Vasc Interv Radiol* 2004;15:347-352. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15064337>.

279. Razavi MK, Wong H, Kee ST, et al. Initial clinical results of tenecteplase (TNK) in catheter-directed thrombolytic therapy. *J Endovasc Ther* 2002;9:593-598. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12431142>.

280. Enden T, Klow NE, Sandvik L, et al. Catheter-directed thrombolysis vs. anticoagulant therapy alone in deep vein thrombosis: results of an open randomized, controlled trial reporting on short-term patency. *J Thromb Haemost* 2009;7:1268-1275. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19422443>.

281. Enden T, Haig Y, Klow NE, et al. Long-term outcome after additional catheter-directed thrombolysis versus standard treatment for



acute iliofemoral deep vein thrombosis (the CaVenT study): a randomised controlled trial. *Lancet* 2012;379:31-38. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22172244>.

282. Kahn SR, Shapiro S, Wells PS, et al. Compression stockings to prevent post-thrombotic syndrome: a randomised placebo-controlled trial. *Lancet* 2014;383:880-888. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24315521>.

283. Kim HS, Preece SR, Black JH, et al. Safety of catheter-directed thrombolysis for deep venous thrombosis in cancer patients. *J Vasc Surg* 2008;47:388-394. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18241762>.

284. Di Nisio M, Ferrante N, De Tursi M, et al. Incidental venous thromboembolism in ambulatory cancer patients receiving chemotherapy. *Thromb Haemost* 2010;104:1049-1054. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20806119>.

285. Khorana AA, Streiff MB, Farge D, et al. Venous thromboembolism prophylaxis and treatment in cancer: a consensus statement of major guidelines panels and call to action. *J Clin Oncol* 2009;27:4919-4926. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19720907>.

286. Akl EA, Vasireddi SR, Gunukula S, et al. Anticoagulation for the initial treatment of venous thromboembolism in patients with cancer. *Cochrane Database Syst Rev* 2011;2:CD006649. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21328285>.

287. Deitcher SR, Kessler CM, Merli G, et al. Secondary prevention of venous thromboembolic events in patients with active cancer: enoxaparin alone versus initial enoxaparin followed by warfarin for a 180-day period. *Clin Appl Thromb Hemost* 2006;12:389-396. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17000884>.

288. Akl EA, Barba M, Rohilla S, et al. Anticoagulation for the long term treatment of venous thromboembolism in patients with cancer.

Cochrane Database Syst Rev 2008:CD006650. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18425959>.

289. Kakkar AK, Levine MN, Kadziola Z, et al. Low molecular weight heparin, therapy with dalteparin, and survival in advanced cancer: the fragmin advanced malignancy outcome study (FAMOUS). *J Clin Oncol* 2004;22:1944-1948. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15143088>.

290. Lee AYY, Rickles FR, Julian JA, et al. Randomized comparison of low molecular weight heparin and coumarin derivatives on the survival of patients with cancer and venous thromboembolism. *J Clin Oncol* 2005;23:2123-2129. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15699480>.

291. Altinbas M, Coskun HS, Er O, et al. A randomized clinical trial of combination chemotherapy with and without low-molecular-weight heparin in small cell lung cancer. *J Thromb Haemost* 2004;2:1266-1271. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15304029>.

292. Klerk CP, Smorenburg SM, Otten HM, et al. The effect of low molecular weight heparin on survival in patients with advanced malignancy. *J Clin Oncol* 2005;23:2130-2135. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15699479>.

293. Akl EA, van Doormaal FF, Barba M, et al. Parenteral anticoagulation may prolong the survival of patients with limited small cell lung cancer: a Cochrane systematic review. *J Exp Clin Cancer Res* 2008;27:4. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18577254>.

294. Kovacs MJ, Kahn SR, Rodger M, et al. A pilot study of central venous catheter survival in cancer patients using low-molecular-weight heparin (dalteparin) and warfarin without catheter removal for the treatment of upper extremity deep vein thrombosis (The Catheter Study). *J Thromb Haemost* 2007;5:1650-1653. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17488349>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

295. Leon L, Giannoukas AD, Dodd D, et al. Clinical significance of superficial vein thrombosis. *Eur J Vasc Endovasc Surg* 2005;29:10-17. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15570265>.

296. van Weert H, Dolan G, Wichers I, et al. Spontaneous superficial venous thrombophlebitis: does it increase risk for thromboembolism? A historic follow-up study in primary care. *J Fam Pract* 2006;55:52-57. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16388768>.

297. Marchiori A, Verlato F, Sabbion P, et al. High versus low doses of unfractionated heparin for the treatment of superficial thrombophlebitis of the leg. A prospective, controlled, randomized study. *Haematologica* 2002;87:523-527. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12010667>.

298. A pilot randomized double-blind comparison of a low-molecular-weight heparin, a nonsteroidal anti-inflammatory agent, and placebo in the treatment of superficial vein thrombosis. *Arch Intern Med* 2003;163:1657-1663. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12885680>.

299. Decousus H, Prandoni P, Mismetti P, et al. Fondaparinux for the treatment of superficial-vein thrombosis in the legs. *N Engl J Med* 2010;363:1222-1232. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20860504>.

300. Condat B, Pessione F, Hillaire S, et al. Current outcome of portal vein thrombosis in adults: risk and benefit of anticoagulant therapy. *Gastroenterology* 2001;120:490-497. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11159889>.

301. Dentali F, Ageno W, Witt D, et al. Natural history of mesenteric venous thrombosis in patients treated with vitamin K antagonists: a multi-centre, retrospective cohort study. *Thromb Haemost* 2009;102:501-504. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19718470>.

302. Hollingshead M, Burke CT, Mauro MA, et al. Transcatheter thrombolytic therapy for acute mesenteric and portal vein thrombosis. *J Vasc Interv Radiol* 2005;16:651-661. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15872320>.

303. Kim HS, Patra A, Khan J, et al. Transhepatic catheter-directed thrombectomy and thrombolysis of acute superior mesenteric venous thrombosis. *J Vasc Interv Radiol* 2005;16:1685-1691. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16371536>.

304. Sharma S, Texeira A, Texeira P, et al. Pharmacological thrombolysis in Budd Chiari syndrome: a single centre experience and review of the literature. *J Hepatol* 2004;40:172-180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14672630>.

305. Smalberg JH, Spaander MV, Jie KS, et al. Risks and benefits of transcatheter thrombolytic therapy in patients with splanchnic venous thrombosis. *Thromb Haemost* 2008;100:1084-1088. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19132234>.

306. Garcia-Pagan JC, Heydtmann M, Raffa S, et al. TIPS for Budd-Chiari syndrome: long-term results and prognostic factors in 124 patients. *Gastroenterology* 2008;135:808-815. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18621047>.

307. Darwish Murad S, Plessier A, Hernandez-Guerra M, et al. Etiology, management, and outcome of the Budd-Chiari syndrome. *Ann Intern Med* 2009;151:167-175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19652186>.

308. Eapen CE, Velissaris D, Heydtmann M, et al. Favourable medium term outcome following hepatic vein recanalisation and/or transjugular intrahepatic portosystemic shunt for Budd Chiari syndrome. *Gut* 2006;55:878-884. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16174658>.

309. Mancuso A, Fung K, Mela M, et al. TIPS for acute and chronic Budd-Chiari syndrome: a single-centre experience. *J Hepatol*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

2003;38:751-754. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12763367>.

310. Perello A, Garcia-Pagan JC, Gilabert R, et al. TIPS is a useful long-term derivative therapy for patients with Budd-Chiari syndrome uncontrolled by medical therapy. *Hepatology* 2002;35:132-139.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11786969>.

311. Rossle M, Olschewski M, Siegerstetter V, et al. The Budd-Chiari syndrome: outcome after treatment with the transjugular intrahepatic portosystemic shunt. *Surgery* 2004;135:394-403. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15041963>.

312. Hemming AW, Langer B, Greig P, et al. Treatment of Budd-Chiari syndrome with portosystemic shunt or liver transplantation. *Am J Surg* 1996;171:176-180; discussion 180-171. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8554136>.

313. Zeitoun G, Escolano S, Hadengue A, et al. Outcome of Budd-Chiari syndrome: a multivariate analysis of factors related to survival including surgical portosystemic shunting. *Hepatology* 1999;30:84-89.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10385643>.

314. Panis Y, Belghiti J, Valla D, et al. Portosystemic shunt in Budd-Chiari syndrome: long-term survival and factors affecting shunt patency in 25 patients in Western countries. *Surgery* 1994;115:276-281.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8128351>.

315. Pisani-Ceretti A, Intra M, Prestipino F, et al. Surgical and radiologic treatment of primary Budd-Chiari syndrome. *World J Surg* 1998;22:48-53; discussion 53-44. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9465761>.

316. Slakey DP, Klein AS, Venbrux AC, Cameron JL. Budd-Chiari syndrome: current management options. *Ann Surg* 2001;233:522-527.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11303134>.

317. Darwish Murad S, Valla DC, de Groen PC, et al. Determinants of survival and the effect of portosystemic shunting in patients with Budd-Chiari syndrome. *Hepatology* 2004;39:500-508. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/14768004>.

318. Perez-Ayuso RM, Valderrama S, Espinoza M, et al. Endoscopic band ligation versus propranolol for the primary prophylaxis of variceal bleeding in cirrhotic patients with high risk esophageal varices. *Ann Hepatol* 2010;9:15-22. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20308718>.

319. Lay CS, Tsai YT, Lee FY, et al. Endoscopic variceal ligation versus propranolol in prophylaxis of first variceal bleeding in patients with cirrhosis. *J Gastroenterol Hepatol* 2006;21:413-419. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16509867>.

320. Schepke M, Kleber G, Nurnberg D, et al. Ligation versus propranolol for the primary prophylaxis of variceal bleeding in cirrhosis. *Hepatology* 2004;40:65-72. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15239087>.

321. Psilopoulos D, Galanis P, Goulas S, et al. Endoscopic variceal ligation vs. propranolol for prevention of first variceal bleeding: a randomized controlled trial. *Eur J Gastroenterol Hepatol* 2005;17:1111-1117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16148558>.

322. Sarin SK, Wadhawan M, Agarwal SR, et al. Endoscopic variceal ligation plus propranolol versus endoscopic variceal ligation alone in primary prophylaxis of variceal bleeding. *Am J Gastroenterol* 2005;100:797-804. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15784021>.

323. Sarin SK, Gupta N, Jha SK, et al. Equal efficacy of endoscopic variceal ligation and propranolol in preventing variceal bleeding in patients with noncirrhotic portal hypertension. *Gastroenterology* 2010;139:1238-1245. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20547163>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

324. Funakoshi N, Segalas-Largey F, Duny Y, et al. Benefit of combination beta-blocker and endoscopic treatment to prevent variceal rebleeding: a meta-analysis. *World J Gastroenterol* 2010;16:5982-5992. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21157975>.

325. Goldhaber SZ. Thrombolytic therapy for patients with pulmonary embolism who are hemodynamically stable but have right ventricular dysfunction: pro. *Arch Intern Med* 2005;165:2197-2199; discussion 2204-2195. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16246980>.

326. Konstantinides S, Geibel A, Heusel G, et al. Heparin plus alteplase compared with heparin alone in patients with submassive pulmonary embolism. *N Engl J Med* 2002;347:1143-1150. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12374874>.

327. Ashton RW, Daniels CE, Ryu JH. Thrombolytic therapy in patients with submassive pulmonary embolism. *N Engl J Med* 2003;348:357-359; author reply 357-359. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12540653>.

328. Thabut G, Logeart D. Thrombolysis for pulmonary embolism in patients with right ventricular dysfunction: con. *Arch Intern Med* 2005;165:2200-2203. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16246981>.

329. Chatterjee S, Chakraborty A, Weinberg I, et al. Thrombolysis for pulmonary embolism and risk of all-cause mortality, major bleeding, and intracranial hemorrhage: a meta-analysis. *JAMA* 2014;311:2414-2421. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24938564>.

330. Dong BR, Hao Q, Yue J, et al. Thrombolytic therapy for pulmonary embolism. *Cochrane Database Syst Rev* 2009:CD004437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19588357>.

331. Tardy B, Venet C, Zeni F, et al. Short term effect of recombinant tissue plasminogen activator in patients with hemodynamically stable acute pulmonary embolism: results of a meta-analysis involving 464

patients. *Thromb Res* 2009;124:672-677. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19493561>.

332. Agnelli G, Becattini C, Kirschstein T. Thrombolysis vs heparin in the treatment of pulmonary embolism: a clinical outcome-based meta-analysis. *Arch Intern Med* 2002;162:2537-2541. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12456225>.

333. Thabut G, Thabut D, Myers RP, et al. Thrombolytic therapy of pulmonary embolism: a meta-analysis. *J Am Coll Cardiol* 2002;40:1660-1667. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12427420>.

334. Wan S, Quinlan DJ, Agnelli G, Eikelboom JW. Thrombolysis compared with heparin for the initial treatment of pulmonary embolism: a meta-analysis of the randomized controlled trials. *Circulation* 2004;110:744-749. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15262836>.

335. Aklog L, Williams CS, Byrne JG, Goldhaber SZ. Acute pulmonary embolectomy: a contemporary approach. *Circulation* 2002;105:1416-1419. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11914247>.

336. Leacche M, Unic D, Goldhaber SZ, et al. Modern surgical treatment of massive pulmonary embolism: results in 47 consecutive patients after rapid diagnosis and aggressive surgical approach. *J Thorac Cardiovasc Surg* 2005;129:1018-1023. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15867775>.

337. Thistlethwaite PA, Kemp A, Du L, et al. Outcomes of pulmonary endarterectomy for treatment of extreme thromboembolic pulmonary hypertension. *J Thorac Cardiovasc Surg* 2006;131:307-313. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16434258>.

338. Bates SM, Weitz JI. Coagulation assays. *Circulation* 2005;112:53-60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16043649>.

339. Bates SM, Weitz JI, Johnston M, et al. Use of a fixed activated partial thromboplastin time ratio to establish a therapeutic range for



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

unfractionated heparin. Arch Intern Med 2001;161:385-391. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11176764>.

340. Raschke R, Hirsh J, Guidry JR. Suboptimal monitoring and dosing of unfractionated heparin in comparative studies with low-molecular-weight heparin. Ann Intern Med 2003;138:720-723. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12729426>.

341. How to validate heparin sensitivity of the aPPT. College of American Pathologists, CAP Today; 2004. Available at: <http://www.cap.org>. Accessed February 11, 2011.

342. Nutescu EA, Spinler SA, Wittkowsky A, Dager WE. Low-molecular-weight heparins in renal impairment and obesity: available evidence and clinical practice recommendations across medical and surgical settings. Ann Pharmacother 2009;43:1064-1083. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19458109>.

343. GlaxoSmithKline. Prescribing Information: Argatroban injection. 2009. Available at: <http://www.gsksource.com/gskprm/htdocs/documents/ARGATROBAN.PDF>. Accessed September 16, 2014.

344. Boehringer Ingelheim Pharmaceuticals, Inc. Prescribing Information: PRADAXA® (dabigatran etexilate mesylate) capsules for oral use. 2012. Available at: <http://bidocs.boehringer-ingelheim.com/BIWebAccess/ViewServlet.ser?docBase=renetnt&folderPath=/Prescribing%20Information/Pis/Pradaxa/Pradaxa.pdf>. Accessed September 16, 2014.

345. van Ryn J, Stangier J, Haertter S, et al. Dabigatran etexilate--a novel, reversible, oral direct thrombin inhibitor: interpretation of coagulation assays and reversal of anticoagulant activity. Thromb Haemost 2010;103:1116-1127. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20352166>.

346. Institute for Safe Medication Practices. QuarterWatch Monitoring FDA MedWatch Reports: Signals for Dabigatran and Metoclopramide.

2012. Available at: <http://www.ismp.org/quarterwatch/pdfs/2011Q1.pdf>. Accessed September 16, 2014.

347. Harper P, Young L, Merriman E. Bleeding risk with dabigatran in the frail elderly. N Engl J Med 2012;366:864-866. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22375994>.

348. Poller L, Keown M, Ibrahim S, et al. An international multicenter randomized study of computer-assisted oral anticoagulant dosage vs. medical staff dosage. J Thromb Haemost 2008;6:935-943. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18489430>.

349. Gosselin RC, Dager WE, King JH, et al. Effect of direct thrombin inhibitors, bivalirudin, lepirudin, and argatroban, on prothrombin time and INR values. Am J Clin Pathol 2004;121:593-599. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15080313>.

350. Warkentin TE, Greinacher A, Koster A. Bivalirudin. Thromb Haemost 2008;99:830-839. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18449412>.

351. Crowther MA, Warkentin TE. Bleeding risk and the management of bleeding complications in patients undergoing anticoagulant therapy: focus on new anticoagulant agents. Blood 2008;111:4871-4879. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18309033>.

352. APP Pharmaceuticals, LLC. Prescribing information: Protamine sulfate injection, USP. 2008. Available at: http://editor.fresenius-kabi.us/Pis/Protamine_Inj_45848E_Jan_08.pdf. Accessed September 16, 2014.

353. Park KW. Protamine and protamine reactions. Int Anesthesiol Clin 2004;42:135-145. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15205644>.

354. Ansell J, Hirsh J, Hylek E, et al. Pharmacology and management of the vitamin K antagonists: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). Chest

2008;133:160S-198S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18574265>.

355. Makris M, van Veen JJ, Maclean R. Warfarin anticoagulation reversal: management of the asymptomatic and bleeding patient. *J Thromb Thrombolysis* 2010;29:171-181. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19882303>.

356. Holbrook A, Schulman S, Witt DM, et al. Evidence-based management of anticoagulant therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012;141:e152S-184S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22315259>.

357. Crowther MA, Ageno W, Garcia D, et al. Oral vitamin K versus placebo to correct excessive anticoagulation in patients receiving warfarin: a randomized trial. *Ann Intern Med* 2009;150:293-300.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19258557>.

358. Lubetsky A, Yonath H, Olchovsky D, et al. Comparison of oral vs intravenous phytonadione (vitamin K1) in patients with excessive anticoagulation: a prospective randomized controlled study. *Arch Intern Med* 2003;163:2469-2473. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/14609783>.

359. Burbury KL, Milner A, Snooks B, et al. Short-term warfarin reversal for elective surgery--using low-dose intravenous vitamin K: safe, reliable and convenient*. *Br J Haematol* 2011;154:626-634. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21751986>.

360. Sarode R, Milling TJ, Jr., Refaai MA, et al. Efficacy and safety of a 4-factor prothrombin complex concentrate in patients on vitamin K antagonists presenting with major bleeding: a randomized, plasma-controlled, phase IIIb study. *Circulation* 2013;128:1234-1243. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23935011>.

361. Warkentin TE, Crowther MA. Reversing anticoagulants both old and new. *Can J Anaesth* 2002;49:S11-25. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12557411>.

362. Vigue B. Bench-to bedside review: Optimising emergency reversal of vitamin K antagonists in severe haemorrhage - from theory to practice. *Crit Care* 2009;13:209. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19486503>.

363. Ilyas C, Beyer GM, Dutton RP, et al. Recombinant factor VIIa for warfarin-associated intracranial bleeding. *J Clin Anesth* 2008;20:276-279. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18617125>.

364. Schulman S, Bijsterveld NR. Anticoagulants and their reversal. *Transfus Med Rev* 2007;21:37-48. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17174219>.

365. Riegert-Johnson DL, Volcheck GW. The incidence of anaphylaxis following intravenous phytonadione (vitamin K1): a 5-year retrospective review. *Ann Allergy Asthma Immunol* 2002;89:400-406. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12392385>.

366. Ehrlich HJ, Henzl MJ, Gomperts ED. Safety of factor VIII inhibitor bypass activity (FEIBA): 10-year compilation of thrombotic adverse events. *Haemophilia* 2002;8:83-90. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11952842>.

367. Mayer SA, Brun NC, Begtrup K, et al. Recombinant activated factor VII for acute intracerebral hemorrhage. *N Engl J Med* 2005;352:777-785. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15728810>.

368. Bijsterveld NR, Moons AH, Boekholdt SM, et al. Ability of recombinant factor VIIa to reverse the anticoagulant effect of the pentasaccharide fondaparinux in healthy volunteers. *Circulation* 2002;106:2550-2554. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12427650>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

369. Elmer J, Wittels KA. Emergency reversal of pentasaccharide anticoagulants: a systematic review of the literature. *Transfus Med* 2012;22:108-115. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22171588>.

370. Lisman T, Bijsterveld NR, Adelmeijer J, et al. Recombinant factor VIIa reverses the in vitro and ex vivo anticoagulant and profibrinolytic effects of fondaparinux. *J Thromb Haemost* 2003;1:2368-2373.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14629471>.

371. Levi M, Bijsterveld NR, Keller TT. Recombinant factor VIIa as an antidote for anticoagulant treatment. *Semin Hematol* 2004;41:65-69.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14872424>.

372. Vavra KA, Lutz MF, Smythe MA. Recombinant factor VIIa to manage major bleeding from newer parenteral anticoagulants. *Ann Pharmacother* 2010;44:718-726. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20233918>.

373. Young G, Yonekawa KE, Nakagawa PA, et al. Recombinant activated factor VII effectively reverses the anticoagulant effects of heparin, enoxaparin, fondaparinux, argatroban, and bivalirudin ex vivo as measured using thromboelastography. *Blood Coagul Fibrinolysis* 2007;18:547-553. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17762530>.

374. Diehl KH, Romisch J, Hein B, et al. Investigation of activated prothrombin complex concentrate as potential hirudin antidote in animal models. *Haemostasis* 1995;25:182-192. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/7557657>.

375. Elg M, Carlsson S, Gustafsson D. Effect of activated prothrombin complex concentrate or recombinant factor VIIa on the bleeding time and thrombus formation during anticoagulation with a direct thrombin inhibitor. *Thromb Res* 2001;101:145-157. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11228338>.

376. Sorensen B, Ingerslev J. A direct thrombin inhibitor studied by dynamic whole blood clot formation. Haemostatic response to ex-vivo addition of recombinant factor VIIa or activated prothrombin complex concentrate. *Thromb Haemost* 2006;96:446-453. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17003921>.

377. Bhagirath VC, O'Malley L, Crowther MA. Management of bleeding complications in the anticoagulated patient. *Semin Hematol* 2011;48:285-294. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22000094>.

378. Benz K, Nauck MA, Bohler J, Fischer KG. Hemofiltration of recombinant hirudin by different hemodialyzer membranes: implications for clinical use. *Clin J Am Soc Nephrol* 2007;2:470-476. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17699453>.

379. Bove CM, Casey B, Marder VJ. DDAVP reduces bleeding during continued hirudin administration in the rabbit. *Thromb Haemost* 1996;75:471-475. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8701410>.

380. Frank RD, Farber H, Lanzmich R, et al. In vitro studies on hirudin elimination by haemofiltration: comparison of three high-flux membranes. *Nephrol Dial Transplant* 2002;17:1957-1963. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12401853>.

381. Frank RD, Farber H, Stefanidis I, et al. Hirudin elimination by hemofiltration: a comparative in vitro study of different membranes. *Kidney Int Suppl* 1999:S41-45. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10560804>.

382. Koster A, Buz S, Krabatsch T, et al. Effect of modified ultrafiltration on bivalirudin elimination and postoperative blood loss after on-pump coronary artery bypass grafting: assessment of different filtration strategies. *J Card Surg* 2008;23:655-658. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18793221>.



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

383. Willey ML, de Denus S, Spinler SA. Removal of lepirudin, a recombinant hirudin, by hemodialysis, hemofiltration, or plasmapheresis. *Pharmacotherapy* 2002;22:492-499. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11939684>.

384. sanofi-aventis US, LLC. Prescribing Information: DDAVP injection (desmopressin acetate). 2007. Available at: http://products.sanofi-aventis.us/DDAVP_IV/DDAVP_IV.pdf. Accessed September 16, 2014.

385. Mannucci PM, Bettega D, Cattaneo M. Patterns of development of tachyphylaxis in patients with haemophilia and von Willebrand disease after repeated doses of desmopressin (DDAVP). *Br J Haematol* 1992;82:87-93. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1419807>.

386. Stangier J, Rathgen K, Stahle H, Mazur D. Influence of renal impairment on the pharmacokinetics and pharmacodynamics of oral dabigatran etexilate: an open-label, parallel-group, single-centre study. *Clin Pharmacokinet* 2010;49:259-268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20214409>.

387. Warkentin TE, Margetts P, Connolly SJ, et al. Recombinant factor VIIa (rFVIIa) and hemodialysis to manage massive dabigatran-associated postcardiac surgery bleeding. *Blood* 2012;119:2172-2174. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22383791>.

388. Marlu R, Hodaj E, Paris A, et al. Effect of non-specific reversal agents on anticoagulant activity of dabigatran and rivaroxaban. A randomised crossover ex vivo study in healthy volunteers. *Thromb Haemost* 2012;108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22627883>.

389. Kaatz S, Kouides PA, Garcia DA, et al. Guidance on the emergent reversal of oral thrombin and factor Xa inhibitors. *Am J Hematol* 2012;87 Suppl 1:S141-145. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22473649>.

390. Nutescu EA, Dager WE, Kalus JS, et al. Management of bleeding and reversal strategies for oral anticoagulants: clinical practice considerations. *Am J Health Syst Pharm* 2013;70:1914-1929. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24128967>.

391. Eerenberg ES, Kamphuisen PW, Sijpkens MK, et al. Reversal of rivaroxaban and dabigatran by prothrombin complex concentrate: a randomized, placebo-controlled, crossover study in healthy subjects. *Circulation* 2011;124:1573-1579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21900088>.

392. Godier A, Miclot A, Le Bonniec B, et al. Evaluation of prothrombin complex concentrate and recombinant activated factor VII to reverse rivaroxaban in a rabbit model. *Anesthesiology* 2012;116:94-102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22042412>.

393. Ghanny S, Warkentin TE, Crowther MA. Reversing anticoagulant therapy. *Curr Drug Discov Technol* 2012;9:143-149. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22023256>.

394. Streiff MB. Long-term therapy of venous thromboembolism in cancer patients. *J Natl Compr Canc Netw* 2006;4:903-910. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17020668>.

395. Merli G, Spiro TE, Olsson CG, et al. Subcutaneous enoxaparin once or twice daily compared with intravenous unfractionated heparin for treatment of venous thromboembolic disease. *Ann Intern Med* 2001;134:191-202. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11177331>.

396. Carrier M, Le Gal G, Cho R, et al. Dose escalation of low molecular weight heparin to manage recurrent venous thromboembolic events despite systemic anticoagulation in cancer patients. *J Thromb Haemost* 2009;7:760-765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19245418>.

397. Schwarz UI, Ritchie MD, Bradford Y, et al. Genetic determinants of response to warfarin during initial anticoagulation. *N Engl J Med*



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

2008;358:999-991008. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18322281>.

398. Teefy AM, Martin JE, Kovacs MJ. Warfarin resistance due to sulfasalazine. *Ann Pharmacother* 2000;34:1265-1268. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11098339>.

399. Luk C, Wells PS, Anderson D, Kovacs MJ. Extended outpatient therapy with low molecular weight heparin for the treatment of recurrent venous thromboembolism despite warfarin therapy. *Am J Med* 2001;111:270-273. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11566456>.

400. Anderson JA, Saenko EL. Heparin resistance. *Br J Anaesth* 2002;88:467-469. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12066718>.

401. Valsami S, Asmis LM. A brief review of 50 years of perioperative thrombosis and hemostasis management. *Semin Hematol* 2013;50:79-87. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24216167>.

402. Linkins LA, Dans AL, Moores LK, et al. Treatment and prevention of heparin-induced thrombocytopenia: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012;141:e495S-530S. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22315270>.

403. Andreescu AC, Possidente C, Hsieh M, Cushman M. Evaluation of a pharmacy-based surveillance program for heparin-induced thrombocytopenia. *Pharmacotherapy* 2000;20:974-980. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10939559>.

404. Warkentin TE, Levine MN, Hirsh J, et al. Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *N Engl J Med* 1995;332:1330-1335. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7715641>.

405. Arepally GM, Ortel TL. Heparin-induced thrombocytopenia. *Annu Rev Med* 2010;61:77-90. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20059332>.

406. Greinacher A, Farnet B, Kroll H, et al. Clinical features of heparin-induced thrombocytopenia including risk factors for thrombosis. A retrospective analysis of 408 patients. *Thromb Haemost* 2005;94:132-135. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16113796>.

407. Warkentin TE, Roberts RS, Hirsh J, Kelton JG. Heparin-induced skin lesions and other unusual sequelae of the heparin-induced thrombocytopenia syndrome: a nested cohort study. *Chest* 2005;127:1857-1861. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15888871>.

408. Opatry L, Warner MN. Risk of thrombosis in patients with malignancy and heparin-induced thrombocytopenia. *Am J Hematol* 2004;76:240-244. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15224359>.

409. Prandoni P, Falanga A, Piccioli A. Cancer, thrombosis and heparin-induced thrombocytopenia. *Thromb Res* 2007;120 Suppl 2:137-140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18023709>.

410. Prandoni P, Siragusa S, Girolami B, Fabris F. The incidence of heparin-induced thrombocytopenia in medical patients treated with low-molecular-weight heparin: a prospective cohort study. *Blood* 2005;106:3049-3054. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16030191>.

411. Morris TA, Castrejon S, Devendra G, Gamst AC. No difference in risk for thrombocytopenia during treatment of pulmonary embolism and deep venous thrombosis with either low-molecular-weight heparin or unfractionated heparin: a metaanalysis. *Chest* 2007;132:1131-1139. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17646239>.

412. Levine RL, McCollum D, Hursting MJ. How frequently is venous thromboembolism in heparin-treated patients associated with heparin-



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

induced thrombocytopenia? Chest 2006;130:681-687. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16963663>.

413. Martel N, Lee J, Wells PS. Risk for heparin-induced thrombocytopenia with unfractionated and low-molecular-weight heparin thromboprophylaxis: a meta-analysis. Blood 2005;106:2710-2715. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15985543>.

414. Pouplard C, May MA, Iochmann S, et al. Antibodies to platelet factor 4-heparin after cardiopulmonary bypass in patients anticoagulated with unfractionated heparin or a low-molecular-weight heparin: clinical implications for heparin-induced thrombocytopenia. Circulation 1999;99:2530-2536. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10330384>.

415. Greinacher A, Alban S, Omer-Adam MA, et al. Heparin-induced thrombocytopenia: a stoichiometry-based model to explain the differing immunogenicities of unfractionated heparin, low-molecular-weight heparin, and fondaparinux in different clinical settings. Thromb Res 2008;122:211-220. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18262226>.

416. Greinacher A. Heparin-induced thrombocytopenia. J Thromb Haemost 2009;7 Suppl 1:9-12. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19630757>.

417. Crowther MA, Cook DJ, Albert M, et al. The 4Ts scoring system for heparin-induced thrombocytopenia in medical-surgical intensive care unit patients. J Crit Care 2010;25:287-293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20149589>.

418. Lo GK, Juhl D, Warkentin TE, et al. Evaluation of pretest clinical score (4 T's) for the diagnosis of heparin-induced thrombocytopenia in two clinical settings. J Thromb Haemost 2006;4:759-765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16634744>.

419. Warkentin TE, Aird WC, Rand JH. Platelet-endothelial interactions: sepsis, HIT, and antiphospholipid syndrome. Hematology Am Soc

Hematol Educ Program 2003:497-519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14633796>.

420. Pouplard C, Gueret P, Fouassier M, et al. Prospective evaluation of the '4Ts' score and particle gel immunoassay specific to heparin/PF4 for the diagnosis of heparin-induced thrombocytopenia. J Thromb Haemost 2007;5:1373-1379. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17362241>.

421. Cuker A, Arepally G, Crowther MA, et al. The HIT Expert Probability (HEP) Score: a novel pre-test probability model for heparin-induced thrombocytopenia based on broad expert opinion. J Thromb Haemost 2010;8:2642-2650. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20854372>.

422. Lewis BE, Wallis DE, Berkowitz SD, et al. Argatroban anticoagulant therapy in patients with heparin-induced thrombocytopenia. Circulation 2001;103:1838-1843. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11294800>.

423. Lewis BE, Wallis DE, Leya F, et al. Argatroban anticoagulation in patients with heparin-induced thrombocytopenia. Arch Intern Med 2003;163:1849-1856. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12912723>.

424. Begelman SM, Baghdasarian SB, Singh IM, et al. Argatroban anticoagulation in intensive care patients: effects of heart failure and multiple organ system failure. J Intensive Care Med 2008;23:313-320. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18701526>.

425. Kiser TH, Jung R, MacLaren R, Fish DN. Evaluation of diagnostic tests and argatroban or lepirudin therapy in patients with suspected heparin-induced thrombocytopenia. Pharmacotherapy 2005;25:1736-1745. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16305293>.

426. Reichert MG, MacGregor DA, Kincaid EH, Dolinski SY. Excessive argatroban anticoagulation for heparin-induced thrombocytopenia. Ann



NCCN Guidelines Version 1.2016

Cancer-Associated Venous Thromboembolic Disease

Pharmacother 2003;37:652-654. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12708939>.

427. The Medicines Company. Prescribing information: ANGIOMAX® (bivalirudin) for injection, for intravenous use. 2010. Available at:
http://www.accessdata.fda.gov/drugsatfda_docs/label/2010/020873s023lbl.pdf. Accessed September 16, 2014.

428. Francis JL, Drexler A, Gwyn G, Moroos R. Successful Use of Bivalirudin in the Treatment of Patients Suspected, or at Risk of, Heparin-Induced Thrombocytopenia [abstract]. Blood 2004;104:Abstract 4077. Available at:
<http://abstracts.hematologylibrary.org/cgi/content/abstract/ashmtg;104/11/4077>.

429. Kiser TH, Fish DN. Evaluation of bivalirudin treatment for heparin-induced thrombocytopenia in critically ill patients with hepatic and/or renal dysfunction. Pharmacotherapy 2006;26:452-460. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16553502>.

430. Kiser TH, Burch JC, Klem PM, Hassell KL. Safety, efficacy, and dosing requirements of bivalirudin in patients with heparin-induced thrombocytopenia. Pharmacotherapy 2008;28:1115-1124. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18752382>.

431. Wester JPJ, Leyte A, Oudemans-van Straaten HM, et al. Low-dose fondaparinux in suspected heparin-induced thrombocytopenia in the critically ill. Neth J Med 2007;65:101-108. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17387236>.

432. Warkentin TE. Fondaparinux versus direct thrombin inhibitor therapy for the management of heparin-induced thrombocytopenia (HIT)--bridging the River Coumarin. Thromb Haemost 2008;99:2-3. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18217128>.

433. Blackmer AB, Oertel MD, Valgus JM. Fondaparinux and the management of heparin-induced thrombocytopenia: the journey

continues. Ann Pharmacother 2009;43:1636-1646. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19737996>.

434. Grouzi E, Kyriakou E, Panagou I, Spiliotopoulou I. Fondaparinux for the treatment of acute heparin-induced thrombocytopenia: a single-center experience. Clin Appl Thromb Hemost 2010;16:663-667. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19825921>.

435. Lobo B, Finch C, Howard A, Minhas S. Fondaparinux for the treatment of patients with acute heparin-induced thrombocytopenia. Thromb Haemost 2008;99:208-214. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18217156>.

436. Chong BH, Chong JJH. Heparin-induced thrombocytopenia associated with fondaparinux. Clin Adv Hematol Oncol 2010;8:63-65. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20351686>.

437. Ratuapli SK, Bobba B, Zafar H. Heparin-induced thrombocytopenia in a patient treated with fondaparinux. Clin Adv Hematol Oncol 2010;8:61-65. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20351685>.

438. Rota E, Bazzan M, Fantino G. Fondaparinux-related thrombocytopenia in a previous low-molecular-weight heparin (LMWH)-induced heparin-induced thrombocytopenia (HIT). Thromb Haemost 2008;99:779-781. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18392338>.

439. Warkentin TE, Maurer BT, Aster RH. Heparin-induced thrombocytopenia associated with fondaparinux. N Engl J Med 2007;356:2653-2655. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17582083>.

440. Srinivasan AF, Rice L, Bartholomew JR, et al. Warfarin-induced skin necrosis and venous limb gangrene in the setting of heparin-induced thrombocytopenia. Arch Intern Med 2004;164:66-70. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14718324>.



441. Warkentin TE, Greinacher A, Craven S, et al. Differences in the clinically effective molar concentrations of four direct thrombin inhibitors explain their variable prothrombin time prolongation. *Thromb Haemost* 2005;94:958-964. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16363236>.

442. Arpino PA, Demirjian Z, Van Cott EM. Use of the chromogenic factor X assay to predict the international normalized ratio in patients transitioning from argatroban to warfarin. *Pharmacotherapy* 2005;25:157-164. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15767231>.

443. Austin JH, Stearns CR, Winkler AM, Paciullo CA. Use of the chromogenic factor x assay in patients transitioning from argatroban to warfarin therapy. *Pharmacotherapy* 2012;32:493-501. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22511112>.

444. Noble S. Management of venous thromboembolism in the palliative care setting. *Int J Palliat Nurs* 2007;13:574-579. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18399382>.

Discussion
update in
progress